link with their hospital to obtain experience which consultants will accept as qualifying them for appointments in the hospital practitioner grade. Greater use could be made of "clinical attachments," which can be funded under Section 63 money but which are not perhaps fully exploited at present. Greater acceptability could, however, be obtained if general practitioners could be taken on as trainee or probationer hospital practitioners. Consultants would be much more willing to utilise this type of assistance if during the first two years the sessions were to be regarded as for training as well as for patient care, and if complete security of tenure were granted only after satisfactory completion of the training period.

The creation of this new category would allow a trial period before irrevocable commitment and would, I believe, promote a much more rapid expansion of this important development in the NHS.

D INNES WILLIAMS

British Postgraduate Medical
Federation,
London WC1N 3EJ

Identifying tablets

SIR,—Every medical practitioner must have been frustrated over the identification of patients' current medication ("Little white water tablets, doctor"). Marked gratitude is felt for the easily recognisable tablets ("Black and red capsules, for my infection").

I would like to suggest that pharmaceutical companies adopt an agreed colour coding for all tablets: three rings, or dots, denoting: the type of drug (for example, major tranquilliser, antibiotic); the specific drug; the strength. A black dot could be placed at one end to denote the starting point (similar to electrical resistors). Problems would arise because of the limited number of easily discriminable colours, and because of difficulty in deciding in which group of drugs certain medications should be included (for example, diazepam—sedative or muscle relaxant?). But perhaps the division could be on the basis of chemical structure instead. None the less, with a little application a workable system could be achieved. The patient might be no better at remembering the colours, but at least if he brings several different tablets in one unlabelled bottle, a quick reference to MIMS would specify them. Furthermore, the chemists' bottles labelled with product but not strength need no longer be unhelpful.

Naturally, the tablets could still be produced in different shapes, colours, and sizes so that the pharmaceutical companies could maintain their distinction. But is there not a strong case for some such simple unification of drug identification as this?

ANTHONY KAISER

Southend General Hospital,
Westcliff-on-Sea, Essex

A new look at examination marking

SIR,—Congratulations to Drs R B Longmore and D A McRae (22-29 December, p 1640) for attempting to lighten a dreary subject. Unfortunately, the variables they use in their comparison of RAPES and CONVENT methods of marking are not independent. The null hypothesis that the algebraic sum of differences in marks obtained by one or the

other method for each of the 50 pairs of papers (in this case a value of -19), averaged out at -0.38 for each pair, does not differ significantly from zero when tested by the relevant *t* test, yields for *t* a value of 3.15 (table), which is significant at $P=0.01$.¹ 'T is

Differences between RAPES (R) and CONVENT (C) assessment of 50 papers (after Longmore and Macrae)

Value of R-C (x)	No of observations (n)	$n_i \times i$	$n_i \times i^2$
-2	6	-12	24
-1	16	-16	16
0	19	0	0
1	9	9	9
Total	50	-19	49

Average = $-19/50 = -0.38$

$s^2 = (49 - 19^2/50)/49 = 0.85$

$t = -0.38 / (0.85)^{1/2} (50)^{1/2} = 3.15$ for 49 df

true, 't is pity...² Still, since the author's data show that RAPES produced a lower mark than CONVENT in 29 out of 50 papers and a higher mark in only two, perhaps dons of limited athletic achievements should wait for further docimological studies before facing a 0.58 probability of a bloody nose.

M C THURIAUX

Project de Renforcement des Services de
Santé et de Formation de Personnel
(Project RESSFOP),
B P 457 Niamey, Niger

¹ Armitage P. *Statistical methods in medical research*. Oxford: Blackwell, 1971:116-8.

² Shakespeare W. *Hamlet, Prince of Denmark*. London: 1604:11,2,97-8.

Cutting down unnecessary x-ray examinations

SIR,—As radiologists we wish to draw the attention of our clinical colleagues to the recent doubling in price of x-ray film owing to the unprecedented rise in the price of silver over the past year. Now more than ever it behoves those in both hospital and general practice before requesting any x-ray examination to ask themselves the vital question: "Will the result of this investigation affect my management of this patient?"

Cutting down on the number of unnecessary examinations will not only preserve scarce resources but also have the even more desirable effect of reducing the overall radiation dosage to the population as a whole.

F W WRIGHT

Chairman, BMA Radiologists Group

MYER GOLDMAN

Vice-chairman, BMA Radiologists Group

Fazakerley Hospital,
Liverpool L9 7AL

Medically qualified preclinical academics

SIR,—I write in the strongest possible support of Dr R R Sturrock's comments (2 February, p 335) on medically qualified preclinical academics. Two points, I think, need to be made—the first to Dr Sturrock and the second to the general medical public.

Firstly, some preclinical academics would undoubtedly find benefit both to their teaching and to their research by clinical involvement; with a research interest in sensory processes

and a teaching commitment to basic neurology, I am one such. But obviously a great deal depends on what kind of clinical involvement—it should not be anything that's going, or someone's unwanted general sessions. However, medical anatomy in particular is (for most doctors) about living people—and therefore any professional commitment by its teachers to examination of the quick by palpation, percussion, and auscultation would add immeasurably to dissection of the dead.

Secondly, and perhaps more importantly, I fail to understand why the question of proper (that is, NHS-equivalent) payment to medically qualified preclinical academics is such an emotive issue. It is often the same people who demand on the one hand that preclinical teachers (or at least a substantial proportion of them) should be medically qualified, and on the other that they should not receive the same salaries as teachers in "clinical" departments (for example, epidemiology?) without clinical justification—that is, participation. Yet no one animadvert against the payment of clinical (or above) salaries to area and regional and other administrative medical officers, or to the medically qualified secretariat of the BMA, the Medical Research Council, and the DHSS; and they are right. The fact remains, Sir, that medically qualified preclinical academics and full-time research workers are the only group of doctors in the country who are not paid in accordance with their professional qualifications. To repeat the unanswered question which you allowed me to put through your columns last year, what do the great and muscular majority of the profession intend to do about us 300?

DAVID BOWSHER

Department of Anatomy,
Liverpool L69 3BX

MDs or PhDs in medical research?

SIR,—With regard to the leading article (2 February, p 274) on MDs or PhDs in medical research, this article indicated that there was a vast difference necessarily between people with PhDs and people with MDs. I think the point should be made that it is not necessarily the degree which is obtained that matters but the subject matter for which the degree was given.

I did some medical research leading to a doctoral thesis in 1977. As I did this under the wing of the University of East Anglia, I was awarded a PhD for this work. If I had done it under the auspices perhaps of London University, and this had passed it, I would have received an MD, not a PhD. The difference therefore is how the degree is awarded, not necessarily the subject matter or even whether the man is a clinician or not. This is not made clear in your article and perhaps the difference between the MD and PhD could be demonstrated only by studying the work for which the degrees are granted.

ADRIAN CARO

Dereham, Norfolk N19 2EN

Revised consultant contract

SIR,—As a longstanding member of the BMA, it seems to me that the letter from Mr David Bolt (19 January, p 202) does not clear up the confusion to which he alludes about the nature of the changes to the consultant contract. In the paragraph headed "Consultants with