and torture. (4) Establishing a network of prominent medical experts in the world who could influence their colleagues in countries acting in breach of the Helsinki agreement to stand firm on the side of human rights.

We would welcome any volunteers sympathising with our aims as new members of Amnesty International's medical group. If, however, any sympathisers feel that they cannot fully participate, on account of their present commitments, we would be grateful for help with an occasional case where the authority of a particular specialist might be of value. We would like to extend this invitation to members of the dental and nursing professions.

Please write to: London Region Medical Group, British Section, Amnesty International, 8-14 Southampton Street, London WC2E 7HF.

J POLLERT

Medical Adviser to London Office of the
International Secretariat

ELIZABETH M GORDON

Co-ordinator, London Region
Medical Group

Amnesty International, British Section,
London WC2E 7HF

Medical audit and clinical competence

SIR,—At the recent meeting of the National Association of Clinical Tutors a straw poll was taken on the question of whether clinical tutors should act as "catalysts" in promoting educational opportunities in their hospitals which could be construed as instituting medical "audit." During the discussion it was accepted that some form of "audit" of a doctor's continuing competence to practise was desirable and that in the first instance this should probably take the form of encouraging clinical case discussions. In the preceding debate the word audit seemed to be used to denote peer review, quality of patient care, cost effectiveness, and clinical competence as well as that instilled professionalism of doctors which has *always* involved a willingness to learn and an anxiety to ensure that personal standards are at least average.

One of the most important ways by which this is achieved is co-operative discussion with colleagues about the management of patients. To refer to this process, which should be absolutely inherent in a doctor's professionalism, by a catchy, new, and misleading name and to suggest that this process may partially placate the supposed demands, legitimate or not, for some form of definable assessment of a doctor's continuing competence is at the best deluding and at the worst deceiving. To proclaim the growth of clinical meetings as part of the new process of "audit" and imply that this is in some way praiseworthy is surely wrong. So much should this process be assumed that we should positively refrain from proclaiming that it may be lacking in some institutions. By agreeing to institute any form of audit we implicitly acknowledge that the need for it exists, whereas we should perhaps be vigorously denying its need rather than appeasing by a process which should be an unquestioned part of continuing self-criticism.

We should realise that audit as generally understood denotes the process of identifying individuals whose standards of practice fall below the minimum acceptable to the majority. It has little or nothing to do with promotion of the highest standards of competence, which

should be the main aim of medical educators, and little to do with the clinical competence of the majority of doctors.

I J T DAVIES

Postgraduate Medical Centre,
Raigmore Hospital,
Inverness IV2 3UJ

Multidisciplinary teams

SIR,—One might take issue with several points in the article by Drs James Appleyard and J G Maden on multidisciplinary teams (17 November, p 1305) and your leading article on the same topic. However, I should like to draw attention to two specific areas.

In the first place, the concept of "a team" is used extremely loosely, to such an extent that it becomes virtually meaningless. In the first example cited by Drs Appleyard and Maden, for instance, "the team" consists of no more than two professionals—who do not necessarily work together on a regular basis but who from time to time must reach agreement on a clinical matter. My edition of the Oxford Dictionary defines a team as "two or more beasts of burden harnessed together; set of persons working together." It is the idea of a group of people bounded by a common purpose who regularly come together to work and to fulfil their task that is central to the concept of a team.

In the second place, it is stated that clinical management by consensus does not work; that "it is impossible for a team to carry out any form of treatment or social work with an individual patient"; and that a multidisciplinary team cannot be "the hand that gives the pill." I have in fact recently reviewed the first eight months' work of a pilot study that has been carried out in this department with a small group of chronic psychiatric patients which directly contradicts these statements. In the course of this study, each of these patients was seen jointly, on an outpatient basis, by a social worker, a community nurse, and myself on from one to five occasions. In addition, these patients were seen individually by at least one of us. Chronic psychiatric outpatients are often seen by workers of different disciplines at different times for overlapping reasons and an important advantage of this joint approach was that management could be discussed concurrently by all the workers concerned, often together with the patient. The managements of these patients' neurotic and social problems were enhanced by this team method, while a recent spell of inpatient care and poor compliance with treatment were felt to be indications for individual management by a psychiatrist.

In general, the multidisciplinary team can enhance the clinical work of individuals, but can probably never replace it. Like any other clinical method, the team approach to management has its indications and its contraindications. Like other methods, it is also liable to abuse. In particular, the practitioner—of whatever discipline—can hide behind the team in an attempt to avoid individual responsibility. The team may be used to "blur roles," as a defence against envy and competition between disciplines, rather than as a means of sharing professional skills. At worst, people working in teams may become preoccupied with internal issues rather than the needs of the community they are meant to serve. These abuses must obviously be guarded against—ensuring that only part of an individual's clinical work is

carried out as a member of a team is one way of doing this—but are no reason to reject multidisciplinary teams out of hand.

There should be careful study and evaluation of the contribution that teamwork can make to clinical practice, aided by clear thinking about the concepts and issues involved. Neither uncritical acceptance nor emotive rejection of teams, or the staunch defence of the rights of doctors, is a valid substitute. Furthermore, working in a multidisciplinary team is a skill which, like other skills, requires training and practice in order to be applied effectively.

T PASTOR

Department of Psychiatry,
St Mary's Hospital Medical School,
St Mary's Hospital (Harrow Road),
London W9 3RL

SIR,—The highly topical article on multidisciplinary teams by Drs James Appleyard and J G Maden (17 November, p 1305) raises important issues. As a consultant psychiatrist with responsibility for the Wilton Unit, an outpatient group psychotherapy unit which is staffed by a multidisciplinary team consisting of four nurses, a social worker, a clinical psychologist, a lay group analyst, two part-time occupational therapists, and a sessional drama therapist (voluntary), I would like to share my own experience of the multidisciplinary team.

We take up to 18 severely disturbed neurotic patients for a time-limited period of 12 weeks in a daily group psychotherapy programme. I spend three sessions weekly in this unit and it is clearly emphasised to both patients and staff that I am clinically responsible for all patients admitted. Signing of sick notes and prescribing are my total responsibility. We do, however, aim to produce an agreed policy in the treatment programme; but should this fail to be agreed there is an overt understanding that the final decision is mine.

At the weekly policy meeting, which has a rotating chairman, policy for the unit is hammered out. The full agenda is submitted by members of staff, and this meeting has top priority in the work programme. It covers matters concerned with the weekly treatment programme, admission, discharge, and the role of the staff members in their differing disciplines; and their specialist points of view are encouraged. Only once in six years has there been an occasion when I overruled a dissident sister. At a further weekly meeting all patients are individually reviewed and their progress monitored. All referrals are accepted initially by me, but before full admission takes place assessment of each patient's suitability for treatment is fully discussed by the team. Some patients are "selected out" through an agreed policy made through the policy meeting.

In addition to these team meetings there is a weekly staff experimental group, led by a consultant colleague, where staff feelings as opposed to opinions are ventilated; these frequently concern anxiety and loss of role identity through the team approach. In this group we are also brought into touch with the power and reality of outside "masters" in our different disciplines, and are reminded that we are not an isolated treatment unit, but part of a wider and infinitely complex series of other systems in hierarchical order.

The National Health Service cannot afford the multidisciplinary team where the consultant abdicates ultimate responsibility for patient care in favour of a multidisciplinary