dence was 1% and 3% in Copenhagen¹ during 1970-2, 1.4% in London² in 1977, and <1% in the Helsinki³ area during the period 1972 to 1977, all studies comprising both outpatients and inpatients.

At the 11th International Congress of Chemotherapy, Boston, in October 1979, P Toivanen and K Dornbusch presented the results of a comparative study of urinary tract pathogens from Stockholm and Turku. In Sweden, where trimethoprim was not used alone, the incidence of resistance during 1977-8 was just under 1%, unchanged throughout the period; while in Turku it was 15% (continuing to increase during the period of study).

For the moment we are again monitoring the sensitivity patterns of urinary tract bacteria at a Copenhagen city hospital. In 1979 we found 3.2% trimethoprim-resistant *E coli* strains—that is, the incidence was unchanged over a period of 10 years in an area where plain trimethoprim has been on the market for one and a half years. Furthermore, we know of a comprehensive series from Aarhus in Denmark (J K Møller *et al.*, personal communication) during 1976-7, in which a sulphonamide-trimethoprim combination, but not trimethoprim alone, has been used since 1972. Among *E coli* strains 2.5% were trimethoprim resistant.

In both reports the authors consider that their findings, together with the rare occurrence of side effects and the convenient dosage scheme, confirm the usefulness of plain trimethoprim for urinary tract infections. In our opinion, all these results demonstrate a coincidence between the use of plain trimethoprim and the development of an *E coli* population resistant to this drug and perhaps also other drugs used in urinary tract infections (sulphonamides, nitrofurantoin, and ampicillin).

The possibility of a causal correlation could justify restriction in the use of plain trimethoprim for long-term treatment until further knowledge has accumulated.

HANS ERIK BUSK
BENT KORNER

Department of Clinical Microbiology,
Bispebjerg Hospital,
DK 2400 Copenhagen NV, Denmark

¹ Fruensgaard K, Korner B. *Chemotherapy* 1974;20:97-101.

² Brumfitt W, Hamilton-Miller JMT, Grey D. *Lancet* 1977;ii:926.

³ Kasanen A, Anttila M, Elfving R, *et al.* *Ann Clin Res* 1978;10,suppl 22:1-39.

Examination of abortuses

SIR,—May I support the recommendations put forward by Dr D I Rushton (15 March, p 767) that all maternal deaths should be adequately necropsied by a recognised panel of pathologists with specific experience in this field?

Such an arrangement is long overdue and I would suggest that the same regional pathology teams should also encompass one other specialised form of necropsy within their remit—namely, the examination of all abortions undertaken on medical grounds. Like maternal deaths, such abortuses are relatively few in number and many are subject to only cursory examination. Not only is it essential to verify the diagnosis for which therapeutic abortion has been performed (usually craniospinal dysraphia or chromosomal abnormality) in order to monitor the accuracy of prenatal diagnostic techniques, but the parents and the

public have a right to an independent reassurance of the propriety of medical practice. Indeed, on these grounds there is a strong case for making all therapeutic abortions notifiable to the coroner in the way which has long been a routine for other deaths occurring in the hospital operating theatre.

A J BARSON

Department of Pathology,
St Mary's Hospital,
Manchester M13 0JH

Mebendazole and hydatid disease

SIR,—We have followed with great interest the correspondence on mebendazole and hydatid disease. Although there is some evidence from Dr Anthony Bryceson (15 March, p 796) and others that with enormous doses by mouth mebendazole can be found in the plasma in low concentrations, it is not reasonable to treat this disease, which requires significant blood levels and indeed cyst fluid levels, orally with a drug developed specifically to be poorly absorbed from the gastrointestinal tract. Mebendazole is an excellent broad-spectrum anthelmintic capable of dealing with a wide range of intestinal parasites; and in this situation, of course, systemic absorption is not required or desirable.

The clinical evidence available to date is not encouraging and confirms that in established hydatid disease no dosage regimen so far devised for giving mebendazole by mouth offers more than an outside chance of causing cysts to shrink and scolices to die. The product licence for mebendazole (Vermox) does not permit us to recommend any dose above 200 mg per day for three days; and hydatid disease is not an approved indication, which accounts for the high apparent cost of Mr D R Osbourne's "full course of mebendazole" (19 January, p 183).

It is not our intention to pursue further studies with mebendazole in hydatid disease—we intend to look instead for alternative compounds and routes of administration.

A L MACNAIR

Janssen Pharmaceutical Limited,
Marlow, Bucks SL7 1ET

Lumbar puncture

SIR,—The value of the otherwise excellent series "Procedures in Practice" is marred by the recent description of lumbar puncture by Drs C Clough and J M S Pearce (2 February, p 297). The article nicely brings us up to date on the indications and contraindications for the technique. The account is, however, too complacent in its acceptance of unwarranted complications.

It is not surprising that 25% of the authors' patients experience a postspinal headache (and in half of those it is described as severe) when an 18-gauge needle is used. Headache is related to needle gauge and, presumably, to the size of the hole in the dura.¹ A 25-gauge needle should be adequate for diagnostic aspiration of cerebrospinal fluid and a 22-gauge needle should suffice otherwise. Disposable needles of these sizes are readily available in anaesthesia departments and have the added virtue of being reliably sterile and sharp, unlike the repeatedly used needles favoured by physicians.

The rare postspinal headache following the use of smaller needles can now be effectively

and safely treated by the use of an epidural blood patch.²⁻⁴ In this technique 10 ml of the patient's blood is taken aseptically and instilled into the epidural space, where it forms a gelatinous plug to seal the cerebrospinal fluid leak.

The authors may be criticised also for failing to stress the importance of a no-touch technique and the value of the patient's sitting in an upright position, particularly if he or she is obese.

It is no wonder that so many patients dread a lumbar puncture, whether for diagnostic or for therapeutic purposes. The time has come for the quality of the technique used on the wards to match that used in the operating rooms.

COLIN E BLOGG

Baylor University Medical
Center,
Dallas, Texas 75246,
USA

¹ Dripps RD, Vandam LD. *JAMA* 1951;147:1118-21.

² Gormley JB. *Anesthesiology* 1960;21:565-6.

³ DiGiovanni AJ, Dunbar BS. *Anesth Analg Curr Res* 1970;49:268-71.

⁴ DiGiovanni AJ, Galbert MW, Wahle WM. *Anesth Analg Curr Res* 1971;51:226-32.

SIR,—May we reply to the points raised by your correspondents about our paper on lumbar puncture (2 February, p 297)? Firstly, we would emphasise that the paper was written primarily for junior doctors in training, many of whom do not share all the facilities of specialised neurological and neurosurgical units; this may modify a counsel of perfection. For example, although we agree with Dr D C Thrush (23 February, p 564) that Queckenstedt's test is not necessary, this is based on the assumption that in any case of suspected spinal cord compression myelography will be performed, making manometry unnecessary. There are, however, doubtful or borderline cases which in a neurological centre would end up with myelography, but which in a smaller hospital may be investigated by a general physician by lumbar puncture; in these circumstances (usually cases of "probable multiple sclerosis" where some diagnostic confirmation is sought) manometry and Queckenstedt's test may provide additional information which could be the first hint of spinal block and the need for more specialised investigation.

We are not unduly impressed with Dr A C Young's observations (15 March, p 796) that manometers cost the Salford authority £660 in a year: this must be infinitesimal in comparison with the multitude of waste which occurs daily in a hospital. Dr Young has a neurological unit and therefore manometry should seldom be necessary, but others are less well equipped.

We agree with Mr J R Gibbs (23 February, p 564) that the correct alignment of the patient is vital, but this was plainly described in our paper—as Dr W H H Calwell points out (p 565). Precise details vary, and a firm mattress with the patient placed right up to the edge proves as satisfactory as a trolley, though the latter is easier on the operator's back.

Dr C E Blogg accuses us of being too complacent in "accepting unwarranted complications"—a nice self-contradiction. The figure of 25% we cite for headache after lumbar puncture (only half of which were severe) is based on actual observations of unselected ward patients,¹ and is not dependent on an unproved—albeit reasonable speculation—that