lationship of solution pH to biological activity. All the available data indicate that alkaline glutaraldehydes, including Cidex Long-Life, are significantly more active against micro-organisms than acid glutaraldehydes, including Sonacide.²⁻⁴ These data, coupled with the fact that Cidex Long-Life is only slightly less chemically stable than Sonacide, enable us to state categorically that Sonacide does not have a longer use life than Cidex Long-Life when compared under identical conditions of use. In addition, Cidex Long-Life has greatly superior anti-corrosion properties throughout an equivalent use period.

A C MAIR

Arbrook,
Livingston, Lothian

¹ Stonehill AA, Krop S, Borick PM. *Am J Hosp Pharm* 1963;20:458-65.

² Miner NA, McDowell JW, Willcockson GW, Bruckner NJ, Stark RL, Whitmore J. *Am J Hosp Pharm* 1977;34:376-82.

³ Collins FM, Montalbano V. *J Clin Microbiol* 1976;4:408-12.

⁴ King JA, Woodside W, McGucken PV. *J Pharm Sci* 1974;63:803-5.

Inflammatory bowel disease in relatives of patients with Crohn's disease

SIR,—I was interested in the report of Dr J F Mayberry and others (12 January, p 84). Their findings that 9% of patients with Crohn's disease in Cardiff had a first-degree relative with inflammatory bowel disease is similar to the 15% previously reported in 186 patients with Crohn's disease in Birmingham.¹

Their statement that only two married couples with Crohn's disease have been reported is erroneous; I am aware of a further such report² and there may well be others. As has been suggested,² the children of these couples should form the basis of a prospective genetic study.

J F FIELDING

Charitable Infirmary,
Dublin 1

¹ Fielding JF. MD thesis, National University of Ireland, 1970.

² Fielding, JF. In: Badenoch J, Brooke BN, eds. *Recent Advances in Gastroenterology*. Edinburgh: Churchill Livingstone, 1972:276.

Polycythaemia vera and central sleep apnoea

SIR,—We were interested to see the report on polycythaemia rubra vera and central sleep apnoea from Dr John F Neil and others (5 January, p 19).

We are currently studying nocturnal hypoxia in patients with chronically obstructed airways. In one such patient marked transient nocturnal hypoxia was reversed following a phlebotomy which reduced the packed cell volume from 0.6 to 0.5. In our case, however, there was neither central nor obstructive apnoea. We think that in some patients with chronic respiratory disease who have secondary polycythaemia and transient sleep hypoxia the high packed cell volume may be the cause of the extra sleep hypoxia rather than vice versa, as has also been proposed.¹ Whether this occurs through hypoventilation or increased ventilation-perfusion mismatch is not clear.

An alternative explanation for the finding of Dr Neil and his colleagues that the sleep apnoea only occurred during non-rapid eye movement (non-REM) sleep is suggested by

Phillipson.² He found in dogs that classical chemical control of ventilation (brain stem) did not seem to regulate respiration during phasic REM sleep and hence interference with brain stem blood flow might not be so critical during this period.

J R STRADLING
DONALD J LANE

Chest Clinic,
Churchill Hospital,
Oxford OX3 7LJ

¹ Douglas NJ, Calverley PMA, Leggett RJE, Brash HM. *Lancet* 1979;ii:1-4.

² Phillipson EA. *Am Rev Resp Dis* 1978;118:909-39.

Day-bed units

SIR,—We have followed with interest the correspondence arising from our papers on day surgery and the work of a day-bed unit (22 September, pp 712 and 714) and should like to thank contributors for their comments.

Mr C A C Clyne and Mr C W Jamieson (27 October, p 1075) appear to believe that the place of outpatient surgery for varicose veins is very limited. They are correct in noting in our data a small relative decline in major day surgery during the last two years, but their interpretation of the cause is not correct. There are several reasons. The most important have been industrial action by hospital and ambulance personnel during the past two years; the opening of a five-day ward, which allowed us to work through a waiting list backlog of short-stay patients; and the special efforts made to enhance the through-put as part of a clinical trial¹ in 1974-5 (hence the peak in those years)—but not any diminished enthusiasm for the day-care system itself. Overall, in fact, the proportion of our varicose vein operations performed on an outpatient basis has steadily increased over the last decade. In the five years 1969-73 53% of our varicose vein surgery was done on an outpatient basis. In the years 1974-8 the figure rose to 70%. It is likely to level out at somewhere between these two figures.

The clinical trial referred to above produced "consumer" responses which showed a higher proportion of responses favourable to day care than to inpatient care in an acute ward or a convalescent hospital,² and the responses were thus different from those reported by Mr Clyne and M Jamieson. We certainly do not accept the suggestion that day surgery "will often provide inadequate treatment." That surely has more to do with the quality of the surgery than the system of care employed. Operations in our trial all involved multiple incisions; roughly 50% included stripping of the upper half of the long saphenous vein (we do not strip the distal long saphenous vein); and the operations lasted on average about one and a half hours.

The valuable contribution to the literature made by Mr T H Berrill (3 November, p 1146) was acknowledged in our second paper. Any comparison between our experience and other published series was intended to relate not particularly to numbers of patients but to the scale of procedures employed, with the object of encouraging others to widen the scope of this type of care. Mr Berrill can be reassured that no persuasion was necessary to encourage colleagues to use the day-bed unit. On the contrary, the problem has been to provide sufficient accommodation for all the enthusiastic claims. He is probably quite

correct in recommending day care for ENT procedures. We cannot comment since that specialty is not represented at the Western General Hospital, nor can we usefully answer Mr Berrill's challenge on the merits of combined medical and surgical day care versus a surgical unit alone since we do not have experience of the latter. There is probably a good place for each type of unit.

Mr H D S Vellacott (17 November, p 1293) and Dr Elizabeth Spalding (15 December, p 1586) are both enthusiastically in favour of day care and correctly point to its unexplored potential. We agree with Mr Vellacott that it is difficult to run day care on an ordinary surgical ward on anything except a minor scale. Specific day care accommodation and staffing are required. Elizabeth Spalding seems to blur the distinction between day care and short stay but there are important practical differences. We were a little puzzled by her reference to doctors removing sutures. District nurses remove most of our sutures except from patients who are too old or too ill, or whose social circumstances prevent discharge from hospital before sutures are due to be removed. Dr Spalding describes an altogether admirable approach to post-operative care in the home, but we suspect that she may not realise that in most acute surgical wards nowadays it would be difficult for her to find patients who could appropriately be sent home earlier to the type of care she describes. In our experience the average acute ward largely contains very ill or very dependent patients. Most of the patients suitable for day care and early discharge are not in acute wards—they are on waiting lists.

IAN A GOULBOURNE
J RAINEY
C V RUCKLEY

Royal Infirmary,
Edinburgh EH3 9YW

¹ Ruckley CV, Cuthbertson C, Fenwick N, Prescott RJ, Garraway WM. *Br J Surg* 1978;65:456-9.

² Garraway WM, Cuthbertson C, Fenwick N, Ruckley CV, Prescott RJ. *J Epidem Comm Health* 1978; 32:219-21.

Chiropractic and the NHS

SIR,—Would you allow me to use your correspondence column to comment on the report "Chiropractic in the NHS." (5 January, p 54)?

Manipulation is already available within the NHS from an ever-growing number of specially trained physiotherapists. Over 200 such expert practitioners in the NHS, industry, and private practice belong to the Manipulation Association of Chartered Physiotherapists, a "specific interest group" of the Chartered Society of Physiotherapists. This body was set up over 10 years ago and is a steadily growing national group dedicated to the extension of postregistration manipulation training for chartered State-registered physiotherapists. Our president is Mr Philip Newman, FRCS. Regular long courses are held in all aspects of manipulative therapy and teachers include internationally recognised medical experts.

There is nothing that a chiropractor could offer a patient with back pain that a suitably trained physiotherapist does not; indeed, we believe the physiotherapist has far more to offer—and in a proper ethical relationship with the referring medical practitioner. The idea

suggested in the report that chiropractors would work "with help from physiotherapists" is a serious misconception and we would strenuously resist any attempt to erode our direct referral relationship with consultants.

P E WELLS

Chairman, Manipulation Association of
Chartered Physiotherapists

Physiotherapy Department,
St Stephen's Hospital,
London SW10 9TH

St Joseph's Hospice

SIR,—I have recently retired from general practice in Hackney and have considerable experience of the work of St Joseph's Hospice and, like Dr B E R Symonds (19 January, p 184), find the strictures of AS (15 December, p 1579) unjustified.

I have never known a patient with terminal illness to be refused admission and at no time have I been asked the religion of my patient. I was always made welcome when visiting my patients, who usually commented on the kindness of the staff. Naturally, the environment is that of a Roman Catholic institution with nuns in attendance.

The McMillan domiciliary service attached to the hospice is in my experience unique. It provides expert medical and nursing care 24 hours a day for patients wishing to die at home.

I am glad of this opportunity to thank the staff of St Joseph's Hospice for their invaluable help to many of my terminal patients. I should like to add that I am a Jew and an agnostic.

DAVID BIELENKY

London NW6 5JH

Cimetidine for pruritus related to systemic disorders

SIR,—I was interested to read the report on the use of cimetidine for pruritus in Hodgkin's Disease (19 January, p 151). However, I feel that some comments are warranted.

It is suggested in the introductory paragraph that cimetidine has been of value in the itching of cholestasis, chronic renal failure, and the myeloproliferative disorders. However, the supporting references do not bear this out. The evidence for its usefulness in the itching of polycythaemia rubra vera is limited to two case reports.^{1,2} On the basis of this, a small uncontrolled trial was carried out on 12 patients with itching associated with polycythaemia; there was no improvement in the itching in any of the patients.² Similarly, there have been only two case reports of the usefulness of cimetidine in the itching associated with cholestasis. The same authors then carried out a small double-blind controlled trial involving six patients with itching related to cholestasis and showed that cimetidine was helpful in only one patient, while two patients responded to placebo.³ There has been one uncontrolled trial on the use of cimetidine in the itching related to chronic renal failure in 10 patients, in which there was no improvement in the itching in any of the patients studied.⁴

Evidence therefore of the usefulness of cimetidine in relieving itching related to cholestasis, chronic renal failure, and polycythaemia rubra vera is limited to a few isolated case reports, and its effectiveness has not been substantiated in the few clinical trials

carried out. The further reporting of the use of cimetidine in pruritus related to Hodgkin's disease in four patients may be of some interest.

It is, however, essential, if any valid conclusions are to be made, that further investigations into the use of cimetidine in pruritus related to systemic disorders should be based on large, properly controlled trials and not on uncontrolled case reports.

D L ROBERTS

Department of Dermatology,
North Staffordshire Hospital Centre,
Stoke-on-Trent ST4 7PA

¹ Easton P, Galbraith PR. *N Engl J Med* 1978;299:1134.

² Scott GL, Norton RJ. *N Engl J Med* 1979;300:434.

³ Harrison AR, Littenberg G, Goldstein L, Kaplowitz N. *N Engl J Med* 1979;300:433-4.

⁴ Zappacosta AR, Howss D. *N Engl J Med* 1979;300:1280.

Emergency admission arrangements in hospital with inadequate resources

SIR—The recent correspondence in the lay and medical press concerning compensation for alleged negligence by doctors reminds one of the increasing problems whereby doctors, generally in the junior grades, are wrongly accused of negligence (sometimes partly more difficult to defend through inadequate records made by them). The defence organisations and the health authorities, if negligence is proved, often both contribute to the damages. In many of these cases negligence is not present but there are "negligent conditions" present.

(1) "*Musical beds*" with patients—Patients are often moved from ward to ward, sometimes late in the night, in order to attempt to obtain a high-dependency bed for an emergency admission. It is quite likely that the patient concerned suffers not only psychologically but physically from being moved to a low-dependency area if his medical condition barely warrants this. The legal responsibility for this move could rest unfairly with the doctor.

(2) "*Bussing*" of patients—This occurs when patients arrive in the casualty department and the junior medical staff are "persuaded" by non-medical Health Service workers to move the patients to another hospital. It has happened that patients have died or deteriorated during transit and actions have been brought against junior doctors as a result.

(3) "*Long waiting periods in casualty*"—Patients can wait a very long period in casualty because the doctors know that a particularly high-dependency bed is needed for a patient but the administration cannot find a suitable available one. Even with consultant support—which, of course, should be requested—the doctor may be left with the alternatives of keeping a very ill patient in casualty, risking transporting him to another hospital, or admitting him to a quite unsuitable bed. If the consultant is not in the hospital he cannot help with regard to immediate clinical decision making. Does a doctor or consultant have any authority to put up an extra bed in these circumstances? Obviously it is unfortunate for extra beds to be put up in any ward but emergency conditions do sometimes arise where this is essential. If the request to put up an extra bed is countermanded by other than medical staff the doctor must write in the notes the name of the person who refused this request.

On any occasion when allegations of negligence in these circumstances are made against medical practitioners they should be firmly

resisted by the defence organisations as the suffering caused to the patient is not due to lack of medical skill or care.

Family doctors have recently been worried concerning their medicolegal responsibilities when they are unable to obtain the admission of their patient to a suitable hospital bed. I gather that the BMA General Medical Services Committee is looking into the matter in relation both to the common law angle and to the family doctor's National Health Service terms of service. But I hope that it is unnecessary to say that the suggestions are not only aimed at protecting the medical profession from a legal standpoint. What perhaps is more important is to try to prevent the quite unnecessary strain which junior doctors suffer through having to work in completely inadequate conditions to the detriment of their patients. Administrators and politicians sometimes appear to think that the admission to any hospital bed resolves the situation; unless adequate resources for diagnosis and treatment are available, however, admission itself is of little use and merely gives a false sense of security to the patient and his relatives.

There comes a time when the medical profession has got to raise a voice against the continuing lowering of standards in the Health Service.

J ANDREWS

West Middlesex Hospital,
Isleworth, Middx TW7 6AF

The threshold of urgency

SIR,—The current dispute about out-of-hours work by the medical laboratory scientific officers is regretted and one must hope that a solution will soon be found. It is essential to keep in sight the fact that some 20 years or so ago out-of-hours work was done only for the seriously ill where life or limb could be saved, but over the years the threshold has become very much lowered. The situation now is that the junior medical staff even at preregistration level ask for a host of tests that would be better done during the routine working week. If the clinical consultant would influence the choice of test in relation to serious need a great deal of time and money would be saved. Furthermore, one must accept that work done during the routine day with all the major apparatus in use is the most accurate.

If there is a case for an extended working day that really is a separate issue.

C A K BIRD

Pathology Department,
Macclesfield Hospital,
Macclesfield, Cheshire SK10 3BL

Patients first

SIR,—Having just spent three frustrating and rewarding years as a district management team consultant representative I have just read the Government's Green Paper *Patients First* with great interest.

It is very clear to me that the district management team in its present form must be the executive body of the new health districts. Its composition is exactly right to reflect the needs of hospital and community. The sensible unification of general practice, community medicine, and hospital practice brought about by the 1974 reorganisation must continue, each