clinicians to examine patients who may be under the care of other doctors.

There is an additional point which I would like to bring forward. The loss of the medical officer of health was a great one. There is no doubt that he could be very helpful to general practitioners, and that general practitioners with an epidemiological background could often alert him to what was going on in their area and allow controlled measures to be applied.

I would like to support the study of epidemiology in clinical practice as a very useful activity.

E J C KENDALL

Epsom, Surrey KT19 8AW

Measles again

SIR,—Dr Hillas Smith (15 March, p 766) believes that the current level of measles vaccination in the UK will result in an increasing incidence of measles in older age groups and this is certainly possible. He quotes as one reason for this the American experience of "loss of vaccine-induced immunity in the group aged over 15," but there is no evidence that this is happening here. The Medical Research Council trial of further attenuated (Schwartz strain) live measles vaccine given to children aged 10 months to 2 years¹ has now completed 15 years of follow-up and the attack rate in the vaccinated group shows no signs of a loss of immunity (see accompanying figure). This was confirmed by estimations of

Medical Research Council investigation of measles vaccines: annual attack rate in children 1964-79.

measles antibody (by haemagglutination inhibition) 10 and 15 years after vaccination. It seems unlikely on this evidence that there will be a sudden loss of vaccine-induced immunity after 15 years. Measles is more likely to occur in adults who have neither been vaccinated nor contracted measles, and in the small proportion in whom vaccine has failed (4-8%).^{1,2}

The UK policy of measles vaccination since May 1968 (set out in CMO 1/68) has been to give one injection of live attenuated measles vaccine in the second year of life. Killed vaccine followed by live vaccine was used in the UK only for comparatively small numbers in trials up to June 1968, and multiple doses of killed vaccine have never been used. The American experience of measles following these vaccine combinations is therefore unlikely to be duplicated here.

Measles vaccine was introduced because of the risk of severe complications of measles.³ These have not changed⁴ and encephalitis, the

most serious, can follow an unremarkable attack of measles at any age. If the policy of selective immunisation were to be adopted, both the incidence of measles—and inevitably the complications—would increase again. Immunisation of "groups at risk" is notoriously difficult both for the decision as to who is at risk and because of the problem of how to reach them.

A more realistic approach to this problem would be to improve on the vaccination acceptance of 50% in the second year of life by offering vaccine at school entry to those who have had neither measles nor vaccine by that age. In the United States this policy has effectively reduced the incidence of measles in school-aged children in the States where it is enforced.⁵

CHRISTINE MILLER

Secretary, Measles Vaccines Subcommittee, Medical Research Council Committee on Development of Vaccines and Immunisation Procedures

Public Health Laboratory Service
Epidemiological Research Laboratory,
Central Public Health Laboratory,
London NW9 5HT

- ¹ Measles Vaccines Committee. *Br Med J* 1966;i:441-6.
- ² Measles Vaccines Committee. *Br Med J* 1965;i:857-60.
- ³ Miller DL. *Br Med J* 1964;ii:75-8.
- ⁴ Miller CL. *Br Med J* 1978;i:1253.
- ⁵ Orenstein WA, et al. *J Infect Dis* 1978;137:847-53.

SIR,—I see that Professor George Dick (10 May, p 1186) has been blandly asserting that measles is becoming a trivial disease in civilised communities.

We have had two cases of febrile convulsions in children under the age of 3 in the prodromal phase of measles infection in the last month. Both children were admitted to hospital, thereby infecting the entire ward, before the Koplik's spots appeared. Both developed severe otitis media.

From the standpoint of the GP, I believe that measles vaccination should be wholeheartedly encouraged in small children. I see no evidence of a decline in the virulence of this virus.

P M J O'DONNELL

Luton, Beds

Prophylaxis of surgical wound sepsis

SIR,—In your recent leading article on prophylaxis of surgical wound infection (19 April, p 1063) you correctly imply that antibiotic prophylaxis should be in addition to and not an alternative to meticulous surgical technique, and that the choice of the antibiotic used should be determined by the organisms likely to be encountered at operation and also on the prevailing hospital bacteriological sensitivity.

However, your comment that "we still do not know how many doses of antibiotic should be given" is rather misleading. As long ago as 1961 Burke¹ demonstrated in animal studies that antibiotics were only effective in dealing with bacterial contamination for three hours following an operation—in other words, over the immediate postoperative period. That this work is relevant to the clinical situation has clearly been demonstrated in several clinical trials. Short courses of preoperative prophylactic antibiotics given either systemically or intracutaneously have been shown to reduce wound sepsis after general²⁻⁵ and orthopaedic⁶ operations and given systemically to reduce chest infection following cardiac operations.⁷

Furthermore, whenever a short course has been compared to a long course of antibiotics no advantage has been shown for the latter.^{8,9}

The use of prophylactic antibiotics for longer than the preoperative period in either one or two doses given systemically, or given intracutaneously at the end of the operation, is therefore unnecessary in terms of efficacy and expense and will lead to a greater incidence of antibiotic toxicity problems and of emergence of resistant organisms. Finally, it should be emphasised that this type of short course is not appropriate when antibiotics are used therapeutically to treat an established infection.

R B GALLAND

Royal Postgraduate Medical School,
Hammersmith Hospital,
London W12 0HS

- ¹ Burke JF. *Surgery* 1961;50:161-8.
- ² Galland RB, Saunders JH, Mosley JG, et al. *Lancet* 1977;ii:1043-5.
- ³ Griffiths DA, Shorey BA, Simpson RA, et al. *Lancet* 1976;ii:325-8.
- ⁴ Stokes EJ, Waterworth PM, Franks V, et al. *Br J Surg* 1974;61:739-42.
- ⁵ Pollock AV, Evans M. *J Antimicrob Chemother* 1975;1, suppl:71-83.
- ⁶ Pollard JP, Hughes SPF, Scott JE, et al. *Br Med J* 1979;ii:707-9.
- ⁷ Galland RB, Shama DM, Prenger KB, et al. *Br J Surg* 1980;67:97-8.
- ⁸ Conte JE Jr, Cohen SR, Roe BB. *Ann Intern Med* 1972;76:943-9.
- ⁹ Strachan CJL, Black J, Powis SJA, et al. *Br Med J* 1977;i:1254-6.

Outbreak of infection in a urological ward

SIR,—We were very interested to read the article about an outbreak of *Acinetobacter* infection in a urological ward (8 March, p 722). We are also isolating this organism.

Acinetobacter calcoaceticus was first isolated from both male and female patients in the urological ward. Since then, with the transfer of these patients to different wards, we have isolated it from patients in the intensive therapy unit and a general surgical and a general medical ward. In only two of these patients has the organism been isolated from blood. Our environmental screening has been unhelpful and the *Acinetobacter calcoaceticus* returned to the intensive therapy unit despite closing, cleaning, and redecorating the unit. It was not isolated from the environment prior to closing or before reopening the unit. None of the new patients was colonised on admission.

This organism rarely causes clinical infection, but the pattern of multiple antibiotic resistance can make treatment very difficult. How can we best prevent its spread throughout the hospital?

MARY ANNE L BURROW

PENELOPE J REDDING

Department of Bacteriology and
Immunology,
Western Infirmary,
Glasgow G11 6NT

Preventing infection in clinical laboratories

SIR,—The contribution by Sir James Howie and Mr C H Collins about the Howie code of practice for preventing infection in clinical laboratories (19 April, p 1071) is both timely and welcome.

If clinical laboratories in the United Kingdom had been constructed to a standard design, no doubt the task of the working party would have been more straightforward; the fact is they have not. In general, clinical