recently taken drugs that are known to induce liver enzymes, and at present this must include recent exposure to enflurane.⁸

If for repeated anaesthesia one substitutes enflurane for halothane one may be replacing the rare case of hepatic disease with the perhaps more common case of renal failure.

MIKE SINCLAIR

Department of Anaesthetics,
Southmead Hospital,
Bristol BS10 5NB

¹ Chase RE, Holaday DA, Fiserova-Bergerova V, Saidman LJ, Mack FE. *Anesthesiology* 1971;35:262-7.

² Corall IM, Knights K, Strunin L. *Br J Anaesth* 1977;49:881.

³ Cousins MJ, Greenstein LR, Hitt BA, Mazze RI. *Anesthesiology* 1976;44:44-53.

⁴ Cousins MJ, Mazze RI. *JAMA* 1973;225:1611-6.

⁵ Cousins MJ, Mazze RI, Kosek JC, et al. *J Pharmacol Exp Therap* 1974;190:530-41.

⁶ Berman ML, Green OC, Calverley RK, Smith NT, Eyer EI. Enzyme induction by Enflurane in Man. *Anesthesiology* 1976;44:496-500.

⁷ Eichhorn JH, Wedley-Whyte J, Steinman I, Kaufman JM, Laasberg LH. *Anesthesiology* 1976;45:557-60.

⁸ Hewer CL, Atkinson RS, eds. *Recent advances in anaesthesia and analgesia* 13. Edinburgh: Churchill Livingstone, 1979:4.

Management of abnormal cervical smears

SIR,—Your leading article on the management of abnormal cervical smears (24 May, p 1239) was informative and encouraging to all concerned with the problem of cervical cancer. But why, oh why, must you spoil it all by saying, "Until a cure for cancer is found early detection offers the best chance for survival." Cervical cancer is in fact an eminently curable type of cancer and in the context of curability (not just "survival") many other cancer types are equally curable. A patient with cervical cancer, properly treated, can be *cured* in the sense of acquiring a life expectancy identical to that of any other woman of the same age. This actuarial definition of cure¹ not only is statistically valid but should also provide great comfort to doctors and patients alike. It is sad therefore that so many workers in the cancer field seem so unaware of the encouraging facts about this disease.

ERIC EASSON

Disley, Cheshire SK12 2BB

¹ Easson EC, Russell MH. *The curability of cancer in various sites*. London: Pitman Medical, 1968.

SIR,—I am in total agreement with your leading article (24 May, p 1239) that more effort should be made to give general practitioners incentive to do follow up routine screening of cervical smears. At present we are paid a fee only if the patient is over 35 years of age—or under 35 if she has had three or more pregnancies—and then it is only once every five years.

I think that in this era of preventive medicine general practitioners should be encouraged to take cervical smears, which take up much time and should attract a fee when performed on any woman of child-bearing age—which they should be yearly, especially in the over-35 age group—and that any follow-up requested by the laboratory or on the opinion of the GP should attract, say, a half fee.

In this way many more GPs would be encouraged to do routine cervical smear screening on most female patients and thus,

hopefully, reduce the incidence of carcinoma of the cervix.

JONATHAN LEVY

London N3 1LD

Indomethacin and perforated duodenal ulcer

SIR,—The criticism by Professor M J S Langman and Dr D A Henry (19 April, p 1088) of the statistical analysis by one of us (MRT, 16 February, p 448) showing that indomethacin is an important cause of duodenal ulceration in elderly women is correct only on the basis that the article may not have made it sufficiently clear that the association between non-steroidal anti-inflammatory drugs and perforated duodenal ulcers was suspected before the study was made. In fact the association has been suspected for many years (many surgeons would claim that it has already been established) and the article did refer to a previous paper¹ which provided statistical evidence for the association. Thus if the article had started, "In view of the previously reported association between non-steroidal anti-inflammatory drugs and perforated duodenal ulcers a retrospective review of..." the statistical analysis performed would have been entirely valid. This apparent lack of clarity in the article should not detract from the important conclusion made.

A further study of gastric ulcer patients has in fact been made as suggested by Professor Langman and an almost identical picture has emerged. Twenty out of 58 (34%) female patients over the age of 60 treated for this condition at the Bristol Royal Infirmary over the last five years were taking either phenylbutazone or indomethacin and eight (14%) were taking aspirin. Most of these patients had chronic ulcers. Further control groups are being collected to determine the significance of these findings. It is likely that they will show that indomethacin is also an important factor in the aetiology of some gastric ulcers, particularly in elderly women.

M R THOMPSON

St Mark's Hospital,
London EC1V 2PS

AUDREY F MORRIS

A O HUGHES

Department of Community Health,
Bristol BS8 2PR

¹ Jørgensen TG. *Br J Surg* 1977;64:247-9.

Pathological investigation of maternal deaths: a cause for concern

SIR,—As the consultant advisers in obstetrics and gynaecology to the Department of Health and Social Security, we write to congratulate Dr R I Rushton (15 March, p 767) for the very positive way in which his paper responds to the concern expressed about the value of some necropsies in the latest report on confidential inquiries into maternal deaths in England and Wales for the years 1973-5.¹

In the preface to that report Sir Henry Yellowlees, the Chief Medical Officer, suggested that more discussion between clinicians and pathologists might help to elucidate some of the problems; and later it was suggested that a consultant adviser in pathology should become a central assessor in maternal mortality studies. This suggestion,

which we both favour, was also recommended by one of us (JST) in a paper² reviewing the confidential inquiries since their inception in 1952.

We therefore now wish to report in your columns that the Chief Medical Officer has appointed the consultant adviser in histopathology to the Department of Health and Social Security, Professor I M P Dawson of the department of pathology, University of Nottingham, as a central assessor in maternal mortality studies.

JOHN TOMKINSON

London W1N 1DG

A C TURNBULL

John Radcliffe Hospital,
Oxford OX3 9DU

¹ Department of Health and Social Security. *Report on confidential enquiries into maternal deaths in England and Wales, 1973-75*. London: HMSO, 1979.

² Tomkinson JS. *Health Trends* 1979;4:77-80.

Contraception by female sterilisation

SIR,—Your leading article (10 May, p 1154) rightly states that "the alleged association between sterilisation and menorrhagia is disputed..." and goes on to quote a number of hospital-based studies. A view from general practice, where gynaecological symptoms are common and many patients are managed without referral, may be of interest to your readers.

In my practice in 1977¹ 18.6% of a random sample of 210 women from 20 to 50 years had been sterilised and the figure is now over 22% and rising. Sterilisation is therefore numerically significant as a contraceptive choice and any associated long-term morbidity may have important effects on the logistics of primary care. Of 368 patients followed up by me, 42% considered menstruation worse after tubal sterilisation and only 13% found it better. The main complaint was of heavier periods, which were often also less regular. Menstrual problems before sterilisation were reported at interview by 101 patients (27.4%). When compared with controls of the same age not taking oral contraceptive, 46% of the sterilised patients complained of current menstrual problems compared with 20% of their matches. While the sterilised group may have been more vulnerable to gynaecological upset before operation, the difference seems large, even allowing for variations in conceptions (mean number of pregnancies 3.5 for the sterilised and 2.6 for controls).

Many authors²⁻⁵ have reported an increase in menstrual disturbances after sterilisation; this has been linked to operation type by Neil *et al*⁶ and withdrawal of cyclic hormone by Chamberlain and Foulkes.² It is not easy to obtain direct data on menstrual dysfunction; menstrual calendars have their limitations, as has the logging of "heavy days" or pad or tampon counts. Kasonde and Bonnar⁷ have directly measured blood loss, before and after operation, in 25 women using the alkaline haematin method of Hallberg and Nilsson.⁸ In clinical practice the use of such painstaking direct methods would present major problems in the handling and transmission of material.

The work of the GP as doctor of first contact and, more important, as doctor of continuing care is quickly affected by any substantial change in the prevalence of menstrual disorders. There is a need for continuing prospective assessment of the effects of sterilisation on menstrual function to test the reality of this

"poststerilisation syndrome,"⁹ which I believe does exist. A combined approach by gynaecologists and general practitioners is likely to be most fruitful.

The analysis of consulting patterns and referrals for gynaecological complaints in sterilised women, both before and after operation, and in controls is currently being explored in this practice.

A F WRIGHT

Glenwood Health Centre,
Glenrothes, Fife KY6 1HL

- ¹ Wright AF. *Health Bull (Edinb)* 1978;**36**:229-34.
- ² Chamberlain G, Foulkes J. *South Med J* 1976;**69**:1474-5.
- ³ Muldoon MJ. *Br Med J* 1972;**i**:84-5.
- ⁴ Neill JG. *Ulster Med J* 1969;**38**:119-22.
- ⁵ Whitehouse DB. *Adv Fertility Control* 1969;**4**:22-6.
- ⁶ Neil JR, Hammond GT, Noble AD, Rushton L, Letchworth AT. *Lancet* 1975;**iii**:699-700.
- ⁷ Kasone JM, Bonnar J. *Br J Obstet Gynaecol* 1976;**83**:572-5.
- ⁸ Hallberg L, Nilsson L. *Scand J Clin Lab Invest* 1964;**16**:244-8.
- ⁹ Swinhoe J. *J Maternal Child Health* 1977;**2**:304-9.

Female sterilisation—no more tubal coagulation

SIR,—Your leading article "Female sterilisation—no more tubal coagulation" (12 April, p 1037) contained the provocative message that the gynaecologist should be careful to ensure that his operation is reversible. If his patient has been properly counselled his main concern should be to ensure that his operation is effective.

Gynaecologists who have used laparoscopic tubal diathermy safely and effectively for many years are likely to be disappointed with the high pregnancy rate with clip sterilisation and with the risk of tearing the tube and broad ligament with ring sterilisation. Despite all the effort and inventiveness, there is still no practical alternative to diathermy, preferably bipolar, for laparoscopic sterilisation.

A KENNEY

St Thomas's Hospital,
London SE1 7EH

Setting up a drip

SIR,—Dr John Aspinall is incorrect in his assumption (3 May, p 1146) that we recommend intramuscular morphine sulphate alone for sedation and analgesia in distressed children prior to venepuncture. Our experience has been that it is of value when used in combination with oral trimeprazine for tranqualgesia as described (29 March, p 942). A suitable alternative to oral trimeprazine would be oral diazepam 0.15 mg/kg, while anaesthetists will be familiar with the use of rectal thiopentane or ketamine by the intramuscular or intravenous routes in refractory cases. Perhaps a solution to this problem for the future may lie in the use of an epicutaneous pad (Ane-Pad, Astra Lakemedel AB) containing the local anaesthetic ketocaine applied to the proposed venepuncture site. While this technique was described¹ for cutting split skin grafts in adults an extension of its use for minor superficial procedures in children deserves investigation.

DERMOT PHELAN
MYLES MACDEVILLY

Dr. Steevens' Hospital,
Dublin 8

- ¹ Ohlson L, Englesson S. *Br J Anaesth*. 1980;**52**:413-7.

Stimulation of dorsal column in multiple sclerosis

SIR,—We apologise for not having cited some of Dr L S Illis's work in relation to dorsal column stimulation (24 May, p 1273). In fairness, however, only one of the six articles quoted were available at the time of final submission of our article (29 March, p 889). Three articles appeared subsequently earlier this year and could not be included. The proceedings of the Belgrade Symposium on External Control of Human Extremities presented difficulty. This was not available through interlibrary loan and we felt it wrong to quote them. Our own group's findings in fact appeared in these proceedings and we deliberately do not refer to them. This leaves us with one reference¹ which we could have included but did not, possibly because only two cases were evaluated.

We made no claim for a "first description" of any major response in dorsal column stimulation. Full credit for most of this goes to Professor A W Cook of Brooklyn, New York. Dr Illis and Dr E M Sedgwick are incorrect in assuming we have simply confirmed their group's findings. We agree broadly that bladder function is benefited but do not agree about voluntary motor function. In their recent article² mention is made of improved motor function in "27.7% of 18 patients," which was felt not to be due to placebo effect, etc. We also found change in gait but thought this was a placebo effect. Furthermore, mention is made² of improved sensation—"27.7% of 18 patients." One hears of descending sensory levels in non-anatomical pattern following dorsal column stimulation. The careful studies on sensation reported in our paper—some in conjunction with Dr J Scadding of Professor P Wall's unit, University College, London—suggested that no change had occurred. Sometimes a patient claimed benefit which could not be substantiated and this again is likely to be a spontaneous fluctuation or placebo effect.

Certain recent publications suggest that dorsal column stimulation has no real effect at all in multiple sclerosis.^{3,4} Perhaps we should devote more time to validating our results than disputing priorities and references to the literature.

C H HAWKES

Department of Neurology,
Ipswich Hospital,
Ipswich IP4 5PD

- ¹ Illis LS, Sedgwick EM, Oygur AR, Sabbahi Awadalla. *Lancet* 1976;**iii**:1383-6.
- ² Illis LS, Sedgwick EM, Tallis RC. *J Neurol Neurosurg Psychiatry* 1980;**43**:1-14.
- ³ Young RF, Goodman SJ. *Neurosurgery* 1979;**4**:225-8.
- ⁴ Rosen JA, Barsoum A. *Ann Neurol* 1979;**6**:66-7.

Beta-blockers in immediate treatment of myocardial infarction

SIR,—We are anxious to comment on the trial of beta-blockers given early after a myocardial infarction that appeared recently (29 March, p 885). It was refreshing to read such a clear and open account of the trial findings. Despite their experience with withdrawals, Dr R G Wilcox and his colleagues emphasise that to minimise potential bias it is essential to include in the main analysis all patients randomised, regardless of their course during the study. We agree wholeheartedly. With the same purpose in mind, we too would advocate the use of total mortality as the primary response variable. It is based on a

simple count with no reliance on definition or interpretation.

We must, however, take issue with the conclusion of the authors. The trial showed that at six weeks there was a 25-33% reduction in total mortality in the atenolol and propranolol groups compared with the placebo group but, because this does not reach statistical significance, it is stated that there is no place for beta-blockers in the acute management of patients with suspected myocardial infarction. We support the plea of Professor Geoffrey Rose (19 April, p 1988) that in any clinical trial with equivocal results either confidence limits or the power—that is, the likelihood of detecting a real difference—should always be given. The trial of the Nottingham group was too small to have adequate power to test the hypothesis that beta-blockers may be of benefit if started immediately after myocardial infarction. With approximately 130 patients in each of the three treatment groups and an $\alpha=0.05$, the chance of detecting a true benefit was less than 30% even when total mortality was reduced by as much as one-third. Should any intervention be shown to have an effect of this degree its impact on public health would indeed be considerable. Unfortunately in small trials a verdict of "not proven" is all too common. The question of whether or not beta-blockers prolong life in heart attack victims can be answered only by a well-designed clinical trial of sufficient size.

CURT FURBERG
GRAHAM MAY
HANS WEDEL

National Heart, Lung, and Blood Institute,
Bethesda, Maryland 20205,
USA

SIR,—Dr R G Wilcox and others (29 March, p 885) have admirably shown how serious bias can be created in small trials if patients are withdrawn after randomisation when they fail to adhere to treatment (a point not always appreciated). We believe that their paper too suffers from some defects in design, so that one cannot conclude, as they do, that beta-blockade is of no practical use in acute myocardial infarction.

Firstly, after oral treatment significant beta-blockade is achieved too late to have any large effect on serious arrhythmias or on infarct size, which are the main causes of mortality and morbidity. Additionally, in the atenolol-treated group a proportion of the patients would have received a placebo tablet initially (midday), thereby causing a further delay of at least eight hours before the patients received any beta-blocker. This therefore reduced the chance of testing the hypothesis that early beta-blockade may reduce mortality by reduction of infarct size or serious arrhythmias.

We too started a similar trial of oral beta-blockade three years ago and abandoned this when we observed that significant bradycardia was not seen for three to four hours after the dose—that is, about six to 12 hours after the onset of infarction. This is perhaps too late to influence infarct size (and hence mortality and morbidity). We therefore embarked on a morbidity (infarct size) study with intravenous beta-blockade. Our statistical advice was that we would need very much larger numbers than our projected trial size of 200-400 to detect any influence on mortality (see also Professor Geoffrey Rose's letter in the *BMJ* of 19 April, p 1080).