Studies using simulated images have shown that it is possible to reduce radiation doses by half or more by having slightly less clear images, without affecting the detection of simulated pathologies.^{9 w4} Organisations such as the International Commission on Radiological Protection and the International Atomic Energy Agency recommend that quality assurance programmes should be implemented,¹⁰ as do the UK regulations.¹¹

One big advance of digital technology is, of course, that it enables the electronic transfer of images to any location. This timely production and transmission of images gives physicians greater access to them during consultations.¹² The option of remote interpretation of images has the potential to ease the burden on hard pressed radiology departments. Images taken in short staffed departments at night can be reported by remote fully staffed departments in the day.¹

Despite the obvious benefits of digital x ray imaging, there is lack of clarity about related ethicolegal issues. Whether the current law relates to telemedicine in the same way that it does to other medical specialties remains controversial.¹ Telemedicine raises licensure questions, such as whether teleradiologists practising medicine in another country need a license for that place. This is complicated by individual hospitals having different standards for accreditation.

In the UK, practice guidelines on legal issues relating specifically to telemedicine or teleradiology are lacking. The next step is to develop and implement guidelines to safeguard both patients and professionals. Until such guidelines exist, practitioners should be

mindful of their position regarding issues such as duty of care, liability, and confidentiality.

Kwan-Hoong Ng *professor and senior consultant*
(ngkh@um.edu.my)

Department of Radiology, University of Malaya, 50603 Kuala Lumpur, Malaysia

Madan M Rehani *radiation safety specialist*

Radiological Protection of Patients Unit, International Atomic Energy Agency, Vienna, Austria

Competing interests: None declared.

- 1 White P. Legal issues in teleradiology—distant thoughts! *Br J Radiol* 2002;75:201-6.
- 2 Andriole KP. Productivity and cost assessment of computed radiography, digital radiography, and screen-film for outpatient chest examinations. *J Digit Imaging* 2002;15:161-9.
- 3 Dalla Palma L, Grisi G, Cuttin R, Rimondini A. Digital vs conventional radiography: Cost and revenue analysis. *Eur Radiol* 1999;9:1682-92.
- 4 Wang J, Langer S. A brief review of human perception factors in digital displays for picture archiving and communications systems. *J Digit Imaging* 1997;10:158-68.
- 5 Samei E, Hill JG, Frey GD, Southgate WM, Mah E, Delong D. Evaluation of a flat panel digital radiographic system for low-dose portable imaging of neonates. *Med Phys* 2003;30:601-7.
- 6 International Commission on Radiological Protection. *Managing patient doses in digital radiology*. Amsterdam: Elsevier, 2005.
- 7 Axelsson B, Boden K, Fransson SG, Hansson IB, Persliden J, Witt HH. A comparison of analogue and digital techniques in upper gastrointestinal examinations: absorbed dose and diagnostic quality of the images. *Eur Radiol* 2000;10:1351-4.
- 8 Reiner BL, Siegel EL, Flagle C, Hooper FJ, Cox RE, Scanlon M. Effect of filmless imaging on the utilization of radiologic services. *Radiology* 2000;215:163-7.
- 9 Mayo JR, Whittall KP, Leung AN, Hartman TE, Park CS, Primack SL, et al. Simulated dose reduction in conventional chest CT: validation study. *Radiology* 1997;202:453-7.
- 10 International Atomic Energy Agency. *International basic safety standards against ionizing radiation and for the safety of radiation sources*. Vienna: IAEA, 1996.
- 11 Department of Health. *The ionising radiation (medical exposure) regulations*. London: DoH, 2000. (IR(ME)R 2000.)
- 12 Jarvis L, Stanberry B. Teleradiology: threat or opportunity? *Clin Radiol* 2005;60:840-5.

doi 10.1136/bmj.38977.669769.2C

Preoperative staging for rectal cancer

Magnetic resonance imaging can accurately predict the success of surgical resection

Colorectal cancer is the second most common cause of death from malignant disease in the United Kingdom, with about 20 000 deaths each year. Around one million new cases (9% of all cancers) are diagnosed each year worldwide (CANCER*Mondial*; www-dep.iarc.fr). As the UK population ages the incidence is predicted to rise.¹ At present the only realistic prospect of cure is complete surgical resection of the primary tumour. The restricted anatomical space in the pelvis makes this technically easier to achieve for cancers of the colon than the rectum. Consequently, local recurrence rates after surgery for rectal cancer have been as high as 50%. Local recurrence is a devastating complication as it is invariably fatal even without disseminated disease. Local recurrence can be reduced by two methods—surgical technique and radiotherapy. Currently, radiotherapy is given to most patients even though only a subgroup will benefit. There is increasing interest in the use of preoperative staging to target high risk patients who will benefit most from radiotherapy. A study in this week's *BMJ* shows the value of magnetic resonance imaging in the preoperative staging of rectal cancer.²

Local recurrence rates can be reduced to less than 5% if the surgeon removes the rectum en bloc with the mesorectum using precise anatomical dissection (total mesorectal excision).³ Crucially, the surgeon must ensure that the circumferential resection margin is clear of tumour because a positive margin predicts a high risk of local recurrence.⁴ The quality of this surgery, which is technically demanding, can be determined by pathological examination of the integrity of the mesorectum in resected specimens. In Sweden, the adoption of total mesorectal excision by surgeons who had attended a training programme reduced the local recurrence rate after surgery for rectal cancer from 20% to 8% and increased survival.⁵

Randomised trials have shown that preoperative radiotherapy reduces local recurrence and is superior to postoperative treatment.^{6 7} The Swedish rectal cancer study showed that "short course" preoperative radiotherapy reduced local recurrence and improved survival, but this study was subsequently criticised because total mesorectal excision was not used.⁸ In a later Dutch trial all patients had total mesorectal excision and were randomised to preoperative radiotherapy or surgery alone.⁹ As expected with high quality

Research p 779

BMJ 2006;333:766-7

surgery the benefit from radiotherapy was less; local recurrence was reduced from 8% to 2%. Furthermore, the size of the benefit depended on the stage of the primary tumour—0.2% for T1 tumours, 5% for T2, and 12% for T3 primary lesions. These data can also be expressed as the proportion of patients who received unnecessary radiotherapy—99.8% of patients with a T1 tumour and 95% of those with T2 lesions. Even for patients with relatively high risk T3 tumours the figure was 88%. This is important because the Swedish and Dutch trials reported higher morbidity, including bowel dysfunction and incontinence, in patients treated with radiotherapy.^{10 11} Ideally, patients at high risk of a positive resection margin because of advanced primary disease should be identified before surgery so they can be targeted with preoperative chemotherapy or radiotherapy.

The study in this week's issue from the MERCURY Study Group shows that pelvic magnetic resonance imaging performed before surgery for rectal cancer has a sensitivity of 94% and a specificity of 92% for predicting negative circumferential margins after surgery.² Crucially, the study showed that when radiologists undergo specific training and the technique is standardised, results are reproducible between centres. The technique was less reliable at predicting positive resection margins mainly due to localised tumour perforation or the presence of adjacent lymph nodes, but this does not limit the clinical value of pelvic imaging.

These data have important implications for the management of patients with rectal cancer. Firstly, although a previous study suggested that preoperative staging may be used to target radiotherapy to high risk patients, the number of patients included was small.¹² The MERCURY Study Group however has now confirmed in a multicentre observational study that magnetic resonance imaging may be used for this purpose. This means that patients with rectal cancer at low risk of local recurrence can avoid unnecessary and harmful chemotherapy or radiotherapy. Also resources can be focused towards those patients at high risk.

Secondly, future studies of chemotherapy and radiotherapy can now be targeted towards a homogeneous group of patients known to be at high risk of local recurrence after surgery. This should make it easier to design trials with sufficient power to determine the most effective treatments in high risk groups, the benefits of short course radiotherapy compared with longer courses, and the role of salvage surgery in both responders and non-responders.

Finally, preoperative staging provides an objective yardstick against which the quality of the surgical technique may be audited.

When used with histological assessment of the integrity of the mesorectum in resected specimens, surgeons will no longer be able erroneously to attribute positive resection margins after rectal cancer surgery to advanced disease rather than poor surgical technique.

Preoperative staging has been advocated in guidelines for the management of colorectal cancer and many clinicians are already using it.^{13 14} The task ahead is to make magnetic resonance imaging mandatory for all patients with rectal cancer before treatment decisions are made at multidisciplinary meetings.

Ian Finlay *consultant colorectal surgeon*

(Ian.Finlay@northglasgow.scot.nhs.uk)

Department of Coloproctology, Lister Surgical Unit, Glasgow Royal Infirmary, Glasgow G3 7ER

Competing interests: None declared.

- 1 Scottish Executive. *Cancer scenarios: an aid to planning cancer services in Scotland (03: colorectal cancer)*. Edinburgh: SE, 2001. www.scotland.gov.uk/Resource/Doc/30859/0012657.pdf (accessed 9 Oct 2006).
- 2 MERCURY Study Group. Diagnostic accuracy of preoperative magnetic resonance imaging in predicting curative resection of rectal cancer: prospective observational study. *BMJ* 2006 doi: 10.1136/bmj.38937.646400.55.
- 3 Heald RJ, Moran BJ, Ryall RD, Sexton R, MacFarlane JK. Rectal cancer: the Basingstoke experience of total mesorectal excision, 1978-1997. *Arch Surg* 1998;133:894-9.
- 4 Adam JJ, Mohamdee MO, Martin IG, Scott N, Finan PJ, Johnson D, et al. Role of circumferential margin involvement in the local recurrence of rectal cancer. *Lancet* 1994;344:707-11.
- 5 Martling A, Holm T, Rutqvist LE, Johansson H, Moran BJ, Heald RJ, et al. Impact of a surgical training program on rectal cancer outcomes in Stockholm. *Br J Surg* 2005;92:225-9.
- 6 Sauer R, Becker H, Hohenberger W, Rodel C, Wittekind C, Fietkau R, et al. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 2004;351:1731-40.
- 7 Bosset JF, Collette L, Calais G, Mineur L, Maingon P, Radosevic-Jelic L, et al. Chemoradiotherapy with preoperative radiotherapy in rectal cancer. *N Engl J Med* 2006;355:1114-23.
- 8 Swedish Rectal Cancer Trial. Improved survival with preoperative radiotherapy in resectable rectal cancer. *N Engl J Med* 1997;337:980-7.
- 9 Kapiteijn E, Marijnen CA, Nagtegaal ID, Putter H, Steup WH, Wiggers T, et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med* 2001;345:638-46.
- 10 Dalberg M, Glimelius B, Graf W, Pahlman L. Preoperative irradiation affects functional results after surgery for rectal cancer: results from a randomised study. *Dis Colon Rectum* 1998;41:543-9.
- 11 Peeters KC, van de Velde CJ, Leer JW, Martijn H, Junggeburst JM, Kranenburg EK, et al. Late side effects of short-course preoperative radiotherapy combined with total mesorectal excision for rectal cancer: increased bowel dysfunction in irradiated patients—a Dutch Colorectal Cancer Group study. *J Clin Oncol* 2005;23:6199-206.
- 12 Horgan A, Finlay IG. Pre-operative staging of rectal cancer allows selection of patients for preoperative radiotherapy. *Br J Surg* 2000;87:575-9.
- 13 Scottish Intercollegiate Guidelines Network. *Guideline 67. Management of colorectal cancer*. Edinburgh: SIGN, 2003. www.sign.ac.uk/guidelines/fulltext/67/index.html (accessed 6 Oct 2006).
- 14 The Association of Coloproctology of Great Britain and Ireland. *Guidelines for the management of colorectal cancer 2001*. London: ACPGBI, 2001. www.acpgbi.org.uk/download/colorectal-cancer.pdf (accessed 6 Oct 2006).

doi: 10.1136/bmj.38996.423102.BE

Use of mobile phones in hospitals

New guidelines are less restrictive but still overcautious

Mobile phones are widely used, but their use is still restricted in certain places including petrol stations, some areas in hospitals, and aircraft. Restrictions have been justified on the grounds of public safety, but the reasons behind these restrictions are often unclear. In hospitals, patients, visitors, and staff routinely breeze through wards with

their mobile phones switched on. As yet we have no evidence that this behaviour has serious consequences for patients. The lack of such evidence has encouraged the Medicines and Healthcare products Regulatory Agency (MHRA) to advise more selective restrictions on the use of mobile phones in hospitals (box).¹ While welcoming this relaxation, we wonder why it has taken