must be cast on the ability of the ACCR to detect or exclude pancreatitis in patients with acute abdominal pain.

A R W FORREST
P McMASTER
C S BOND

Departments of Biochemistry and
Surgery,
Addenbrooke's Hospital,
Cambridge

¹ Durr, H K, *et al*, *New England Journal of Medicine*, 1977, **296**, 635.

Stiff-neck syndrome

SIR,—During the past three months, six female members of a hospital staff of approximately 1600—three nurses, two domestics, and one administrator—have presented with a similar clinical picture. This consisted of the sharp onset of severe and persistent pain in the cervical region, which was rigidly held in the erect central position and resisted the least movement, active and passive, in any direction, the pain being greatly augmented by such attempts. With one exception resolution was complete within a few days and the impression gained was that this was spontaneous rather than attributable to the local application of heat and the collar supplied. One recurrence has taken place, of identical nature and duration. Systemic upset, pyrexia, other abnormal physical signs, and radiological changes were all absent.

Although chest pains have not infrequently been reported among the staff during this period, no example suggestive of Bornholm disease has been encountered nor has inquiry in the medical wards yielded information of its presence or that of the stiff-neck syndrome among the patients.

The clustering of the cases raises the possibility of an infective aetiology. It would be of interest to learn of experience of like nature.

J SHAFAR

Staff Health Centre,
Burnley General Hospital,
Burnley, Lancs

Suffocation and sudden infant death

SIR,—I wish to report the case of a 10-week-old male infant who was brought to the accident and emergency department of this hospital last November.

He was born by normal delivery to a healthy gravida 2 mother after an uneventful pregnancy and had previously appeared normal and healthy. On this afternoon he was put outside in his pram wrapped in an infant's sleeping bag with the hood loosely drawn around his head. When his mother next looked she found him to be white, rigid, and not breathing. The nylon lining of the sleeping bag was damp and stuck across his nose and mouth. He started to breathe when she turned him upside down and slapped his back. On arrival at the department he was pale and breathing with difficulty and had generally increased tone, especially in the right arm. He was given oxygen by face mask and rapidly became completely normal. Chest x-ray showed no evidence of inhalation and he has since been quite well.

The sleeping bag is made of two layers of closely woven nylon with a safety lining between. Adults who have experimented find that, *when wet*, the nylon sticks to the face and is quite impossible to breathe through. Although the child may have had a convulsion, either initially or as a result of the anoxia,

we were concerned that the way in which he was wrapped might have led to a fatal outcome. In recent years suffocation in clothes and bedding has largely been discounted as a cause of sudden unexpected death in infancy, but this incident suggests that it should still be considered.

The baby did not struggle or cry out, and if he had died one wonders what evidence would have remained to explain death. This case underlines the importance of trying to construct an exact picture of all the circumstances surrounding any unexplained death in infancy as a number of factors, some of them innocent and harmless in themselves, can prove overwhelming in combination.

ANN JAY

Sheffield Children's Hospital,
Sheffield

Krukenberg's chopsticks

SIR,—I was delighted to see your leading article on Krukenberg's chopsticks (21 January, p 129), and this is not because of any personal experience in the performance of this operation. Those of us who have had the privilege of working for a short time in the Orthopaedic Hospital in Bangladesh have all seen the quite remarkable results of this procedure as carried out by Ronald Garst in Dacca. Once you have felt the force of the pincer grip achieved by one of his patients, you do not come back a second time.

Over many years in Britain we have all been grilled against the place of this operative procedure for severe forearm injuries, but Garst's success with many patients in Bangladesh is a tribute both to him and to the value which this operation may have at the present time in many places in the world.

H D W POWELL

High Wycombe, Bucks

Ink caps and alcohol

SIR,—As a native of Switzerland, where edible fungi are much sought after and highly prized, I would like to add my support to Dr A P Radford's defence of *Coprinus comatus* (14 January, p 112).

This fungus, aptly known as the "shaggy ink cap" or "lawyer's wig," should not be confused with its close relative *C. atramentarius*, which has a smooth cap. Drs M J Caley and R A Clark (24-31 December, p 1633) draw attention to a serious interaction between *C. atramentarius*, which they describe as an edible fungus, and alcohol. In Europe symptoms due to this type of interaction are well known and are generally considered to be the result of acetaldehyde formation during detoxication of alcohol. It has been claimed that the fungus contains disulfiram,¹ but other workers have denied this.² It is worth noting, however, that despite the widespread culinary use of wild fungi in many areas of Europe, reports of serious cardiovascular symptoms due to such interaction are extremely uncommon. Mild hypotension, sometimes with alarming orthostatic features, may occur and has been reported elsewhere,³ but cardiac arrhythmias are very rarely seen.

Although many textbooks describe *C. atramentarius* as an edible fungus, such interactions as can follow its ingestion together

with alcohol (even if the latter is taken many hours later) cause it to be widely viewed with suspicion. *C. comatus*, however, is entirely free from this unpleasant complication and is not only safely edible but delicious.

LOTTE BROADHURST-ZINGRICH

Great Barton, Suffolk

¹ Simandl, J, and Franc, J, *Chemie Listy*, 1956, **50**, 1862.

² Wier, J K, and Tyler, V E, *Journal of the American Pharmaceutical Association* (Scientific edn), 1960, **49**, 426.

³ Buck, R W, *New England Journal of Medicine*, 1961, **265**, 681.

Treatment of hepatic osteomalacia

SIR,—The careful study by Dr R G Long and others of four patients with biliary cirrhosis (14 January, p 75) illustrates well the problems of hepatic osteodystrophy. It also raises one or two questions, particularly the authors' interpretation of the results of 1,25-dihydroxycholecalciferol (1,25-(OH)₂D₃) therapy.

While their patients had clinical bone disease the extent of osteomalacia was uncertain: their x-rays indicated osteoporosis and not osteomalacia; previous fractures either had healed or were healing, suggesting adequate vitamin D activity; metabolic myopathy may be obscure in patients near to death; and virtually all biochemical indices of osteomalacia, without evidence of secondary hyperparathyroidism and including calcium absorption and phosphorus excretion, were normal. It must be hard to be sure of clinical improvement following 1,25-(OH)₂D₃ treatment when the second biopsy specimens were obtained post mortem in two subjects. Nevertheless, if one accepts significance of histological changes following 1,25-(OH)₂D₃ and of the stated clinical improvement the possible reasons need to be more carefully considered.

The authors' data suggest that patients with biliary cirrhosis may have an increased requirement for vitamin D above that necessary to overcome malabsorption of the vitamin. But they fail to provide grounds for concluding that either 1 α -hydroxylation is at fault or that vitamin D₂ is inferior to vitamin D₃. Many other possibilities are at least as likely. In particular there is good evidence that deficiency of active metabolites may result from increased urinary loss in this disease.¹ Years of parenteral vitamin D therapy in the authors' patients had failed in two of them to raise plasma 25-hydroxyvitamin D (25-OHD) levels above 11.4 μ g/l (misprinted as pg/l in the paper); higher levels might be expected following oral therapy in patients without malabsorption.² One might expect similar urinary loss of other active metabolites. Furthermore, comparison of different parenteral routes of vitamin D administration as a means of raising plasma 25-OHD levels suggests that availability of vitamin D following injection of an oily bolus directly into storage organs (muscle and adipose tissue) may be limited.³ (Defective hepatic 25-hydroxylation of vitamin D, which would also explain inappropriately low 25-OHD levels, has been largely excluded by recent work.⁴) Whatever circulating activity is maintained in the face of increased loss and restricted availability, immediate substitution of 1,25-(OH)₂D₃ will simply be additive and tells little of the nature of any "resistance".

Additional mechanisms which are theoretically neither more nor less likely than the

authors' conclusions include increased faecal loss of active metabolites, increased intestinal breakdown by side-chain cleavage,⁴ defective synthesis of binding proteins, and target-organ resistance. The last might result from impaired nutrition and/or atrophy such as is presumably associated with resistance to parenteral treatment in coeliac osteomalacia.⁵ Alterations in side-chain cleavage by liver or intestine⁴ might also produce target-organ resistance from accumulation of interfering vitamin D metabolites, and other circulating compounds (bilirubin, ammonia, etc) might also interfere with vitamin D activity.

If further studies do confirm the suggestion that patients with biliary cirrhosis may show true resistance to parenteral vitamin D then it is important that all the questions should be asked and critically evaluated, since consequent research may prove any one (or more) of them to be correct.

T C B STAMP

Royal National Orthopaedic
Hospital,
Middlesex

¹ Krawitt, E L, *et al*, *Lancet*, 1977, 2, 1246.

² Stamp, T C B, Haddad, J G, and Twigg, C A, *Lancet*, 1977, 1, 1341.

³ Whyte, M, Stamp, T C B, and Haddad, J G, in *Endocrinology 1977*, ed I MacIntyre *et al*. In press.

⁴ Kumar, R, and DeLuca, H F, *Biochemical and Biophysical Research Communications*, 1977, 76(2), 253.

⁵ Nassim, J R, *et al*, *Quarterly Journal of Medicine*, 1959, 28, 141.

Household products and poisoning

SIR,—I was interested to see the paper by Dr Roy Goulding and others (4 February, p 286). I was not surprised by his findings, as the range of items included in the research are responsible for only about 16.5% of the poisonings from household products. If the research had been widened to include turpentine and white spirit, paraffin, disinfectant, surgical spirit, perfume, and alcohol he would have covered a further 35% of the problem.

Dr Goulding's conclusions are misleading in that he implies that the dangers of household products are being exaggerated. He does not say who he feels is exaggerating the problem. We would like to see more done to prevent poisonings from both medicines and household products.

While ingestion of these substances may well require no treatment, Dr Goulding makes no allowance for the distress to the child and his family that such an incident can cause. Prevention is still better even if no cure is needed.

ELSPETH G MACLEAN
Director for Home Safety,
Royal Society for the
Prevention of Accidents

Birmingham

Glucagon levels and vagotomy

SIR,—Dr L V Campbell and others (10 December, p 1527) suggest that the glucagon peak expected during hypoglycaemia is absent in some diabetics because of vagal neuropathy. Russell *et al*¹ demonstrated that under similar conditions the glucagon rise was significantly less after truncal vagotomy than after selective vagotomy. These results implied that provided the coeliac and hepatic branches of the vagus remained functional a normal glucagon response to hypoglycaemia should be expected. It would seem therefore that glucagon assay during hypoglycaemia is

potentially an index of the selectivity achieved during vagotomy. This may prove useful in assessing and comparing operative techniques.

Dr Campbell and his colleagues suggest that increased glucagon release is essential for recovery from hypoglycaemia. Russell *et al* found no difference in the rise in plasma concentration of enteroglucagon (which stimulates insulin release but does not have the hepatic glycogenolytic/gluconeogenic capacity of pancreatic glucagon) after a glucose load between patients who had undergone truncal and those who had had selective vagotomy. Unfortunately they did not state if any patients suffered from "late dumping" or developed it after the glucose load.

Thus, postvagotomy and postprandially, enteroglucagon could stimulate insulin release and the low pancreatic glucagon level would be unable to offset the impending hypoglycaemia. Could this situation contribute to the pathogenesis of late (hypoglycaemic) dumping? Would it be of therapeutic value to attempt to raise postprandial glucagon levels in these patients?

D P MIKHAILIDIS

London N6

¹ Russell, R C G, Thomson, J P S, and Bloom, S R, *British Journal of Surgery*, 1974, 61, 821.

The new Sinbad syndrome

SIR,—I have read with interest and amusement the recent column "The new Sinbad syndrome" by our recent distinguished past president, Dr Peter Banks (18 February, p 423).

Dr Banks states that my observations regarding the growing "antigovernment paranoia" exhibited in his writing was "not becoming" to him. He also suggests that this change in the tone of his writings (a more basic, naked distrust of governments) does not indicate a progressively worsening paranoid state but rather a recognition of reality.

Verily, Peter, I did not suggest that your written observations on Canadian government, on how we are grossly overgoverned, our out-of-control Civil Service, the detrimental effect of these facts on life in general and medicine in particular, etc, were unbecoming or incorrect. I simply pointed out that if you continue to write of their existence in a systematic, albeit on occasion slightly exaggerated, fashion they are so extensive and grotesque that the resultant chronicle may jeopardise your credibility with UK readers.

The transmission of a message across the oceans is a very fragile operation more frequently requiring, and benefiting from, the approach and techniques of the neurosurgeon rather than those of the orthoped.

D A GEEKIE
Director of Communications
Canadian Medical Association

Ottawa, Ontario

General practice expenses

SIR,—Our partnership has just received our audited accounts for 1976-7.

For the past two years our accountants have calculated the percentage of our costs against our income and the frightening situation now shows that this figure has risen to 45%, an increase of 10% over the past year. This figure does not take into account each partner's

personal expenses for his car and maintenance for emergency facilities in his own home.

We wonder how many of our fellow general practitioners have found it impossible to increase their banker's orders over the past three years? This is the sorry tale we have to tell, and the sad part about it is that it is not due to a static income—in fact our work load has increased, our numbers have increased, and extra work done has claimed extra fees. We have tried to look at the problem objectively, and in our particular case we feel that the three problems outlined below perhaps are responsible for this rather sad situation.

(a) Our local pathological laboratory is 26 miles away from here and all telephone calls made before 2 pm are at the high tariff rate. In 1975 our telephone costs were running at £1200, in 1976 they had risen to £1750, and now in 1977 the figure is £1960.

(b) Our heating expenses have risen by over 100% to £650 per annum during these past three years.

(c) Our maintenance costs—painting, replacement of carpets, etc—have risen from an average of £250 to £550 per annum.

This practice is an urban and rural one; 32% of its patients, approximately 1500, are over 65. Each partner interests himself in local organisations, among which are listed St John's, the Round Table, and the Lifeboat Institution, and each participates in the local welfare of this area. This year has seen a drop in our net income of £1000 for each of us; this now means that two of us have to have extended overdrafts because the cost of living and maintaining a family has risen by some 30% and yet our income in real terms has fallen by this amount.

We do not want to lower our standards of medical care and attention; we can only hope that the Review Body and the general public will realise just how serious our plight now is. Unless the Government can find ways and means of reimbursing us adequately then we must ask the question, "Where is it all going to end?"

A WEST COUNTRY GENERAL PRACTITIONER

Community Medicine Consultative Committee

SIR,—A very confused debate took place at the meeting of the Central Committee for Community Medicine held on 10 February.

It concerned the formal constitution and inauguration of the all-important Community Medicine Consultative Committee between the BMA, the Faculty of Community Medicine, and the Department of Health and Social Security. I regret to have to inform colleagues in community medicine that a formal constitution has not been agreed even after four years have elapsed since reorganisation, and it may not be clear from the report of the CCCM proceedings quite why this is so important.

I therefore urge all community physicians to question their regional representatives on this matter and to mandate them very clearly for the next full meeting of the CCCM on 12 May this year, when this subject will no doubt be discussed further. An early decision must be made.

PAUL J HEATH

Sheffield Area Health Authority
(Teaching),
Sheffield