need to complete the randomised controlled trials currently under way.

R R H LOVELL
R D ULMAN

University of Melbourne,
Department of Medicine,
The Royal Melbourne Hospital,
Victoria

W B STEPHENS

Albury,
New South Wales

¹ Reader, R, *Acta Medica Scandinavica*, 1975, **197**, 5765.

² Princeas, R J, Stephens, W B, and Lovell, R R H, *Medical Journal of Australia*, 1973, **1**, 5.

³ Lovell, R R H, et al, *Australian and New Zealand Journal of Medicine*, 1976, **6**, 398.

SIR,—There is little doubt that most physicians working in geriatric departments would endorse Dr Anthony Martin's scepticism (27 November, p 1326) regarding the value of hypotensive therapy on the elderly and many will have seen catastrophes resulting from such treatment, sometimes for one reason or another leading to the patient's premature death.

Dr Martin's prospective study showing significantly better survival among patients with diastolic blood pressures of 100 mm Hg or more is interesting but does not really surprise me. After all, as long ago as 1964 a leading neurologist¹ considered that a drug (if one was available) for maintaining a higher blood pressure would probably be of more benefit to the elderly than hypotensive agents had been.

My own observations in a geriatric unit where perhaps above average use is made of the sphygmomanometer suggest that hypertension in the elderly is very often a peculiarly transient and intermittent condition. Diastolic levels of 120 mm Hg and more are frequently encountered in patients without cardiomegaly or other evidence of target-organ disease. In the vast majority of these the diastolic level falls to normal or very near normal within a few days with nothing more active in the way of treatment than bed rest and a diuretic and I am even inclined to think that both rise and fall are probably wholly spontaneous events. Certainly it has not been found possible to blame emotional disturbance or other stress for the initial rise, neither have catecholamine levels been increased. Such hypertensive episodes may be symptomless but not infrequently are associated with a little vertigo, a feeling of anxiety or tension which is endogenous rather than reactive, and sometimes a mild confusional state. Anginal pain and occasionally heart failure have also been observed, but symptoms promptly disappear when the blood pressure falls. The patient will then remain normotensive and completely well for weeks or months, but usually further episodes eventually occur with the same pattern of symptoms and speedy resolution. It is easy to see how an attempt at continuous hypotensive therapy might be disastrous.

There seems to be a substantial difference between this type of case and the "legitimate hypertensive" with a large left ventricle and perhaps progressive neurological and renal deficit which may not be just a matter of severity and duration of disease. At the same time it is possible that in these transient cases the blood pressure may remain elevated sufficiently long or frequently for them to be included in studies such as Dr Martin has carried out and it would then not be sur-

prising if their expectation of life did not correspond with that forecast for hypertensives in life tables.

At any rate, without apology, I still have complete confidence in my sphygmomanometer whatever the age.

S L O JACKSON

Harold Wood Hospital,
Romford, Essex

¹ Williams, D, *British Medical Journal*, 1964, **1**, 84.

SIR,—The difficulty in deciding whether to treat asymptomatic and uncomplicated hypertension in the elderly is illustrated by Dr A Martin's figures on patient survival (27 November, p 1326). On the other hand one does not want to deny the benefit of treatment just on the grounds of age. His scepticism, however, on the wisdom of treating elderly hypertensives is reinforced by a clinical trial on hypotensive therapy in stroke survivors¹ which showed that there was no reduction in mortality with hypotensive treatment in people over the age of 65. In comparison, in the under-65 group the mortality for treated hypertensives was less than half that of the untreated.

D R SWINSON

Rheumatology Unit,
Wrightington Hospital,
Wigan, Lancs

¹ Barham Carter, A, *Lancet*, 1970, **1**, 485.

"Nurse consultants"

SIR,—I was interested to read of Dr D W Eyre-Walker's consternation over a nurse presuming to call herself a consultant (4 December, p 1386). Could it be that we have a latterday Rip Van Winkle in our midst who is just awakening to modern terminology? The local plumber now calls himself a central heating consultant and the farm worker who used to do a bit of earth heaving in his spare time now advertises his services as a landscaping consultant.

I do not think Dr Eyre-Walker need "fight to the death" to preserve his God-like status. As long as there are ill and frightened people in hospital this will be maintained—and the nurses will continue to aid and abet the consultant in this role as it is essential for the wellbeing of their patients.

JEAN KELLY

Felmersham, Beds

Chemoprophylaxis of malaria

SIR,—I have been reading with great interest your leading article on this subject (20 November, p 1215) and am rather puzzled by some of the statements that I find there.

In the middle of the second column there is a statement comparing pyrimethamine and proguanil with chloroquine. The sentence reads: "Both drugs, unlike chloroquine, act slowly against established infections of malaria, and this may reflect their chemosuppressive potency." I am rather puzzled by the part of the sentence I have italicised since it seems to me to be a non sequitur. I wonder what you had in mind? I am disturbed too by the inclusion of the last sentence of this paragraph which states: "Unofficially it is sometimes recommended that both should be taken—

chloroquine weekly and proguanil daily." I think it would have been much better to have left out this very illogical sentence.

As regards the use of Fansidar (pyrimethamine + sulfadoxine), I am not sure that enough studies have been made of the correct prophylactic dose for non-immunes. While two tablets once every two weeks is probably reasonable, I would think that one tablet once a week is almost certainly as good. The WHO report cited¹ does in fact suggest one tablet weekly or two tablets every two weeks. The majority of trials with Fansidar as a prophylactic have been in semi-immune populations and we cannot necessarily translate the results obtained in such trials into a firm recommendation for non-immunes. When I last looked into the matter the makers of Fansidar, Roche Products, were not actually recommending it for general prophylaxis for the simple reason that they did not feel that they had enough data on the dosage to make a general recommendation. Perhaps they have now come to some conclusion on this and, if so, I would be interested to see the reference. I certainly approve of this particular combination in areas where chloroquine-resistant falciparum malaria is present and am interested to see that WHO is now also recommending it (but please note, as I have mentioned above, that the WHO recommendation gives the alternative weekly or fortnightly dosage). It is regrettable that the WHO report still gives the alternative of taking chloroquine as 300 mg base once a week, 300 mg twice a week, or 100 mg base daily. I never did agree with the last two of these three alternatives and I am glad that you have taken up this question in some detail at the top of your second column.

Finally, in your last paragraph it is recommended that drug prophylaxis be continued for eight weeks following the departure of a traveller from an endemic area. There must surely be some confusion here. As far as the prevention of falciparum malaria is concerned the continuation of appropriate prophylaxis for four weeks is really adequate. The eight-week regimen is one that was established for the prevention of relapses due to Korean *Plasmodium vivax*, the prophylactic against which was the US Army regimen of 300 mg chloroquine base plus 45 mg primaquine base once per week. Taken regularly for eight weeks this was very effective in preventing late relapses due to this particular parasite. One must be quite clear on this point.

W PETERS

Department of Parasitology,
Liverpool School of Tropical Medicine,
Liverpool

¹ World Health Organisation, *Information on Malaria Risk for International Travellers, Weekly Epidemiological Record*, 1976, **51**, 181.

*.*Only a controlled prospective study will confirm or deny our suggestion that the chemosuppressive potency of a drug is related to its chemotherapeutic potency. Chemosuppression is a better term than chemoprophylaxis because a true prophylactic (a killer of sporozoites) does not exist.

Malaria does occur in some people despite the regular ingestion of daily proguanil or weekly chloroquine. The risk of retinal damage is greater if chloroquine is taken at more frequent intervals. One well-known expert in tropical medicine developed malaria despite daily proguanil but was protected by daily proguanil on weekdays and chloroquine at

weekends. Many experts do not approve of the use of a long-acting sulphonamide (as in Fansidar) or sulphone in Africa at all, or in other areas for long periods of time. Thus there is a current reluctance to recommend Fansidar or any similar combination in Africa even if chloroquine or proguanil have failed in prophylaxis. Therefore in any prospective study it would seem reasonable to include a regimen comprising both chloroquine and proguanil (for example, proguanil 200 mg daily on weekdays with chloroquine 150 mg daily on weekend days), especially in people who have suffered an attack of malaria despite the regular ingestion of one of the drugs given alone. Sequential chemosuppression with proguanil and chloroquine could also be used in practice.

As noted in our article, Fansidar is effective as a chemosuppressant against malaria when administered every one or two or four weeks. Several studies in chloroquine-resistant areas have indicated that dosing every four weeks gives a high degree of protection^{1,2} and this is supported by the lengthy elimination half-lives of pyrimethamine and sulfadoxine.³ However, in one study in a chloroquine-sensitive area (Africa) fewer breakthroughs occurred when Fansidar was given every week or two weeks rather than every four weeks.⁴ In a recent extensive study in a chloroquine-resistant area Fansidar was very effective when given every two weeks,⁵ but there has been little experience with weekly dosing of Fansidar in a chloroquine-resistant area. People often prefer a firm recommendation, and fortnightly dosing (two tablets in adults) would appear to be the optimum dosage interval for Fansidar. Weekly dosing (one tablet in adults) would probably be equally satisfactory.

Since so many people develop malaria after return from endemic areas current prophylactic practice is obviously inadequate. Much of this malaria is due to failure to take the drugs. However, if people are instructed to take their drugs for eight weeks rather than four weeks then it may be hoped that the drugs will be taken on average for a longer period and the prevalence of malaria reduced. The WHO report mentioned by Professor Peters recommends drug prophylaxis for at least four weeks and preferably for six to eight weeks.

Professor Peters notes that prophylaxis for four weeks after leaving an endemic area is sufficient to prevent falciparum malaria. What is his evidence for that statement? He also notes that continued prophylaxis for eight weeks is needed to prevent vivax breakthroughs. This supports the eight-week recommendation since prophylaxis is intended to prevent both falciparum and vivax malaria.—Ed, *BMJ*.

¹ Lewis, A. N., and Ponnampalam, J. T., *Annals of Tropical Medicine and Parasitology*, 1975, **69**, 1.

² O'Holohan, D. R., and Hugoe-Matthews, J., *Southeast Asian Journal of Tropical Medicine and Hygiene*, 1971, **2**, 164.

³ Peck, C. C., Lewis, A. N., and Joyce, B. E., *Annals of Tropical Medicine and Parasitology*, 1975, **69**, 141.

⁴ Shafiei, A. Z., *Journal of Tropical Medicine and Hygiene*, 1975, **78**, 190.

⁵ Pearlman, E. J., et al, in preparation.

Risks of exotic infections

SIR,—Professor N R Grist's plea for a sense of proportion over the risks of imported infections (4 December, p 1382) is timely. That viruses should be projected in the public mind as species of invisible locusts swarming everywhere in relentless pursuit of unwitting

victims correlates with the media's desire to scare people (Minerva, 20 November, p 1267) but not with more orthodox views of the nature and properties of viruses. Even official pronouncements on safety in laboratories seem to reflect greater experience at the committee than the laboratory bench. Why, with our knowledge of viral characteristics, we should be reverting to an era reminiscent of the red flag for the motor car is a mystery. Is the virologist, booted and wrapped in his swaddling clothes and with hands numbed by multilayered gloves, any more fitted to do cultural manipulations than is the surgeon to operate with one hand tied behind his back? It does not appear so.

Of course viruses, being infectious, are capable of the unexpected. Yet examples of lack of secondary spread in the face of numerous opportunities of exposure to persons acutely ill with virus infections are commonplace. The later reports of the recent outbreak of Marburg-like disease in Africa indicate a more limited spread even within families than was at first thought. The precipitate evacuation of some 60 patients from Coppett's Wood Hospital because of the presence of the unfortunate scientist from Porton was therefore startling.

Fascination with safety is laudable. Like crossing a busy road, a code is one thing; its implementation, however, does need trust, not exasperation.

A D MACRAE

Public Health Laboratory,
City and Sherwood Hospitals,
Nottingham

Comparative diagnostic accuracy of barium meal and endoscopy

SIR,—We were interested to read the paper by Dr G W Stevenson and others (25 September, p 723) on the comparison of double-contrast barium-meal examination and fiberoptic endoscopy in acute upper gastrointestinal haemorrhage and the subsequent correspondence from Dr M W Dronfield and others (23 October, p 1010) and Dr P B Cotton (13 November, p 1197). We would like to make a number of comments on the points that they have raised.

The paper by Dr Stevenson and his colleagues might lead many of your readers to believe that endoscopy had made a precise and accurate diagnosis of the bleeding source in 89% of their cases. We would suggest that this is not a proper interpretation of their results. In our opinion¹ the demonstration of a non-bleeding lesion is not acceptable as a positive diagnosis and one must, either by endoscopy or barium-meal examination, have evidence that a lesion is actively bleeding or has recently been bleeding to constitute a positive diagnosis. Lesions not showing these features are of interest but do not constitute a positive diagnosis. Dr Stevenson and his colleagues comment that active bleeding or adherent clot was seen at endoscopy in 36 of their patients out of a total of 53 and thus in our view their positive diagnostic rate was 68% rather than 89%. This compares very favourably with our own previously published series.¹ We would suggest that the remaining 11 of the 53 patients in whom a lesion was found had non-bleeding lesions only.

With regard to the barium-meal examination they comment that the 11 radiological failures or partial failures occurred in patients

in whom endoscopy showed active bleeding or adherent clot. This suggests that this examination has been accurate in only approximately 25 of the 36 patients who had by our criteria a positive endoscopic diagnosis. This represents a positive rate for the barium-meal examination of approximately 25 of the 53 patients—that is, around 50%. We feel it is most important that physicians and surgeons appreciate that the recognition of a non-bleeding lesion does not mean that that lesion has been the site of blood loss.

Our experience of reviewing deaths in acute upper gastrointestinal haemorrhages has shown that patients who die have severe concomitant disease.² It is therefore of interest that in Dr Stevenson's paper the five deaths occurred in patients who were not suitable for surgery. We have already pointed out that it is unlikely that early endoscopy will benefit these patients² and to this extent we entirely agree with the views expressed by Dr Dronfield and his colleagues. Dr Cotton rightly points out that looking at the overall mortality across the broad spectrum of upper gastrointestinal bleeding may well obscure improvements in one area and adverse trends in another. We entirely agree with him that endoscopy will be essential in therapeutic trials studying the management of particular bleeding lesions. However, this is not a justification for early endoscopy as part of the routine clinical management of every patient with upper gastrointestinal bleeding. Our experience leads us to believe that the establishment of an early and accurate diagnosis in those cases in which surgical intervention is not already contemplated is at best of marginal benefit to the patient. Despite the publications of Hoare³ and Hellers⁴ there is as yet no convincing evidence that early endoscopy improves mortality in this situation.

JOHN A H FORREST

R F A LOGAN

Gastrointestinal and Liver Service,
Royal Infirmary,
Edinburgh

¹ Forrest, J A H, Finlayson, N D C, and Shearman, D J C, *Lancet*, 1974, **2**, 394.

² Logan, R F A, and Finlayson, N D C, *Lancet*, 1976, **1**, 1173.

³ Hoare, A M, *British Medical Journal*, 1975, **1**, 27.

⁴ Hellers, G, and Ihre, T, *Lancet*, 1975, **2**, 1250.

Labelling of foodstuffs for patients with coeliac disease

SIR,—There is a proposal for an EEC council directive (COM(76)107 Final) on the approximation of the laws of member States relating to the labelling, presentation, and advertising of foodstuffs for sale to the ultimate consumer currently being considered in Brussels. The Irish Coeliac Society considered that the directive if adopted would allow foodstuffs to contain gluten without any reference to its presence in the listing of the ingredients. The society was also worried that dietary food directives R 3244/73 would not be in a position to countermand such a directive if adopted.

Accordingly we sought a meeting with the Irish joint committee on secondary legislation of the European communities (the All Party Parliamentary Committee on the EEC). I am pleased to report that following this meeting the committee asked for an amendment of the directive so that food manufacturers would have to print on all packets a statement on whether or not there was any gluten in the product. I am writing this letter in the hope