based on H titres of 1/80 or 1/160, and the medical officer of health was informed that he had an epidemic on his hands. *Salmonella paratyphi* B was subsequently isolated from all four persons. Rapid diagnosis was also helpful in patients living outside the main epidemic area and one of these provided the key to cause of the outbreak: 12 days before onset she had been into the area to play tennis and had drunk a glass of water.

Infection was confirmed by my laboratory in 49 cases, in 24 of which serum was examined. H antibodies were present in all sera taken more than five days after onset, and in two of four taken earlier. In four cases H antibodies to *S. typhi* were also found and diagnosis had to await the result of stool or clot culture. Clot culture was positive in 11 of 14 cases examined in the first week of illness; conventional blood cultures, even if used, could not have been much more successful. Faeces were not available on demand and often reached the laboratory several days later.

The exceptional value of serology in this outbreak is attributable to the fact that most patients were children or young adults, and in only four cases was the significance of a raised H titre to *S. paratyphi* B in doubt. In contrast, food handlers examined serologically often showed evidence of past immunisation; serology would therefore have been of much less value had the outbreak occurred in the adult population.

J G WALLACE

Public Health Laboratory,
Lincoln

¹ George, J T A, et al, *British Medical Journal*, 1972, 3, 208.

Shoes for growing feet

SIR,—I write to support the views of Dr M J Illingworth (11 March, p 651) regarding the difficulty of buying suitable shoes for children.

I have twin daughters aged 7 years who have very slim feet. It is virtually impossible to purchase shoes which fit them properly, and one is usually forced to buy shoes which are slightly short in length because of the excess width. On asking shopkeepers about this they state that we are not isolated in this situation and have to send away frequently for narrow shoe fittings. There has recently been the suggestion that shoe manufacturers should adjust the width scale of fittings to make wider the narrowest fitting at present manufactured. This means that it will become impossible to obtain any narrow shoe fittings of any description.

Surely, in the interests of our children, the medical profession should make its views clear.

ANDREW DUNCAN

City Hospital,
Nottingham

Do people smoke for nicotine?

SIR,—In your leading article on this subject (22 October 1977, p 1041) the nicotine-dependence view of smoking was brought into question. One piece of evidence which is claimed to show that smoking behaviour over a period of years is unrelated to the availability of nicotine from a cigarette is that the consumption of cigarettes has not doubled over the 15-20-year period in which the nicotine yield of cigarettes has dropped by about half.

On a crude self-regulation of nicotine intake view one might indeed expect smokers to compensate for the reduced availability of nicotine per cigarette by increasing the number smoked. The recent release of cigarette consumption figures by the Tobacco Advisory Committee prompted me to take a closer look at the available information on this point.^{1,2}

I take it that the figures central to the argument are those relating to the number of cigarettes consumed per adult smoker, since any figures for overall consumption will confound changes in the individual's smoking habits with those of the population as a whole. The accompanying figure strongly suggests that as the average nicotine delivery of UK cigarettes has declined the annual consumption of manufactured cigarettes per smoker has in fact greatly increased. Data on consumption

Average nicotine delivery and annual consumption per smoker of UK-manufactured cigarettes, 1935-75.

per smoker (based on consumer surveys) seem not to be available until 1949 and information on nicotine delivery is very incomplete. However, if the clear trends shown in cigarette consumption since 1949 are extrapolated back to 1935 it can be seen that over the period in which the average nicotine delivery of cigarettes decreased by 53% the number of cigarettes smoked per annum more than doubled in the case of men and increased over six-fold in women. The rise in cigarette consumption by women is presumably due in part to the increasing social acceptability of such behaviour, but no similar explanation is likely in the case of men.

More complete and reliable information covering the past 10 years also reveals an association between the declining nicotine delivery of cigarettes and increased consumption per smoker. In the period 1965-75 the sales-weighted nicotine levels of cigarettes decreased by 35.1%. Over the same period cigarette consumption per smoker increased by 17.9% in men and 30.7% in women.

The smoker has ways of compensating for reduced nicotine delivery by altering the way he smokes individual cigarettes as well as by increasing total number smoked.^{3,4} In view of this the association between increased cigarette consumption and decreased nicotine delivery is perhaps rather greater than might be expected. Such an association cannot, of course, be used to establish any causal relationship, and a way of explaining the rise in consumption per smoker without reference to nicotine delivery has been suggested. If the number of smokers giving up the habit completely has been drawn more from light than from heavy smokers, heavy smokers will have come to form a larger proportion of the total smoking population and per caput consumption will have increased. However, until this is shown to have happened the association between reduced nicotine delivery and increased consumption will tend rather to support than undermine the nicotine-dependence view of smoking.

ROBERT STEPNEY

Clinical Psychopharmacology Unit,
University of Newcastle upon Tyne

¹ *Statistics of Smoking in the UK*, ed P N Lee. Tobacco Research Council Research Paper No 1, 7th edn, tables 7 and 17. London TRC, 1976.

² Todd, G F, *Changes in Smoking Patterns in the UK*, Tobacco Research Council Occasional Paper No. 1. London, TRC, 1975.

³ Ashton, H, and Watson, D W, *British Medical Journal*, 1970, 3, 679.

⁴ Stepney, R, *British Journal of Clinical Pharmacology*, 1977, 4, 653p.

Gastrointestinal bleeding in acute respiratory failure

SIR,—Your leading article (4 March, p 531) left many things unsaid. Stress ulceration is commonly found in patients who require artificial ventilation of the lungs, especially when this is prolonged, and seems to be identical with that found in burns, trauma, etc. It has a high mortality and in 1938 Harkins¹ reviewed all the published reports of Curling's ulcer he could find and reported 107 patients, only 13 of whom had recovered.

Because stress ulceration occurred so commonly among battle casualties during the Vietnam war the American National Research Council held a conference on the subject.² They could come to no clear conclusion as to its aetiology. It has, however, become evident that stress ulceration can develop in the absence of any free hydrochloric acid (HCl) and that a low titratable acidity of the gastric contents and high pH is by no means an uncommon finding.³⁻⁶ The pH of the gastric contents before massive haemorrhage can be as high as 7.0 and is frequently 5.6 or 6.0.⁶

Patients who have undergone open heart surgery and develop postoperative respiratory failure are known to suffer from stress ulceration. A continuous infusion of sodium bicarbonate, either in water or in milk, through a nasogastric tube into the stomach maintaining the pH above 7.0 has been ineffective. When in America I measured the pH of the gastric fluid and found that when it had a pH of 2.0 or less it was sterile, even after it had been discarded and left exposed for a long period. If the pH was 5.0 it contained numerous bacteria. Patients whose gastric fluid contained bacteria were prone to develop stress ulceration. Those patients whose gastric fluid had no bacterial content did not suffer from this problem. Numerous organisms were found and it appeared that the lack of acidity in the