CORRESPONDENCE

Effect of maternal diazepam on the newborn R J Rowlatt, MRCP.....	985	Cigarette smoking and spontaneous abortion J Murphy, MRCOG, and R Mulcahy, FRCP... ..	988	Devolution and the BMA E R C Walker, FRCPED; A M K Barron, MRCPGLAS.....	990
Reactions after pertussis vaccine L H Collier, FRCPATH.....	985	Genital herpes and cervical carcinoma J S Cornes, FRCPATH; J K Oates, FRCPED; Constance A C Ross, FRCPATH.....	988	Hospital day nurseries Sheila C Hollins, MRCPsych, and others....	991
"Human Growth and its Disorders" W A Marshall, MB.....	986	SI, moles, and drugs A C Hutchinson, FIMECHE.....	989	Subconsultant grade I M Librach, MB.....	991
Beta-blockers and plasma triglycerides J Shaw, and others.....	986	HLA antigens in polymyalgia rheumatica P C Mattingly, MRCP, and others.....	989	Payment of district management teams J S Gilmore, MRCP.....	991
Price of survival in childhood leukaemia P G F Swift, MRCP, and others.....	986	Accuracy of data based on hospital discharge diagnoses A J R Scott-Samuel, MFCM.....	990	Dental emergencies and the GP A Hough, MB.....	991
Transmission of agent of Creutzfeldt-Jakob disease D H Adams, DPHIL, and W M Edgar, PHD..	987	Overdose of Lomotil M A McGuigan, MD, and F H Lovejoy, jun, MD.....	990	Distinction awards M Rosen, FFARCS, and M D A Vickers, FFARCS.....	991
Cutaneous reactions to labetalol A Y Finlay, MRCP, and E Waddington, FRCP; Ruth L Savage, MB, and others.....	987	Drug compliance in diabetics A Melander, MD.....	990	Hospital practitioner grade A B Shrank, FRCP.....	991
College of Anaesthetists J A T Duncan, FFARCS.....	987	Foreign intervention in the private sector J W W Studd, MRCOG.....	990	Points from letters Coughing and huffing (H E Schablin; P Skrabanek); Who discovered the circulation of the blood? (S Al-Dabbagh); Postcoital test (J J Slome); Posts in the Middle East (M Doddridge); Errant suffix (J M Henderson)	992
Blood transfusions and renal allograft survival D P S Sengar, PHD, and A Rashid, FRCP(C)...	988				

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters must be signed personally by all their authors.

Effect of maternal diazepam on the newborn

SIR,—Diazepam seems to be the favourite sedative among obstetricians in the management of mild pre-eclamptic toxæmia according to the survey conducted by Mr G V P Chamberlain and his colleagues (11 March, p 626). May I point out what diazepam does to the newborn baby?

High doses (30 mg or more) of diazepam administered during labour cause, in the infant, failure to start breathing, shallow, inadequate respirations, periodic cessation of respiration, floppiness, subnormal temperature, and poor sucking.¹⁻³ These effects last several days and significant plasma levels of diazepam and of its active metabolites persist for up to eight days.¹ Diazepam accumulates in the tissues of the fetus,⁴ and is metabolised and excreted slowly by the newborn baby. It is therefore reasonable to assume that chronic administration of diazepam given in therapeutic doses could cause similar effects, and indeed there is some evidence that this is so.⁵⁻⁷ Symptoms from withdrawal of diazepam have been described in the newborn.⁸

The popular hypnotic nitrazepam is closely related to diazepam and could have similar effects.⁶ The depressant effects of pethidine and other drugs given during labour would be made worse by diazepam; such depression is not always obvious from the relatively crude Apgar score. Diazepam is excreted in breast milk,⁹ which may sedate the baby and cause feeding difficulty.¹⁰ Finally, there is the fear of impairing future intellectual development by exposing the developing brain to the influence of tranquillisers.

Obstetricians are already under attack for embracing new developments without proper

prospective trials.¹¹ How often must we be reminded of the ancient precept "primum non nocere"?

R J ROWLATT

Paediatric Department,
North Lonsdale Hospital,
Barrow-in-Furness, Cumbria

¹ Cree, J E, Meyer, J, and Hailey, D M, *British Medical Journal*, 1977, **4**, 251.

² McCarthy, G T, O'Connell, B, and Robinson, A E, *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1973, **80**, 349.

³ Owen, J R, Irani, S F, and Blair, A W, *Archives of Disease in Childhood*, 1972, **47**, 107.

⁴ Erkkola, R, Kanto, J, and Selman, R, *Acta Obstetrica et Gynecologica Scandinavica*, 1974, **53**, 135.

⁵ Gillberg, C, *Lancet*, 1977, **2**, 244.

⁶ Speight, A N P, *Lancet*, 1977, **2**, 878.

⁷ Kanto, J, Erkkola, R, and Selman, R, *Annals of Clinical Research*, 1973, **5**, 375.

⁸ Rementeria, J C, and Bhatt, K, *Journal of Pediatrics*, 1977, **90**, 123.

⁹ Ayd, F J, *International Drug Therapy News Letter*, 1973, **8**, 38.

¹⁰ Patrick, M J, Tillstone, W J, and Reavy, P, *Lancet*, 1972, **1**, 542.

¹¹ Chalmers, I, and Richards, M P M, in *Benefits and Hazards of the New Obstetrics*, ed M T Chard and M P M Richards, pp 34-61. London, Heinemann, 1977.

Reactions after pertussis vaccine

SIR,—Dr A H Griffith (1 April, p 809) has provided a comprehensive and careful review of adverse reactions associated with the use of Wellcome pertussis vaccine. He rightly draws attention to the possibility that variations in the vaccines made by different manufacturers may in some way affect the incidence of adverse reactions, but in considering this topic it is important that clear distinctions be drawn between fact and suppositions or

inferences. If this is not done there is a danger of confusing the medical profession still further about the complex technical problems surrounding the manufacture, testing, and use of pertussis vaccines. The differences in the vaccines discussed in the article relate mainly to *efficacy* and *toxicity*: several points call for comment.

(1) Dr Griffith makes the bald statement (p 810), reiterated with emphasis in your leading article in the same issue (p 806), that "Wellcome pertussis vaccine... has been found to be more protective than another manufacturer's vaccine." It so happens that the other vaccine was made by Glaxo; this is no secret, since the names of the manufacturers concerned were published in the report referred to.¹ It is, however, unfortunate to say the least that neither Dr Griffith's paper nor your leading article makes it clear that this report, although published in 1973, was entitled "Efficacy of Whooping Cough Vaccines Used in the United Kingdom before 1968" (my italics). In the 1971 addendum to the *British Pharmacopoeia* the minimum standard of potency was specified as 4 international units per dose; needless to say, all vaccines made in the United Kingdom have since then—if not indeed before—complied with this statutory requirement.

(2) There are implications (pp 810 and 812) that vaccines other than Wellcome do not necessarily contain agglutinin 3, which appears to be important for protection. Again, however, Dr Griffith refers only to papers dating from the 1960s that deal with vaccines made before the value of this antigen was widely recognised. It has for some time been the general practice to include it in pertussis vaccines; the Lister Institute has certainly done so for many years.

(3) Dr Griffith states (p 811) that pertussis vaccines are prepared "either dispersed in liquid medium in large tanks or on the surface of a solid medium containing charcoal" (the Wellcome method). Lister vaccine is prepared by neither of these techniques: the organisms are grown in suspension in 8-litre roller bottles. So far as comparative efficacy is concerned, it is worth mentioning that in the 1953-6 MRC trials a direct comparison was made of a solid-medium and two liquid-medium vaccines, all containing the same

strains of *Bordetella pertussis*.² One of the liquid-medium vaccines (coded V15) was made at the Lister Institute; the other liquid-medium vaccine was coded V14 and the solid-medium vaccine V12. In this field trial the Lister vaccine, V15, performed consistently better than V14, V12 being intermediate in efficacy. In laboratory tests V15 was significantly more potent than both V14 and V12. But having made this point I should not think of using it to suggest that any one method of manufacture is "better" than any other; another trial with different batches of vaccine would possibly give somewhat different results.

(4) Telephone calls to these laboratories following the appearance of Dr Griffith's paper and your leading article indicate that the impression has been created among some that pertussis vaccines other than Wellcome's may be less potent or more toxic. Practitioners should be aware that all such vaccines made in Britain have to meet identical statutory specifications and that no batch of vaccine containing an adsorbed pertussis component may be distributed until a sample has been submitted to the National Institute of Biological Standards and Control for examination before release.

(5) On the question of toxicity, the Wellcome method of growing vaccine on medium containing charcoal may indeed absorb out "bacterial toxins" (p 811); but the association of these putative toxins with adverse reactions in children is, as I am sure Dr Griffith would agree, a very moot point. In this connection—and in any consideration of reactions to pertussis vaccine—it is important to be clear as to what is meant by "toxicity." *B. pertussis* can induce a remarkable range of responses in small animals; some of them appear to be associated with specific components of the organism, but none has as yet been unequivocally linked with reactions in children, which fall into two main groups: first, the sort of local inflammatory response, with or without transient fever and malaise, that can be caused by any bacterial vaccine, including the diphtheria and tetanus toxoids contained in the triple preparation; and second, the more serious reactions such as collapse, persistent screaming, convulsions, and encephalopathy. There is evidence³ that toxicity tests in animals are of some use in predicting the liability of a given batch of vaccine to induce reactions in the first category; but—for the moment begging the question of a cause-and-effect relationship between pertussis vaccine and central nervous system (CNS) reactions—there is at present simply no evidence that the reactions in the second category are determined by the strains of *B. pertussis* used, method of manufacture, or performance in any laboratory test. The comparative rarity of these serious reactions bedevils any attempt to relate them to such variables in pertussis vaccine—if indeed such a relationship exists; in addition to the possibility, referred to by Dr Griffith, that some reactions may be purely coincidental, susceptibility to convulsions or other CNS complications could be determined by factors pertaining to a particular child rather than to a particular vaccine. Again, such reactions might not be specific to pertussis; were it customary to inject large numbers of infants with vaccines made from other Gram-negative organisms such as cholera or typhoid, might not similar accidents occur from time to time?⁴

I share Dr Griffith's concern about the difficulty of obtaining reports of untoward reactions and about the poor quality of some of the information that is received. In view of the low incidence of major adverse reactions to pertussis vaccine every case is potentially valuable as a source of information and should be adequately recorded by a central agency; and means should be found of making such records available to manufacturers. For their part, manufacturers should obtain and record as much information as is practicable about the vaccines they distribute. It is the collation of the data from the field with those from the laboratory that provides the major challenge; this is an outstanding example of a problem in which the ability to pose the right questions at

the outset will determine the validity of the answers.

L H COLLIER

Vaccines and Sera Laboratories,
Lister Institute of Preventive Medicine,
Elstree, Herts

¹ Public Health Laboratory Service Whooping Cough Committee and Working Party, *British Medical Journal*, 1973, **1**, 259.

² Report of the MRC Whooping-Cough Committee, *British Medical Journal*, 1959, **1**, 994.

³ Perkins, F T, et al, *Symposia Series in Immunobiological Standardization*, 1970, **13**, 141.

⁴ Cameron, J, *Advances in Applied Microbiology*, 1976, **20**, 57.

"Human Growth and its Disorders"

SIR,—May I draw your readers' attention to an unfortunate error in my book "Human Growth and its Disorders" published by Academic Press? On p 166 the text states that testosterone oenanthate should be given intravenously. It should, of course, be given intramuscularly. Intravenous injection would be highly dangerous.

Sales of the book have been discontinued pending the insertion of errata and every effort is being made to contact those who have already purchased copies. I should be grateful if any readers who possess copies of the book but have not received correction slips would contact me.

My thanks are due to Dr A Stuart Mason for mentioning this error in your review columns (1 April, p 845).

W A MARSHALL

Department of Human Sciences,
University of Technology,
Loughborough, Leics

Beta-blockers and plasma triglycerides

SIR,—We have been conducting a trial comparing the effects of propranolol (160–240 mg), pindolol (15–30 mg), and atenolol (100 mg) given as a single daily dose for the control of hypertension. A randomised block design was used so that each patient received all of the treatments. We observed small but significant increases in fasting plasma triglyceride levels after four weeks of treatment with these drugs when compared with the placebo phases. These rises were not accompanied by changes in plasma cholesterol or blood glucose levels. Although the rise was greatest with atenolol, the changes were significant with all three drugs (see table). Subsequently we examined the effects of metoprolol in the same group of 17 patients. After a further period on placebo patients were given metoprolol 100 mg daily. Again a small but significant rise in fasting plasma triglyceride levels was observed.

While food and alcohol intake may well account for major fluctuations in triglyceride

levels, the evidence from plasma glucose and insulin determinations, particularly in the latter experiment, indicates that all of the subjects were fasting at the time of blood sampling. From direct questioning and our knowledge of the patients only three are regular heavy alcohol users and the fluctuations in their plasma triglyceride levels observed during these studies did not correlate with periods of heavy alcohol intake.

Previous authors^{1–3} have reported varying changes in plasma triglyceride levels in patients taking β -adrenoceptor antagonists. Most have reported the effect of a single drug and have not compared the effects of different drugs within a group of patients.

In view of the associations between hypertension, coronary artery disease, and blood lipids these findings may have important implications. If plasma triglyceride levels remain elevated during long-term treatment of hypertension with β -adrenoceptor antagonists, do the benefits outweigh the risks?

J SHAW
J D F ENGLAND
A S P HUA

University Department of Medicine,
Royal Melbourne Hospital,
Melbourne, Victoria

¹ Barboriak, J P, and Friedberg, H D, *Atherosclerosis*, 1973, **17**, 31.

² Nilsson, A, Hansson, B-G, and Hökfelt, D, *British Medical Journal*, 1977, **2**, 126.

³ Waal-Manning, H J, *Drugs*, 1976, **11**, suppl 1, p 121.

Price of survival in childhood leukaemia

SIR,—Your leading article (11 February, p 321) emphasises the need for long-term surveillance of children who have completed two or three years of intensive therapy for lymphoblastic leukaemia. This is already standard practice in children who appear to have been cured of solid tumours and there is now substantial evidence for clinically significant growth hormone deficiency in children who have received high-dose (5000–6000 R) irradiation for the treatment of intracranial tumours.¹ These children, however, have often had the associated problems of raised intracranial pressure and neurosurgical intervention.

The effects of leukaemia therapy on growth and the endocrine system is not so clear-cut and the results you quote from the Manchester group of Shalet *et al* have not as yet been substantiated by other centres. Shalet *et al*² have described a blunted growth hormone response to hypoglycaemia rather than growth hormone deficiency except in one girl who received more than the now conventional doses of radiation. The quoted cortisol responses³ (not corticotrophin (ACTH) as suggested in your article) to hypoglycaemia are all within the normal limits and the

Mean fasting plasma triglyceride, plasma cholesterol, and blood glucose concentrations ($\pm$ SEM) and fasting insulin levels in 17 patients during treatment with placebo and beta-blockers

	Placebo	Atenolol	Pindolol	Propranolol	Metoprolol
Triglycerides (mmol/l)	1.35 $\pm$ 0.11	2.19 $\pm$ 0.44*	1.73 $\pm$ 0.19*	1.85 $\pm$ 0.28*	1.79 $\pm$ 0.22*
Cholesterol (mmol/l)	5.5 $\pm$ 0.21	5.6 $\pm$ 0.33	5.8 $\pm$ 0.28	5.7 $\pm$ 0.27	5.6 $\pm$ 0.22
Glucose (mmol/l)	4.3 $\pm$ 0.21	4.1 $\pm$ 0.16	4.0 $\pm$ 0.11	4.1 $\pm$ 0.15	4.1 $\pm$ 0.15
Insulin (mU/l)	12.2	12.5	11.8	12.5	13.3

P < 0.05 by paired *t* test

Conversion: SI to traditional units—Triglycerides: 1 mmol/l $\approx$ 88 mg/100 ml. Cholesterol: 1 mmol/l $\approx$ 39 mg/100 ml. Glucose: 1 mmol/l $\approx$ 18 mg/100 ml.