the nation on these controversial matters? At regional and area level there is feedback that can never exist centrally—not only through Cogwheel but through irate colleagues stomping into offices and beating on desks. Dear Royal Commissioners, how about some devolution for England as well?

DAVID KERR

University Department of Medicine,
Royal Victoria Infirmary,
Newcastle upon Tyne

Neck injuries in seatbelt wearers

SIR,—Mr M S Christian's report (27 November, p 1310) of a larger incidence of soft-tissue injuries to the neck in seatbelt wearers is probably the first time that this has been shown in this comparative way. It is, however, important that this is not interpreted as a greater risk of a broken neck with all that that implies. In fact Burke¹ reports in Victoria, Australia, since compulsory wearing of belts was introduced, a decrease in spinal cord injuries from motor vehicle accidents and an overall decrease in severity of cord damage.

The relationship between neck injury, belts, and head restraints expressed in the article has been corrected by Professor W Gissane and Dr J P Bull (18 December, p 1505) and a similar misunderstanding of the same subject was also corrected by them in a letter to you in 1972.² Owing to the emotive potential behind any association between seatbelts and neck injury, with its connotations of severe disability or death, I feel that medical men should be particularly careful not to confuse issues as their views play such a part in general public education. It is unfortunate that much of the research material on which proper understanding depends is not readily available to most doctors.

The acute hyperextension of the neck caused by the sudden forward acceleration of the torso in a rear-end impact was first named "whiplash" by Gay and Abbott.³ Not until the advent of upper torso restraint with a diagonal strap was the opposite neck movement, acute flexion, encountered. But flexion could never be as extreme as extension owing to contact between chin and chest.

Recent bioengineering research on the neck conducted jointly by the biomedical science department of the General Motors Corporation and the biomechanics research centre of Wayne State University^{4,5} has revealed a further significant factor mitigating against injury in flexion. Mertz summarises this as follows.⁵ "The neck appears to be at least three times stronger in resisting flexion than extension. This conclusion is consistent with the anatomical fact that there are more muscles located posterior to the cervical spine, and thus, these muscles should generate greater forces in resisting flexion than the limited number of prevertebral muscles can generate in resisting extension."

As Mr Christian himself points out, sudden hyperextension (he uses the term "whiplash") only rarely causes fracture. How much less likely therefore should serious neck injury be from acute flexion due to wearing a seatbelt? These considerations, however, pale into insignificance when compared with the devastating effects of head impacts with car interiors both on the brain and, through transmitted forces, on the cervical spine in the absence of belts.

Lastly, since the relationships of the occurrence of soft-tissue neck injuries to frontal and rear-end collision respectively are not known it is not possible to attribute injury to flexion or extension and hence to deduce the role of the seatbelt, when worn.

C P DE FONSEKA

Accident and Emergency Department,
Royal United Hospital,
Bath, Avon

- ¹ Burke, D C, *Medical Journal of Australia*, 1973, 2, 801.
- ² Gissane, W, *British Medical Journal*, 1972, 2, 288.
- ³ Gay, J R, and Abbott, K H, *Journal of the American Medical Association*, 1953, 152, 1968.
- ⁴ Mertz, H J, and Patrick, L M, in *Proceedings of 15th Stapp Car Crash Conference*, p 207. New York, Society of Automotive Engineers, 1972.
- ⁵ Mertz, H J, in *Biomechanics and Its Application to Automotive Design*. Neck Injury, p 24. New York, Society of Automotive Engineers, 1973.

Counselling and the student

SIR,—In his article (20 November, p 1245) Mr Brian Thorne describes his basic stance in counselling as listening to his client in an accepting way and helping him to make his own decisions. These ideas will sound very familiar to dynamic psychiatrists and general practitioners who operate on dynamic principles. Mr Thorne shows no awareness that he is working within a tradition that has been built up over a generation by workers of the older disciplines. How difficult it is, too, to use these principles is illustrated in Mr Thorne's example of his clinical work.

There is a great deal of overlap between the counsellor and dynamic psychiatrist, in both their tasks and clientele. Mr Thorne's student is described as one who "fears intimacy and cannot relate to others except in a dependent and compulsive way that leads to inevitable rejection" and surely (unless the student is exaggerating his difficulties as a result of his depression, a possibility which Mr Thorne does not appear to consider) he is suffering from a severe character disorder; and as such he might be considered for long-term individual or group psychotherapy if seen by a dynamically orientated psychiatrist. In the matter of diagnosis Mr Thorne makes a virtue of not attaching a diagnostic label, and, while there is something to be said for this as a way of preserving awareness of the individuality of the client, the real task is to make an assessment which is as refined and accurate as possible of the forces within the personality of the client and his circumstances which result in his distress. It would be most regrettable if the new profession of counselling were to develop without appreciation of the body of thinking and experience which has already been accumulated.

Many basic problems of diagnosis and therapy cry out for solution which not even the most skilled and learned in the present state of knowledge are able to supply. Psychiatry is already discredited in some quarters because too much is expected of it. The same fate will undoubtedly overtake counselling if vast expectations are aroused. This would be a pity, for a well-trained body of counsellors co-operating with general practitioners and psychiatrists could have the potentiality to be of great value in reducing psychiatric morbidity and in lightening the load of the medical profession.

ISOBEL H HUNTER-BROWN

Psychological Advisory Service,
Student Health Service,
University of Leicester

General practitioner's role in the management of labour

SIR,—Professor Peter Curzen and Dr Ursula M Mountrose have written a clear account of a retrospective survey on their unit (11 December, p 1433). Like the good authors they are they have kept it short and some may miss the more obvious steps in the argument; like good debaters they have slightly overstated their case to make a better argument. May I comment on two aspects of the paper?

No delivery can be considered high- or low-risk until it is well over. This is a retrospective differentiation only and should be kept as such. In using these terms an arbitrary dividing line is drawn, but this is only a point made in a continuous curve of risk. Further, there is no uniformity of risk in the high-risk group but an accumulation of high-risk factors, the precise summation of which is made impossible by lack of measurement and knowledge of many of the risks. Professor Curzen and Dr Mountrose quote in their table a standard list with which few would quarrel but, for example, cigarette-smoking by the mother would have a greater chance of affecting the weight-gestation index of her baby than some of the factors on their list. The level at which high risk is taken varies from one population to another and, perhaps more practically, it relates to the local availability of diagnostic and therapeutic facilities. High risk in Roehampton might be low risk in Stockton.

The authors have excluded from their report women who were induced (37.9%) and who presumably had a greater proportion of the quoted high-risk factors. Of the rest, the apparently low-risk group, 7.1% had a fetal or maternal emergency in labour. These situations came unexpectedly and needed prompt treatment. The general practitioner obstetrician, like his colleague the consultant obstetrician, is not present for every moment of labour. They both need other skilled staff to help deal with such emergencies. This implies that the patient should be in a place where appropriate treatment can be given. Surely the data presented provide an argument for general practitioner obstetricians to book their patients in centres where such staff and facilities are available. Obviously a small number of women live in remote areas, but the majority of patients in England, Wales, and Scotland could be looked after in hospitals equipped for this task and not in isolated units. Let any practitioner who doubts this tell a pregnant woman booked with him that even though she is thought to be at low risk there is a greater than 1 in 14 chance of her putting her life or that of her unborn child at hazard by not being in a well-equipped hospital. The answer may convince the practitioner.

GEOFFREY CHAMBERLAIN

Department of Obstetrics,
Queen Charlotte's Maternity Hospital,
London W6

SIR,—Professor Peter Curzen and Dr Ursula M Mountrose (11 December, p 1433) conclude that there is no place in modern hospital obstetric practice for the independent general practitioner. Unfortunately, the results they present show no evidence to support this conclusion. The results show that 7% of low-risk mothers develop problems during labour. What they did not show was, firstly, what pro-