no complications. An abnormal pancreatogram absolutely defines pancreatic disease. There are some difficulties in interpretation of very minor changes and in distinguishing obstruction due to cancer or pancreatitis, at least when clinical information is withheld. A normal pancreatogram does not completely exclude cancer. We have seen one patient with a normal main duct radiograph (with poor branch filling) among 55 cases of pancreatic cancer. Equally, a normal pancreatogram does not exclude an early stage of chronic pancreatitis, as judged histologically. Here, however, the clinical picture is usually one of recurrent pain with a raised serum amylase. The diagnosis is already made and a normal pancreatogram, far from being misleading, is helpful in protecting the patient from pancreatic surgery. Conversely, an abnormal pancreatogram provides a map on which to plan surgical triumphs.

When faced with a patient with obscure upper abdominal pain we are concerned for the diagnosis and exclusion of peptic ulcer disease, gall stones, gastric and pancreatic cancer, and chronic pancreatitis. In this context ERCP (which includes gastroduodenoscopy, retrograde cholangiography, and pancreatography) is a powerful tool.

P B COTTON
R B STERN

Gastrointestinal Unit,
Middlesex Hospital,
London W1

Lomotil in acute diarrhoea

SIR,—In his article on this subject (20 November, p 1240) Dr D R Bell has given a paragraph to diphenoxylate with atropine (Lomotil) which may be misleading.

As he so rightly says, Lomotil is popular nowadays. This is because it is found to be effective and commonly free from side effects. Like most other drugs of therapeutic value, it has unwanted additional actions, but these are not frequently encountered. Warnings are provided that Lomotil should not be given with sedatives and cerebral depressants, but one wonders where he derived his information that "it can potentiate the effects of alcohol dramatically."

The broad statement that the drug "is no longer recommended for children" is unacceptable. It must be asked, On whose recommendation? How old are the children? What are the facts in support? There are no references at the end of the article. The Committee on Safety of Medicines accept its use in children from 1 year of age.

If the drug has the dangers he implies Dr Bell should have been as precise about dosage as the manufacturers. Equally as important, the drug is not related to morphine but to pethidine and there is no evidence that long-term administration leads to habituation. When overdosage does occur with respiratory depression the drug of choice in treatment is naloxone.

MAURICE COHEN
Medical Director,
Searle Laboratories

High Wycombe, Bucks

*.*We sent a copy of this letter to Dr Bell, whose reply is printed below.—ED, *BMJ*.

SIR,—Of course it is true that Lomotil is commonly free from side effects, but a responsible review of current practice cannot

afford to ignore side effects that do sometimes occur. The statement that Lomotil "can potentiate the effects of alcohol dramatically" is based on personal observation and is intended to warn against giving Lomotil with alcohol or any other central nervous system depressant drug. The manufacturer's literature recommends "caution" under these circumstances; I think my recommendation to avoid such combinations is more helpful.

The use of Lomotil in children below the age of 12 is a matter for the physician's own discretion. My article was intended to reflect the consensus of current medical practice in Britain and I believe I have done this.

That prolonged use of diphenoxylate can cause dependence of the morphine type is stated in Martindale's *Extra Pharmacopoeia*.¹ The same source also refers to a report of coma and cardiac arrest in a 2-year-old child after ingestion of as few as six Lomotil tablets.²

I agree that the manufacturer's 1973 recommendation of nalorphine as the preferred antidote to diphenoxylate poisoning has now been superseded and that naloxone, being free of depressive effects itself, is the drug of choice.

DION BELL

Department of Tropical Medicine,
Liverpool School of Tropical Medicine,
Liverpool

¹ *Martindale: The Extra Pharmacopoeia*, ed N W Blacow, 26th edn, p 1117. London, Pharmaceutical Press, 1972.

² Henderson, W, and Psaila, A, *Lancet*, 1969, **1**, 373.

The hospitals we need

SIR,—Dr R M Emrys-Roberts has summarised the Oxford cost-effectiveness study concisely (4 December, p 1385). The thoroughness of the Oxford experiment deserves wide recognition and it is sad that our colleagues there have had to give the financial kiss of death to their brain-child so early in the life of the community hospital. What is not clear is why the Department of Health and Social Security accepted this experiment as the blueprint for their own discussion document.¹ Experience of some 350 general practitioner hospitals containing some 9000 beds, much of it now fully recorded by Hospital Activity Analysis, has been available for some time. The scope and costs of providing hospital services in small units could usefully have been subjected to open debate before any discussion document was produced.

The main difference between the more active GP hospitals and the Oxford Community Hospital Programme is one of emphasis on the hospitals' acute role. Arbitrary limitation of this aspect of hospital work is likely to lead to a severely limited service to the community in question and subsequent loss of cost-effectiveness. In Brecon the casualty:admissions ratio is 5:1, and the x-ray:admissions ratio is 3:1. The casualty department is the busiest and relatively most efficient in the hospital, with a referral rate to district general hospitals of less than 10%. This compares with an inpatient referral rate of 30% to DGHs from the community as a whole, 70% being treated in Brecon Hospital. Transfer costs for patients in the Oxford area were a significant cause of loss of cost-effectiveness. The more acute services that are provided locally, the less this factor will predominate.

The chief loss in the debate so far has been the potentially useful term "community hospital," which has been elevated to a great

height only to be shot down on financial grounds—the most devastating of all at the moment. The Association of General Practitioner Hospitals is about to produce a document in similar terms to those of the DHSS for further discussion. Could not the name "community hospital" be viewed afresh from the time of its publication?

A J M CAVENAGH

Brecon, Powys

¹ Department of Health and Social Security, *Community Hospital. Their Role and Development in the National Health Service*. London, HMSO, 1974.

SI: two years on—a lesson for the Royal Commission

SIR,—I share the lack of enthusiasm in your leading article (1 January, p 5). Molar concentrations are largely irrelevant to clinical medicine—I can count on my toes the times I have used them in clinical decisions in the past two years. If I want the plasma osmolality I do not add six experimental errors and a few false assumptions: I measure it.

New graduates acquire facility with the new system as we did with the old, but for my generation the switch to SI wiped out, at a stroke, 25 years of experience in interpreting results. Two years later I respond emotionally to creatinine and urea in SI units but I still convert dextrose, calcium, phosphate, urate, bilirubin, and blood gases into "old units" before making decisions. This was entirely predictable by anyone who had watched the more necessary introduction of multi-equivalent mEq for the common electrolytes 20 years earlier. We were told that the transition to SI would be quick and we would soon be wondering what all the fuss was about, but for those of us who have 10-20 years' clinical practice ahead and know we will never be as sharp with the new units as we were with the old the transition is pestilentially slow.

At the outset I wrote to Sir George Godber protesting against the new system and urging him at least to preserve the mm Hg for blood pressure. I received a bland reply from one of his colleagues saying there were no major problems with the changeover; I could only conclude that he never treated patients or read the *BMJ* and the *Lancet*. Had he been brain-washed by a handful of biochemists and failed to realise that laboratories can express their results in eils and furlongs at the touch of an electronic calculator, but they are interpreted by clinicians? We were assured that 20+ official bodies had been consulted, but the Department of Health and Social Security, having created Cogwheel, never used its own system to sound the opinion of those who would have to use SI in their daily work.

It is this lack of contact between the Elephant and Castle and the front line of medicine that I hope the Royal Commission will notice. Some decisions (including the future of SI) have to be made centrally but a lot more could be devolved, provided resources are shared fairly between regions and cash limits are applied impartially across the country. If these objectives are achieved is it really necessary for the minutiae of building plans, painstakingly drawn up with local consultation, to be rescrambled in London? Indeed, does the DHSS need to be involved at all in decisions about the siting and size of hospitals or health centres? Need we have uniform policies across

the nation on these controversial matters? At regional and area level there is feedback that can never exist centrally—not only through Cogwheel but through irate colleagues stomping into offices and beating on desks. Dear Royal Commissioners, how about some devolution for England as well?

DAVID KERR

University Department of Medicine,
Royal Victoria Infirmary,
Newcastle upon Tyne

Neck injuries in seatbelt wearers

SIR,—Mr M S Christian's report (27 November, p 1310) of a larger incidence of soft-tissue injuries to the neck in seatbelt wearers is probably the first time that this has been shown in this comparative way. It is, however, important that this is not interpreted as a greater risk of a broken neck with all that that implies. In fact Burke¹ reports in Victoria, Australia, since compulsory wearing of belts was introduced, a decrease in spinal cord injuries from motor vehicle accidents and an overall decrease in severity of cord damage.

The relationship between neck injury, belts, and head restraints expressed in the article has been corrected by Professor W Gissane and Dr J P Bull (18 December, p 1505) and a similar misunderstanding of the same subject was also corrected by them in a letter to you in 1972.² Owing to the emotive potential behind any association between seatbelts and neck injury, with its connotations of severe disability or death, I feel that medical men should be particularly careful not to confuse issues as their views play such a part in general public education. It is unfortunate that much of the research material on which proper understanding depends is not readily available to most doctors.

The acute hyperextension of the neck caused by the sudden forward acceleration of the torso in a rear-end impact was first named "whiplash" by Gay and Abbott.³ Not until the advent of upper torso restraint with a diagonal strap was the opposite neck movement, acute flexion, encountered. But flexion could never be as extreme as extension owing to contact between chin and chest.

Recent bioengineering research on the neck conducted jointly by the biomedical science department of the General Motors Corporation and the biomechanics research centre of Wayne State University^{4,5} has revealed a further significant factor mitigating against injury in flexion. Mertz summarises this as follows.⁵ "The neck appears to be at least three times stronger in resisting flexion than extension. This conclusion is consistent with the anatomical fact that there are more muscles located posterior to the cervical spine, and thus, these muscles should generate greater forces in resisting flexion than the limited number of prevertebral muscles can generate in resisting extension."

As Mr Christian himself points out, sudden hyperextension (he uses the term "whiplash") only rarely causes fracture. How much less likely therefore should serious neck injury be from acute flexion due to wearing a seatbelt? These considerations, however, pale into insignificance when compared with the devastating effects of head impacts with car interiors both on the brain and, through transmitted forces, on the cervical spine in the absence of belts.

Lastly, since the relationships of the occurrence of soft-tissue neck injuries to frontal and rear-end collision respectively are not known it is not possible to attribute injury to flexion or extension and hence to deduce the role of the seatbelt, when worn.

C P DE FONSEKA

Accident and Emergency Department,
Royal United Hospital,
Bath, Avon

- ¹ Burke, D C, *Medical Journal of Australia*, 1973, 2, 801.
- ² Gissane, W, *British Medical Journal*, 1972, 2, 288.
- ³ Gay, J R, and Abbott, K H, *Journal of the American Medical Association*, 1953, 152, 1968.
- ⁴ Mertz, H J, and Patrick, L M, in *Proceedings of 15th Stapp Car Crash Conference*, p 207. New York, Society of Automotive Engineers, 1972.
- ⁵ Mertz, H J, in *Biomechanics and Its Application to Automotive Design*. Neck Injury, p 24. New York, Society of Automotive Engineers, 1973.

Counselling and the student

SIR,—In his article (20 November, p 1245) Mr Brian Thorne describes his basic stance in counselling as listening to his client in an accepting way and helping him to make his own decisions. These ideas will sound very familiar to dynamic psychiatrists and general practitioners who operate on dynamic principles. Mr Thorne shows no awareness that he is working within a tradition that has been built up over a generation by workers of the older disciplines. How difficult it is, too, to use these principles is illustrated in Mr Thorne's example of his clinical work.

There is a great deal of overlap between the counsellor and dynamic psychiatrist, in both their tasks and clientele. Mr Thorne's student is described as one who "fears intimacy and cannot relate to others except in a dependent and compulsive way that leads to inevitable rejection" and surely (unless the student is exaggerating his difficulties as a result of his depression, a possibility which Mr Thorne does not appear to consider) he is suffering from a severe character disorder; and as such he might be considered for long-term individual or group psychotherapy if seen by a dynamically orientated psychiatrist. In the matter of diagnosis Mr Thorne makes a virtue of not attaching a diagnostic label, and, while there is something to be said for this as a way of preserving awareness of the individuality of the client, the real task is to make an assessment which is as refined and accurate as possible of the forces within the personality of the client and his circumstances which result in his distress. It would be most regrettable if the new profession of counselling were to develop without appreciation of the body of thinking and experience which has already been accumulated.

Many basic problems of diagnosis and therapy cry out for solution which not even the most skilled and learned in the present state of knowledge are able to supply. Psychiatry is already discredited in some quarters because too much is expected of it. The same fate will undoubtedly overtake counselling if vast expectations are aroused. This would be a pity, for a well-trained body of counsellors co-operating with general practitioners and psychiatrists could have the potentiality to be of great value in reducing psychiatric morbidity and in lightening the load of the medical profession.

ISOBEL H HUNTER-BROWN

Psychological Advisory Service,
Student Health Service,
University of Leicester

General practitioner's role in the management of labour

SIR,—Professor Peter Curzen and Dr Ursula M Mountrose have written a clear account of a retrospective survey on their unit (11 December, p 1433). Like the good authors they are they have kept it short and some may miss the more obvious steps in the argument; like good debaters they have slightly overstated their case to make a better argument. May I comment on two aspects of the paper?

No delivery can be considered high- or low-risk until it is well over. This is a retrospective differentiation only and should be kept as such. In using these terms an arbitrary dividing line is drawn, but this is only a point made in a continuous curve of risk. Further, there is no uniformity of risk in the high-risk group but an accumulation of high-risk factors, the precise summation of which is made impossible by lack of measurement and knowledge of many of the risks. Professor Curzen and Dr Mountrose quote in their table a standard list with which few would quarrel but, for example, cigarette-smoking by the mother would have a greater chance of affecting the weight-gestation index of her baby than some of the factors on their list. The level at which high risk is taken varies from one population to another and, perhaps more practically, it relates to the local availability of diagnostic and therapeutic facilities. High risk in Roehampton might be low risk in Stockton.

The authors have excluded from their report women who were induced (37.9%) and who presumably had a greater proportion of the quoted high-risk factors. Of the rest, the apparently low-risk group, 7.1% had a fetal or maternal emergency in labour. These situations came unexpectedly and needed prompt treatment. The general practitioner obstetrician, like his colleague the consultant obstetrician, is not present for every moment of labour. They both need other skilled staff to help deal with such emergencies. This implies that the patient should be in a place where appropriate treatment can be given. Surely the data presented provide an argument for general practitioner obstetricians to book their patients in centres where such staff and facilities are available. Obviously a small number of women live in remote areas, but the majority of patients in England, Wales, and Scotland could be looked after in hospitals equipped for this task and not in isolated units. Let any practitioner who doubts this tell a pregnant woman booked with him that even though she is thought to be at low risk there is a greater than 1 in 14 chance of her putting her life or that of her unborn child at hazard by not being in a well-equipped hospital. The answer may convince the practitioner.

GEOFFREY CHAMBERLAIN

Department of Obstetrics,
Queen Charlotte's Maternity Hospital,
London W6

SIR,—Professor Peter Curzen and Dr Ursula M Mountrose (11 December, p 1433) conclude that there is no place in modern hospital obstetric practice for the independent general practitioner. Unfortunately, the results they present show no evidence to support this conclusion. The results show that 7% of low-risk mothers develop problems during labour. What they did not show was, firstly, what pro-