(7) Contrary to much practice streptomycin has been relegated to deal with "difficult" patients, ethionamide has been placed in "reserve," and thiacetazone, much used in the developing countries, has not been mentioned.

(8) Perhaps most significant is the failure to discuss the importance of monitoring the patient's progress through sputum cultures and drug sensitivity testing.

Dr Seaton rightly notes some of the pitfalls met with in the treatment of individual patients. However, even the most affluent countries might have difficulty in justifying the use of expensive chest physician time to check on the accuracy of the pharmacist or the drug handling of the patient. It is the community nurse or auxiliary health worker who should play the primary role in the treatment of all ambulatory patients, not just the vagrants and alcoholics.

DEREK ROBINSON

Liverpool School of Tropical Medicine,
Liverpool

¹ WHO Expert Committee on Tuberculosis, *Ninth Report*, Technical Report Series No 522. Geneva, WHO, 1974.

² Sewell, E M, O'Hara, D O, and Kendig, E L, *Pediatrics*, 1974, **54**, 650.

³ Evans, D A P, *Annals of the New York Academy of Sciences*, 1968, **151**, 723.

⁴ Center for Disease Control, *Morbidity and Mortality Weekly Report*, 1971, **20**, 231.

⁵ Springett, V H, *Practitioner*, 1975, **215**, 480.

Pneumonia

SIR,—Dr Colin M Ogilvie starts off well when discussing the management of pneumonia (25 March, p 771), but how I wish he had consulted an experienced general practitioner before he sent you the completed manuscript. To take 12 ml of blood for four separate containers before giving antibiotics is a counsel of perfection which is unlikely to be carried out routinely in many hospitals, let alone in general practice, where few doctors carry syringes larger than 10 ml. How many bacteriologists would be pleased to receive all these specimens as well as sputum samples from every case of pneumonia in their catchment area? Dr Ogilvie advises chest radiographs during the illness to rule out underlying lesions. These are difficult to arrange in the home and are more likely to be helpful during convalescence. If amoxycillin is prescribed "thrice daily" 15 h may elapse between the teatime dose and the breakfast dose in some households. Eight-hourly dosage should be advised. Transfer to hospital after fever has persisted for more than four or five days is unnecessary in viral pneumonia unless the patient is toxic. Finally, Dr Ogilvie's first-choice simple analgesic, dihydrocodeine, combined with the codeine of his first-choice linctus, will cause constipation in most patients confined to bed.

DENIS CRADDOCK

South Croydon, Surrey

What kind of cot death?

SIR,—We are probably not the only investigators into sudden death in infancy to have noticed the resemblance between the three types described in section 7 of the 1965 "Enquiry into Sudden Death in Infancy"—"(1) cause of death established, (2) possible cause of death found, and (3) cause of death not established" with those in your leading

article "What kind of cot death?" (18 March, p 671).

That "this heterogeneity reduces the value of the epidemiological facts about cot death that continue to be gathered" is a comment whose validity is better assessed at the conclusion of the studies to which you refer. The fact that "no study has attempted to differentiate its findings for the three clinicopathological categories of cot death" is surely a reason for continuing epidemiological studies until this problem is solved.

One benefit of cot death fact gathering is that the recently bereaved parents are put in touch with skilled personnel with an interest in the death of a specific baby. From comments made to us during our studies and from elsewhere these interviews both comfort and relieve—activities which we regard still as important in late 20th century medicine. Furthermore, those engaged in community health services themselves benefit by participating in prospective studies, for in this way they become more aware of the possibilities for prevention described in the article.

We agree that, for reasons not entirely clear but which may be related to the fact of a much greater awareness of the vulnerability of certain categories of baby, the infant death rate is falling. In our study we have been heartened to find that in one of the four boroughs concerned there has been no instance of cot death in the year just ended at a time when the birth rate was rising.

In our view this sort of study should be encouraged in those areas in which it is not yet practised, for the benefits are bound to outweigh the organisational problems. Not least of those benefits is the bringing together of all those who must be involved when a sudden infant death occurs in a manner which indicates to the parents that their particular personal tragedy has evoked concern.

D R CHAMBERS
ELIZABETH WATSON
J M CAMERON

Department of Forensic Medicine,
London Hospital Medical College,
London E1

¹ Ministry of Health, *Reports on Public Health and Medical Subjects*, No 113. London, HMSO, 1965.

SIR,—Your leading article (18 March, p 671) states that the recent reduction in the post-neonatal mortality rate "is hard to explain at a time when the economic climate has been at its worst since the war."

Postneonatal mortality rates are available for England and Wales since 1905. Since then there have been four occasions on which the unemployment rate has more than doubled to over 5% in less than two years: 1907-8, 1920-1, 1929-31, and 1974-6. The average postneonatal rates in England and Wales for the two years immediately preceding the rise in unemployment and the following 2 years were:

83.8 (1906-7) — 74.5 (1908-9)
51.2 (1919-20) — 43.4 (1921-2)
37.8 (1928-9) — 32.0 (1930-1)
5.4 (1973-4) — 4.7 (1975-6)

The percentage decline in each period was 10%, 15%, 15%, and 13% compared with an average decline of 10% every two years since 1905. It appears that unemployment does not slow the rate of decline of the postneonatal mortality rate and may accelerate it.

This phenomenon could be due to a decline

in fertility. The fertility rate (births per 1000 women aged 15-44) for the corresponding periods were:

108 (1906-7) — 105 (1908-9)
86 (1919-20) — 85 (1921-2)
68 (1928-9) — 67 (1930-1)
68 (1973-4) — 63 (1975-6)

There is a decline in fertility in each of these periods but it is consistent with the general decline in the respective period.

It has been known since 1927¹ that the total standard mortality rate in the USA and Britain has tended to decline more rapidly when there is an increase in unemployment. This has been confirmed recently for the USA.² The standard mortality rates for relevant periods were (1930 = 100):

159 (1906-7) — 151 (1908-9)
133 (1919-20) — 126 (1921-2)
121 (1928-9) — 112 (1930-1)
73 (1973-4) — 72 (1975-6)

The percentage decline in each period was 5%, 5%, 7%, and 1% compared with an average decline of 2% every two years since 1905. There is an increased rate of decline on the first three occasions, which Eyer² has argued is due to the "moderate predominance of the stresses associated with migration, community breakdown, and overwork during the boom over the stress associated with unemployment during the depression." One result is that "the suppression of immunity in stress may thus be responsible for the increased population susceptibility to disease prior to epidemics."

A smaller reservoir of adults with infectious disease may explain the earlier acceleration in the rate of decline of the postneonatal mortality rate but does not explain the most recent acceleration.

PETER ELTON

Salford Area Health Authority
(Teaching),
Manchester

¹ Thomas, D S, *Social Aspects of the Business Cycle*. New York, Knopf, 1927.

² Eyer, J, *International Journal of Health Services*, 1977, **1**, 125.

Age and urine concentration after desmopressin

SIR,—We have previously demonstrated a relative impairment of urine concentrating ability in a group of hospital inpatients compared with a younger group of healthy volunteers (7 January, p 24). The differences noted might be attributed to the effects of illness or to deterioration in concentrating capacity with advancing age.

To investigate this further we selected 43 healthy volunteers (31 men and 12 women; age range 20-59 years). Each underwent a urine-concentration test, receiving 40 µg of desmopressin intranasally at 0900 on the day of the test as previously described.¹ Osmolality was measured on all urine specimens passed over the next 9 h. Age groups and mean maximum urine osmolality for each group are shown in the accompanying table.

The mean maximum osmolality for group A was significantly greater than for groups E ($P < 0.01$ by Student's t test), G ($P < 0.02$), and H ($P < 0.05$). Other differences did not reach statistical significance. Results for male and female subjects were similar in each age group.

There was no significant difference between mean maximum osmolality in the 24 healthy subjects aged 40-59 (894 ± 105 mmol/kg) and that previously measured in 11 convalescent