events), the Benton Visual Retention Test, Inglis PALT, Digit Symbol and Block Design Subtests, reaction times, Mill Hill Vocabulary Test, and Raven Progressive Matrices. Computerised axial tomography, serological tests for syphilis, thyroid function tests, serum vitamin B₁₂ and folate estimations, and liver and renal function tests were performed on all patients. Two patients were found to have low folate levels, but correction of this deficiency failed to improve their dementia.

Treatment was begun with bromocriptine 2.5 mg daily and increased by 2.5 mg increments at 4-day intervals to a maximum dose of 20 mg daily. Patients were assessed clinically at fortnightly intervals and after eight weeks of maximum tolerated doses of bromocriptine psychological testing was repeated. Placebo was then substituted for bromocriptine unknown to the patient and a further assessment made after two weeks.

Bromocriptine failed to improve cognitive disabilities. Side effects including confusion and nausea were common, only six patients attaining a dose of 20 mg. The relatives of three patients, however, reported improvement in the patients' ability to cope with their disease as a result of a mild sedative effect. It seems unlikely, therefore, that presently available dopaminergic agonists will benefit the intellectual and memory deficits of presenile dementia. The effects of a dopaminergic receptor agonist cannot be entirely equated with those of levodopa and, as Dr Lewis and his colleagues suggest, further studies with levodopa on more patients with presenile dementia for longer periods would seem justified.

We are grateful to Sandoz Products Ltd for supplies of bromocriptine.

P PHUAPRADIT
M PHILLIPS
A J LEES
G M STERN

University College Hospital,
London WC1

- ¹ Drachman, A D, and Stahl, S, *Lancet*, 1975, **1**, 809.
² Kristensen, V, Olsen, M, and Theilgaard, A, *Acta Psychiatrica Scandinavica*, 1977, **55**, 41.
³ Loranger, A W, et al, *Archives of General Psychiatry*, 1972, **26**, 163.

VD journals

SIR,—I note that in your leading article on genitourinary medicine (25 March, p 740) you refer to the *British Journal of Venereal Diseases* as the only specialist journal dealing exclusively with sexually transmitted diseases.

I think that the *Journal of the American Venereal Disease Association*, started in 1974, also deals exclusively with this subject.

JOYCE WRIGHT

Tillingham, Essex

Exotic diets and the infant

SIR,—Some of the problems in exotic and cult diets could be reduced if doctors and nutritionists made more use of the experience of indigenous communities eating and breeding on well-balanced vegetarian and vegan diets. We already provide such services, in the form of nutritionally assessed recipes, cookery demonstrations, and information. Immigrants in alien circumstances and young parents

wrestling with spiritual challenges but unversed in nutrition might be helped more understandingly than under the stress of a professionally clinical and uncomprehending (and, maybe, even unsympathetic) environment. Such people can honour their beliefs and enjoy the benefits of vegetarianism. Unfortunately many of the victims of malnutrition are suffering from the corrupting influences of Western junk-foods, which, alas, are demotic.

ALAN LONG
Honorary Research Adviser
Vegetarian Society of the
United Kingdom

London W8

Looking inside arteries

SIR,—There are two points in your leading article (18 February, p 389) which require comment. You are right to take a cautious and critical view of endoscopy in vascular surgery.

The paper by Towne and Bernhard¹ describes the use of a Hopkins choledochoscope for inspecting the end result of various vascular operations. Nearly half of these were carotid endarterectomies. The additional time taken was 1-5 min in 53 cases and 5-10 min in 29 cases, not important perhaps in aortoiliac procedures but probably an unjustified extension of carotid occlusion time in carotid endarterectomy. The return for this was the finding of atheromatous shreds in the internal carotid artery in two patients, but most would consider that these can be adequately recognised and removed before arteriotomy closure. The 40% incidence of distal intimal flaps in the external carotid artery can hardly be considered important in practice. Vollmar, the originator of this technique, used it mainly for inspection of arteries after closed endarterectomy, particularly in the iliac and femoral regions. Endarterectomy of the femoropopliteal segment is now rarely performed by most vascular surgeons as the results of femoropopliteal vein bypass, erroneously called saphenofemoral bypass in your article, are much better and there is little indication from Towne and Bernhard's paper that endoscopy will help in the construction of the distal anastomosis, the principal problem in this operation. Likewise, closed aortoiliac endarterectomy is being performed less frequently because of the superior results of aortofemoral grafting and it is hard to see how endoscopy could help in this procedure. Perhaps this technique might be useful for inspection of thrombectomised aortofemoral graft limbs, when it is often difficult to remove adequately the adherent neointima that forms in all prosthetic grafts. However, I suspect the authors have their doubts about the technique—expressed in the paper's subtitle: "Useful tool or interesting toy."

The second point in the article concerns available methods for determining technical faults in vascular reconstructions. In this respect, as you state, the flowmeter has little to offer as there are many other factors which govern the flow in femoropopliteal vein grafts. However, it may have a predictive value. Also flowmeters give unreliable results when used on prosthetic materials, particularly the new expanded polytetrafluoroethylene (PTFE), where the flowmeter is virtually useless.

Arteriography is certainly useful, though it adds to the length of the procedure and fails

to show many intimal flaps because it visualises one plane only. Also it is questionable whether correction of all abnormalities seen actually improves graft patency, as re-exploration produces its own problems. One method not mentioned in your article is the pulse volume recorder (PVR). This is simply a segmental plethysmographic method similar to oscillography. It is a simple, non-invasive method which requires a cuff placed on the limb distal to the reconstruction; monitoring can be continued in the recovery room postoperatively. A comparison of the preoperative and postreconstruction PVR tracing gives a good indication of the functional result achieved. It is particularly valuable when there is disease distal to the reconstruction where peripheral pulses are unlikely to be felt. Experience at this hospital suggests that the best compromise is to use the PVR in all reconstructions and to reserve angiography for technically difficult reconstructions, such as femorotibial bypass grafts, or when the PVR shows a poor response.

ANDREW MAY

Massachusetts General Hospital,
Boston, Massachusetts

- ¹ Towne, J B, and Bernhard, V M, *Surgery*, 1977, **82**, 415.

Treatment of tuberculosis

SIR,—Physicians trained or practising tuberculosis care in other parts of the world may be surprised by some of the emphasis in Dr A Seaton's summary of the current United Kingdom attitude towards this disease (18 March, p 701).

(1) Not least by the assertion that extensive controlled trials enable the practitioner to be dogmatic about treatment.

(2) By the assumption that the effects of *Mycobacterium tuberculosis* and the syndrome of the mycobacterial diseases we call "tuberculosis", including infections with *M bovis* and the so-called "atypical" mycobacteria, are the same thing. As a corollary, the mycoses are among other organisms giving rise to a similar disease.

(3) By the statement that the subdisease actually reviewed should be necessarily treated by a specialist chest physician. The principal contributions to the modern successful cure of the tuberculosis syndrome come not from the supervisory physician but from the laboratory which cultures the sputum and determines the drug sensitivity of the causative organism and from the community workers who ensure that the patient remains under adequate and appropriate therapy and free of side effects.

(4) By the unqualified advice that contacts should be "visited," especially as the final paragraph on the same subject is somewhat confused. For determining tuberculin sensitivity most non-British physicians accept the WHO advice that the intradermal Mantoux test with its quantifiable dose is the procedure of choice.^{1,2}

(5) Isoniazid side effects, especially in slow acetylators, are not rare, as the investigation of liver function at the onset of therapy will confirm.³ Death from isoniazid in the United States has occurred in young adults and is most likely with previous liver damage or in chronic alcoholism.⁴ For all the antituberculous drugs it is most important to warn the patient or family of the possible side effects and to provide immediate review should any occur.

(6) The use of costly rifampicin as a first-line drug, though now widespread, may in the long term prove unwise. It dramatically clears the sputum and it was a useful reserve drug for resistant organisms. Trials do not as yet prove that it has a therapeutically unique ability to shorten treatment. There are many contraindications to its use, including pregnancy.⁵