Of 64 patients with severe menopausal symptoms who were allocated to a four-month trial in which Premarin and placebo were each taken for two months, 44 had severe vasomotor symptoms; a further 56 patients with less severe symptoms were allocated to a longer, 12 month trial in which Premarin and placebo were each taken for six months. It is stressed that neither the patients nor the attending doctors knew which treatment was being given. Graphic rating scales,^{3,4} which are a widely accepted and sensitive method of measuring change, were used to assess the effects of therapy, and thus the argument that a flush count is not an objective way of evaluating therapy cannot apply to these trials.

During placebo therapy, with respect to control values, there was a small, non-significant reduction in hot flushes in patients on the four-month trial, and in patients on the 12-month trial this reduction was just statistically significant ($P < 0.05$). However, during oestrogen therapy, with respect to placebo values, there was a statistically significant reduction in hot flushes in patients on both the four-month ($P < 0.001$) and 12-month ($P < 0.001$) trials.

Furthermore, in the 64 patients on the four-month trial benefits of oestrogen therapy over placebo were observed for 11 additional symptoms. The improvement in such a large number of symptoms is to some extent a domino effect—that is, a reduction in vasomotor symptoms creates a favourable response in a large number of associated symptoms. This could be demonstrated by studying the 20 patients who had severe menopausal symptoms but no hot flushes; in this group significant improvements on oestrogen therapy were observed for only five (vaginal dryness, memory, anxiety, worry about self, and worry about age) of the 11 additional symptoms. However, the improvement in memory and anxiety in this subgroup suggests that the “mental tonic” effect of oestrogens described by Utian⁵ is an entity and is independent of vasomotor symptoms.

The dearth of good placebo studies on the evaluation of oestrogen therapy in the treatment of the menopausal syndrome initiated these trials. Our results indicate that a significant reduction in hot flushes and a significant improvement in wellbeing, both psychological and symptomatic, will be achieved by oestrogen therapy.

STUART CAMPBELL
M WHITEHEAD

Department of Obstetrics and
Gynaecology,
King's College Hospital Medical
School,
London SE5

¹ Campbell, S, et al, in *Management of the Menopausal and Postmenopausal Years*, ed S Campbell, p 149. Lancaster, MTP, 1976.

² Campbell, S, and Whitehead, M, in *Clinics in Obstetrics and Gynaecology*, vol 4, No 1, ed R Greenblatt and J Studd. Philadelphia, Saunders. In press.

³ Guilford, S, *Psychometric Methods*. New York, 1936.

⁴ Lader, M H, and Wing, L, *Physiological Measures, Sedative Drugs and Morbid Anxiety*. Institute of Psychiatry Maudsley Monographs. London, Oxford University Press, 1966.

⁵ Utian, W H, in *Frontiers in Hormone Research: Oestrogens in the Postmenopause*, ed P A van Keep and C Lauritzen, p 74. Basel, Karger, 1975.

Clonidine and facial flushing in rosacea

SIR,—A recent survey in this department¹ showed an association between migraine and rosacea, 44% of patients with rosacea having migraine, compared with only 13% of control subjects. As both conditions are associated with

abnormal vascular reactivity, drugs which are helpful in migraine could be of benefit in rosacea. In that same paper it was suggested that low doses of clonidine might reduce facial flushing. It was therefore decided to perform an oral and topical double-blind trial of clonidine in the treatment of rosacea with particular reference to flushing.

Seventeen patients entered the oral trial, which lasted for six months; all subjects had rosacea with some degree of facial erythema and flushing and were given either clonidine 0.05 mg twice daily or matched placebo tablets, the therapy being crossed over after three months. Twenty rosacea patients in whom flushing was the main feature entered the topical trial, in which 2% clonidine in a cream base was applied to one side of the cheek and the base alone to the other side four times daily for two weeks.

Both oral and topical active therapies were, on the whole, without effect, but five patients in the oral trial did notice an improvement in the frequency and severity of their attacks of facial flushing while on clonidine. In four patients with severe rosacea the oral trial had to be stopped because of a flare-up of the pustular element of their rosacea. In three of these patients this occurred during the three-month clonidine regimen. The lack of effect of the topical therapy could be due either to clonidine as a drug being totally ineffective in this condition or, alternatively, to the base in which the clonidine was incorporated being inappropriate.

From this evidence it would seem that there is no place for clonidine therapy in the routine treatment of rosacea, but it may occasionally be of benefit to patients whose predominant feature is flushing.

W J CUNLIFFE
BARBARA DODMAN
JULIA G BINNER

Department of Dermatology,
General Infirmary,
Leeds

¹ Tan, S G, and Cunliffe, W J, *British Medical Journal*, 1976, 1, 21.

D-Penicillamine and fatal obliterative bronchiolitis

SIR,—As stated by Dr D A Brewerton (18 December, p 1507), the death from obliterative bronchiolitis of four patients with rheumatoid arthritis was reported recently at a meeting of the Heberden Society. Two of these patients had been given D-penicillamine for their arthritis at Westminster Hospital, where some 60 rheumatoid patients had been maintained on the drug, and two had not.

In order to find out whether the same sort of thing was happening elsewhere I wrote to those in charge of 37 clinics asking (a) for the approximate numbers of patients who had received penicillamine for more than six

months, (b) for the mortality (from all causes) among them, and (c) for the numbers of such deaths that were from cardiopulmonary disease, excluding acute myocardial infarction.

The results of this inquiry are summarised in the accompanying table. The relatively high mortality from Wilson's disease and scleroderma is almost entirely the result of failures of the drug to control these diseases. Note also that most of the patients with Wilson's disease or cystinuria have been taking penicillamine for many years (up to 21), whereas most of the others have been on it for less than two years. Of the 15 non-acute pulmonary deaths, only two were definitely attributable to obliterative bronchiolitis and these were the two at Westminster Hospital. This diagnosis remains a possibility in two more, though an unlikely one in the case of the cystinuric patient.

There is therefore no reason to suppose that penicillamine causes death from obliterative bronchiolitis.

This letter has been drafted with the help of those who contributed the data upon which it is based, for which much thanks.

W H LYLE
Medical Director,
Dista Products Ltd

Speke,
Liverpool

Prostatic cancer

SIR,—We agree with Dr A Rostom (16 October, p 942) and Mr E P N O'Donoghue and Mr G D Chisholm (27 November, p 1324) on the need to improve the treatment of carcinoma of the prostate. We also agree that lymphography is not necessarily a sufficiently accurate technique on which to base a therapeutic protocol.¹ We are, however, concerned that clinical trials are apparently being considered for so-called “localised” prostatic carcinoma involving as a basis for therapeutic protocols currently available techniques to fulfil TNM requirements for classification. As with virtually every other solid tumour, the whole problem of prostatic carcinoma is that therapeutically essential staging—that is, the distinction between those patients who have truly localised or regional disease as distinct from those who have disseminated micro-metastases not detectable by current screening methods—is not possible at present. The concept that this therapeutically vital distinction cannot be made is not recent, having been pointed out by Franks in 1956,² and the same problem also applies to other tumours such as lung, breast, and head and neck cancers.

In our opinion any logical trial aimed at improving the survival time in prostatic cancer should compare local or regional techniques such as surgery or radiotherapy with local therapy plus systemic therapy with antitumour drugs. It is a depressing fact that only 10% of the currently available antitumour drugs

Mortality of patients under treatment with D-penicillamine

	Approximate No of patients treated	Deaths (all causes)	Deaths from non-acute cardiopulmonary disease	Deaths from obliterative bronchiolitis
Rheumatoid arthritis ..	2861	64	8	2 (+ 1 uncertain)
Scleroderma ..	76	8	6	0
Wilson's disease ..	300	37	0	0
Cystinuria ..	69	3	1	1 (uncertain)
Other ..	50	0	0	0
Total	3356	112	15	4

have been adequately tested in prostatic cancer.^{3,4} Such little experience with chemotherapy in prostatic cancer is regrettable since it is the only method potentially capable of killing tumour cells disseminated in most areas of the body.

At present, since we are able by staging techniques to detect only cell numbers of greater than 1000 million, the need for a trial comparing local or regional therapy versus local therapy plus systemic therapy with drugs shown to be occasionally effective in prostatic cancer, such as cyclophosphamide, nitrogen mustard, 5-fluorouracil, and possibly adriamycin,^{5,6} is overwhelming. Since it is now known that adjuvant chemotherapy protocols can be given perfectly safely⁷⁻⁹ there is a priority for a logical trial integrating chemotherapy with local treatment to see if the survival time in these cancers can be increased.

L A PRICE

Royal Marsden Hospital,
London SW3

BRIDGET T HILL

Imperial Cancer Research Fund,
London WC2

¹ Gittes, R F, and Chu, T M, *Seminars in Oncology*, 1976, 3, 123.

² Franks, L M, *Lancet*, 1956, 2, 1037.

³ Murphy, G P, et al, *Seminars in Oncology*, 1976, 3, 103.

⁴ Carter, S K, and Soper, W T, *Cancer Treatment Reviews*, 1974, 1, 1.

⁵ De Wys, W D, *Seminars in Oncology*, 1976, 3, 189.

⁶ Wasserman, T H, et al, *Cancer Chemotherapy Reports*, 1975, 6, 399.

⁷ Price, L A, and Hill, B T, *Lancet*, 1976, 2, 1195.

⁸ Hill, B T, and Baserga, R, *Cancer Treatment Reviews*, 1975, 2, 159.

⁹ Price, L A, and Goldie, J H, *British Medical Journal*, 1971, 4, 336.

Use of cardiac pacemakers in Britain

SIR,—The title of your leading article on pacemakers (13 November, p 1158) quotes from *The Merchant of Venice*:

Tell me where is fancy bred
Or in the heart or in the head?
It is engendered in the eyes
With gazing fed, and fancy dies
In the cradle where it lies.

Portia's father knew well the value of the randomised controlled trial to test flights of fancy, but his daughter was equally determined that any trial of her suitors should not be double-blind or random and, having broken the code, she imposed her own choice on unsuspecting Bassanio by her imperious gaze.

Dr Edgar Sowton (p 1182) is equally reluctant to put himself to a valid test and tries to impose his choice upon the NHS. He strongly advocates the doubling or the trebling of the use of cardiac pacemakers at a cost of £4m or £6m for materials alone with larger sums for hospital costs, but he gives little justification for this expenditure except that "there are still many patients with congenital heart block, cardiomyopathy, surgically induced or associated block, or idiopathic [conduction system] fibrosis." How many? What benefit? What costs?

His paper fails to state how he conducted his national study of pacemakers or what was the response rate to his inquiry, he does not state how large was the subsample on which he calculated his 48-h mortality rate, and his list of co-operating centres seems remarkable for its exclusions. He is unable to quote the market percentages of devices used "for commercial reasons," but he is able to quote

actuarial curves of the proportion of functioning pacemakers from a specific company without reporting the methods used to calculate these curves or how intercurrent deaths were handled.

His study of 55 patients at Guy's Hospital who sought driving licences, of whom 52 were successful, merely indicates that the Swansea computer is not programmed to deal with this matter. His inquiry of 16 insurance companies, none of whom had any knowledge of a motor accident claim involving someone with a pacemaker, must have been addressed to Antonio; Shylock could have told him instantly that at standard risks the probability that in a single year none of the 52 patients in his small study would make a claim is less than 1 in 1000.

If there is to be an increase in the number of pacemakers inserted let it be based upon valid cardiological and epidemiological grounds which are tested by valid trials. Let it not be based upon the gossip of the Rialto.

H CAMPBELL

Department of Medical Statistics,
Welsh National School of Medicine,
Cardiff

SIR,—I read with great interest the review by Dr Edgar Sowton (13 November, p 1182). As an anaesthetist I was concerned that in this detailed review, which includes a section on the method of implantation, there is no mention of any provision of either local or general anaesthesia. Let me assure your overseas readers that even in the present state of the country pacemaker implantations in Britain are accompanied by some form of pain relief.

Can Dr Sowton tell us what percentage of the patients reviewed were given general anaesthetics and if the mortality rate was the same with different methods of pain relief?

EDWARD MATHEWS

Anaesthetics Department,
Queen Elizabeth Hospital,
Birmingham

* * * We sent copies of these two letters to Dr Sowton, whose reply is printed below.—
Ed, *BMJ*.

SIR,—Dr Mathews is, of course, quite correct to point out that patients were given pain relief, but any comparison of mortality rates would be invalid since in the more complicated and difficult cases, including those having epicardial systems, the patient would be more likely to have a general anaesthetic.

Professor Campbell has misunderstood the purpose of the paper, in which I provided up-to-date figures of actual pacemaker practice in Britain. There is no attempt to impose increased pacing on the NHS. This must depend upon interpretation of the facts by referring doctors. Most cardiologists and physicians who have experience of pacemaker patients would agree that a double-blind randomised trial of this form of treatment is now unethical in patients with heart block.

Professor Campbell asks how many patients, what benefit, and what costs would be involved in cardiac pacing, but the answers to these three questions are given in the review. The survey was carried out by means of a questionnaire, as noted in the acknowledgments. This

was sent to all major pacing centres (including his own) and to all agents selling pacemakers in Britain. Complete figures were returned on all apparatus sold, but the different manufacturers were naturally reluctant for their own sales to be made known to their competitors. The actuarial curves apply to one company because 100% follow-up was available from nine large centres and because most of the pacemakers used in Britain are supplied by Devices. Similar data were not available for other generators, although smaller samples showed similar performance, as I reported. The curves were calculated by the standard methods widely used in the literature and by many pacemaker manufacturers. The actuarial curves deal with pacemakers, not patients, and if a patient died with a normally functioning pacemaker then the data were included from the time of implantation up to the time of death, as is standard practice. The list of co-operating centres relates only to those which provided data for these actuarial curves of functioning pacemakers and does not include the very large number of other centres which contributed other data.

There was no subsample for calculation of the 48-h mortality rate since all reports from the cardiac centres provided this information and all were included in the calculation.

The problem of driving licences is one of human decision rather than programming the Swansea computer. Professor Campbell has misunderstood the risk of a pacemaker patient making an accident claim since all pacemaker patients in the country who drive are at risk and not merely the 52 in the Guy's survey.

EDGAR SOWTON

Guy's Hospital,
London SE1

Quis whasit?

SIR,—Dr P G T Bye (25 December, p 1560) purports to correct me in relation to one kind of gerundial usage while allowing me to be right in a second. He states that a noun attached to a gerund must *always* be in the possessive. I disagree. English usage is more flexible than that. He twice misquotes me; but no matter. The crucial point is that he cites in his own support Fowler in the Gowers revision. He fails, however, to state that, having given Fowler's view of the matter at issue, Sir Ernest Gowers added, and with emphasis: "The foregoing article is reproduced just as Fowler wrote it, except that some illustrations have been omitted and others shortened. It provoked some controversy. Jespersen . . . vigorously defended the construction condemned by Fowler [the one used by myself that Dr Bye criticises]. He gave numerous examples of its use by famous authors from Swift to Shaw; he made light of the argument that it defied grammatical analysis, and maintained that it represented 'the last step in a long line of development, the earlier steps of which . . . have for centuries been accepted by everybody.'" What did Fowler have to say to this?—simply that he had an "instinctive repugnance" to any deviation from the rigid rule. Alas! instinctive repugnance, even when Fowlerian, is hardly enough. We must move with the times, anyway the centuries.

Sir Ernest goes on to state that the House of Lords itself later supported my Bye-