CORRESPONDENCE

The other crisis of health care

A E Finnigan, MB..... 1211

Psychiatric problems after mastectomy

Tina Morris, BA, and others..... 1211

If I had . . .

J H McKillop, MRCP, and others; J M Last, FRACP; G T Watts, FRCS; C Wayte; H H G Eastcott, FRCS; R D James, MRCP; N G Buchan, FRCSed..... 1212

Diet and heart disease

J N Morris, FRCP, and others; R R O'Moore, MRCPATH, and others..... 1213

Hazards of Diconal

D H Marjot, MRCPsych..... 1214

Whooping cough

I Capstick, FRCGP..... 1214

Management of self-poisoning

D H Wilson, FRCSed; Jane G B Newson-Smith, MB..... 1214

One drug for epilepsy

K D MacRae, PHD; S D Shorvon, MRCP and others..... 1215

Happy to starve

D P Addy, MRCP..... 1215

Safety in laboratories

C H Collins, MB; D R Christie, MB..... 1215

Treatment of paracetamol poisoning

T J Meredith, MRCP, and others..... 1215

What kind of cot death?

A S Cunningham, MD..... 1216

Fetal malformation not associated with Debendox

A D Kohn, MB, and A C Parsons, MRCP; D H A Redford, MB, and I J Lewis, MB... 1216

Effect of maternal diazepam on the newborn

J M B Burn, FFARCS..... 1216

Side effects of cyproheptadine

J Wortsman, FACP, and others..... 1217

Alcohol and cirrhosis

H G Kinnell, MRCPsych..... 1217

Bronchiectasis and cystic fibrosis

A P Norman, FRCP..... 1217

Hyperbaric oxygen

B H Bass, FRCP..... 1217

Payment to clinical members of district management teams

H G Penman, FRCPATH..... 1217

Payment of district management teams

H B Devlin, FRCS..... 1218

Hospital practitioner grade

R Hole, FRCS; A L Hovenden, FRCR.... 1218

Abuse of Form Med 3

A S Urquhart, MRCPG..... 1218

Joint announcement by BMA and Secretaries of State

J R Donaldson, MRCPATH..... 1218

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters must be signed personally by all their authors.

The other crisis of health care

SIR,—I read your leading article "The other crisis of health care" (15 April, p 939) with interest. It is, of course, very difficult for a nation to budget the amount of money spent on its health, and I entirely agree that the problems of the National Health Service are not merely a shortage of money. The French and the Belgians, with whom I spent 12 months on a prolonged study leave, similarly have their problems regarding finance. The French spend approximately one and a half times the British figure on health. Perhaps, however, your readers may be interested to hear what they get for it. The main points are:

(1) The patient has complete freedom of choice to see any doctor he wishes, and can go directly to a specialist if he wants to. This means he can visit a doctor not only in the same town but also in the next *Département* if he so wishes. It seems that in practice 85% approximately stay with the same GP. The patient can also choose his hospital—for example, a Parisian can have his gall bladder removed in Montpellier.

(2) A specialist can invariably be seen within a week for a consultation, and admission to hospital for an operation or investigation takes three to four weeks at the most.

(3) Many doctors are now tending to form groups, but in a different way from Britain. It is now commonplace for a group of GPs to link up with specialists such as psychiatrists, paediatricians, cardiologists, and pathologists. In one practice in Montpellier, for example, dentists, speech therapists, and home nurses also were included in the group.

(4) GPs in particular tend to work very long hours and the consultation rate as compared with the visiting rate is approximately 3 to 2. It is not uncommon for a GP to do 20-25 visits per day. Many GPs work a 12-hour day, including Saturday mornings, there being a rota for Sundays. (The average French GP earns approximately two and a half times the British rate.)

(5) The patient pays for his services, be this to GP, specialist, dentist, chiropodist, or nurse. *Sécurité Sociale* covers 99% of the population and pays 75-80% of the fee and costs. *Aide Sociale* ensures that the poor pay nothing, and there are maximum fees in existence, in addition to a long list of exemptions from any charges for certain diseases. French law also states that a doctor if summoned must attend the patient in an emergency. 50-55% of patients are covered for the rest of the costs by various private *mutualités*. Fees are controlled by the Government.

(6) Dentists provide a 24-hour cover including Sundays and public holidays. Similarly, pharmacies are open by rota day and night. Home nursing is available at any time and home physiotherapy and chiropody are commonplace. There is also an emergency doctor service should one's doctor of choice be unavailable.

(7) For emergencies there is the *Serviced'Assistance Médicale d'Urgence*. SAMU is a casualty service with a difference. It is based at the main teaching hospitals, which are equipped with a fleet of ambulances, helicopters, nurses, and doctors. Teams of doctors go out if necessary to attend to accidents and urgent medical situations, and it is indeed a "flying squad."

(8) Maternity allowance is paid depending on attendance at antenatal clinic for at least three examinations. This may explain why the French perinatal mortality rates and infant mortality rates have been consistently lower than the British for a number of years now.

(9) The French live slightly longer than we do (despite the scourge of alcoholism).

(10) Non-salaried doctors and paramedical workers can live where they wish within France without Government influence or interference.

My impression of medicine in France (and Belgium is very similar) is one of a contented, dedicated, hard-working profession in which the patient "calls the tune." Indeed, a non-salaried doctor gets no income if he sees no patients and has to compete with colleagues for

a livelihood. Though a certain amount of "playing to the gallery" no doubt takes place, it is good medicine and good manners which are the essential requirements. I have never seen a better doctor-patient relationship anywhere. There is no doubt that the French system not only works, but works a great deal better than ours. The average "Frenchman in the Street" is infinitely better off than his British counterpart. Perhaps, however, the Australian system of the bills being sent to the Government would be more morally acceptable.

My final comment is this. If an individual or a nation wishes to spend a lot of money on health why should this be such a bad thing?

A E FINNIGAN

Oxford

Psychiatric problems after mastectomy

SIR,—Dr G P Maguire (15 April, p 693) has drawn attention to the reluctance of doctors to recognise and treat psychiatric problems in patients who have recovered from mastectomy and therefore may be presumed to be "well." Our findings¹ are essentially similar to Maguire's, but the conclusions we have drawn are somewhat different.

In a consecutive series of 160 women (69 with operable breast cancer and 91 with benign breast disease) we assessed prior to and at three, 12, and 24 months following operation the marital, sexual, interpersonal and work adjustment, depression, and personality characteristics by means of rating scales and standard psychological tests. By two years, there were no significant differences in social adjustment between breast cancer patients and benign breast disease controls. However, 22% of cancer patients were rated moderately or severely depressed on the Hamilton Rating Scale for Depression² at two years compared with 8% of benign breast disease controls, and this figure was not related to the presence of clinically advancing disease in the cancer patients. Patients therefore do not experience spontaneous resolution of their

difficulties with the passage of time. Indeed, we are at present examining data from the five-year follow-up of our cohort, and our evidence so far suggests that difficulties may gradually increase in intensity. A number of recent studies have shown similar levels of emotional morbidity (that is, 20-30%) in newly diagnosed cancer patients.³⁻⁶

It is none the less clear that most patients recover remarkably well from traumatic experiences such as mastectomy and diagnosis of cancer. A "blanket" service for all such patients is not only unnecessary but may be unwanted. It is also clear from the studies we have cited and from our own work that clinical symptoms of depression and inability to return to a previous life style are not a "natural" response even to a life event as threatening as a diagnosis of cancer, whatever the prognosis. What is needed is to identify predictors of poor psychosocial outcome following mastectomy. In our own study, patients who failed to recover psychologically or who were unable to resume their social roles fell into clear at-risk categories, detectable at the time of operation:

(1) Patients likely to experience depressive symptoms needing medical attention were those who had clear clinical signs of depression at the time of operation, regardless of their previous psychiatric history.

(2) Patients most likely to experience sexual difficulties were those who were biologically or chronologically (that is, aged about 50) perimenopausal.

(3) Patients likely to remain stressed by anxieties about their diagnosis were those who had a strong component of emotional lability as measured by the Eysenck Personality Inventory Neuroticism Scale in their personalities: that is, the visibly anxious patient with somatic evidence of anxiety.

Thirty-three per cent of our cancer patients were also dissatisfied with the information they received.

No service can adequately assuage private grief at the loss of a valued body part. What is important is that surgeons, physicians, radiotherapists, and general practitioners treating breast cancer patients recognise and themselves treat or refer depression, anxiety, or sexual problems so that patients may resume their previous level of functioning as quickly as possible. Skill in recognition, at least, in these areas should be part of the armoury of any medical practitioner. Creation of a milieu where patients feel free to voice their anxieties will assist this recognition and therefore the provision of appropriate help.

TINA MORRIS
H S GREER
KEITH W PETTINGALE

Faith Courtald Unit for Human
Studies in Cancer,
King's College Hospital,
London SE5

- ¹ Morris, T, Greer, S, and White, P, *Cancer*, 1977, **40**, 2381.
- ² Hamilton, M, *British Journal of Social and Clinical Psychology*, 1967, **6**, 278.
- ³ Schonfield, J, *Journal of Psychosomatic Research*, 1972, **16**, 41.
- ⁴ Weisman, A D, *American Journal of the Medical Sciences*, 1976, **271**, 187.
- ⁵ Winick, L, and Robbins, G F, *Cancer*, 1977, **39**, 478.
- ⁶ Plumb, M M, and Holland, J, *Psychosomatic Medicine*, 1977, **39**, 264.
- ⁷ Eysenck, H J, and Eysenck, S B G, *Manual of the Eysenck Personality Inventory*. London, University of London Press, 1964.

If I had . . .

SIR,—In a recent article in this journal (8 April, p 896), Professor Harold Ellis indicated that if his wife developed clinically early breast cancer he would wish a preoperative skeletal survey to be carried out, presumably to exclude asymptomatic bony metastases. There is now considerable evidence that the x-ray skeletal survey is an insensitive means of detecting bony metastases in breast cancer and that an isotope bone scan is superior.¹⁻³

There is some debate over the exact incidence of scan-detectable bony metastases in clinically early breast cancer (British Breast Group, unpublished data), but such abnormalities have been shown to have grave prognostic significance.^{4,5}

Like Professor Ellis we would wish to exclude occult skeletal metastases if our wives presented with clinically early breast cancer and suggest that the best means at present available for assessing this, both at the time of presentation and during follow-up, is a bone scan using a ^{99m}Tc-labelled phosphate agent. If this is normal no further studies are necessary. If the scan shows abnormalities, then selected x-rays of the abnormal areas should be performed to exclude benign skeletal disease.

JAMES H MCKILLOP
C B WOOD
L H BLUMGART

University Departments of Medicine
and Surgery,
Royal Infirmary, Glasgow

- ¹ Roberts, J G, *et al*, *Lancet*, 1975, **1**, 237.
- ² Gerber, F H, *et al*, *New England Journal of Medicine*, 1977, **297**, 300.
- ³ Citrin, D L, *et al*, *Clinical Radiology*, 1977, **28**, 107.
- ⁴ Galasko, C S B, *British Journal of Surgery*, 1975, **62**, 694.
- ⁵ Citrin, D L, *et al*, *Surgery, Gynecology and Obstetrics*, 1975, **143**, 360.

SIR,—Your new series got off to a promising start with a thoughtful consideration by a distinguished surgeon of his actions and reactions if his wife had breast cancer. I hope that later we may have a woman doctor's essay on the same theme. There are unmistakable sex differences in attitudes to this disease, which may sometimes be ill understood by men. About four years ago, I asked an examination question in a first-year medical course on behavioural science which illustrates this point. I stated the fact that a high proportion of women who discover lumps in the breast delay seeking treatment, sometimes for many months, and asked the students for a brief statement of the possible reasons. The question was answered by virtually all of the 30-odd women in the class and by about an equal number of the men. The men gave the conventional, expected answer (fear of cancer, the hope that the lump would go away, reluctance to leave the family to enter hospital, economic reasons, and guilty feelings that the lump was a punishment for sexual sin—all of this had been mentioned in the class). The women students gave the same reasons with an additional one on which many waxed eloquent. Recall that these women were in their late teens or early 20s. The extra reason they gave was that the current methods of treatment are an assault on a woman's sexual identity. Men can best understand this by considering how they would react if they suspected they had cancer of the penis. Perhaps the treatment of breast cancer, including its psychological implications, will improve when more women graduates enter specialties such as surgery and clinical oncology.

JOHN M LAST

Epidemiology and Community
Medicine Department,
University of Ottawa,
Ottawa, Ontario

SIR,—The criticisms in your issue of 22 April (p 1050) of Professor Harold Ellis and his views on cancer of the breast seem a little

unfair. Professor Ellis was presenting a sound balanced view which is acceptable for most thinking members of the profession. It is, perhaps, unfortunate that he omitted to consider reconstruction of the breast, but this was, perhaps, part of his brief. It surely seems an obsolete approach now to do a lumpectomy if the sole reason is a cosmetic one. It is now widely accepted that it is possible to remove the breast and reconstruct it using an implant, and this would seem a more logical approach. Many surgeons now perform this procedure and your correspondent might be interested to know that there was to be a symposium on the subject at York on 5 May.

GEORGE T WATTS

General Hospital,
Birmingham

SIR,—Surely Dr T B Brewin (22 April, p 1050) rather misses the point about Professor Harold Ellis's essay (8 April, p 896) "If my wife had breast cancer." If a doctor is in the position written about by Professor Ellis, it is only normal that he would be emotional and even irrational about things—he's a human being like everyone else. It is easy to sit back and be rational when you're not personally involved, but as soon as it is yourself or your wife, child, father, or mother involved rationality is not exactly your first concern in the situation. You don't run a series entitled "If I had . . ." just to get people's clinical opinions.

CHRISTOPHER WAYTE

Bath

SIR,—Dr J M S Pearce (15 April, p 969) has put the cart before the horse. If I had a right ocular transient ischaemic attack like his I would put my stethoscope bell on the middle of my right carotid and not beat about the bush in the manner he described. A good GP friend of mine referred himself to me some years ago in just this manner and I operated on his stenosed carotid a couple of days later.

H H G EASTCOTT

London NW1

SIR,—How strange that Professor H A F Dudley (22 April, p 1035) should omit a good radiotherapist from his list of experts to manage his rectal cancer. As long ago as 1963¹ surgeons had recognised the potential in radiotherapy for eradicating local recurrence from this disease. Postoperative radiotherapy was tried and failed (as many had predicted from their experience in irradiating the female pelvis) because the complication rate is too high.

The logical alternative, preoperative radiotherapy, followed by radical excision, seemed a much more attractive proposition from practical as well as radiobiological considerations. The sceptical rectal surgeon could have taken heart from his "commando" colleagues specialising in the head and neck who regularly perform radical neck dissection after high doses merely by getting their timing right. This approach has now been vindicated and a Medical Research Council trial set up. Local recurrence is reduced to zero and no increase in operative morbidity or mortality occurs—indeed the combined operation is said to be easier.²

There remains the objection that his pelvic nodes might have been needlessly irradiated