regrettable that the medical profession's journal should publish a leading article containing comments which are bound to give encouragement to that body of 19th-century opinion in the Department of Health and Social Security prepared to condemn disabled persons, able to drive, to be second-class citizens with their isolation still further confirmed by restricting their mobility to a "solo vehicle" that is cheap; while your repeated reference to a "motorised wheelchair"—in other words, great-grandfather's Bath chair with some form of power—is an insult to disabled drivers at the close of the 20th century when all the wonders of electronics are available to Government to produce a four-wheel car that at the very least is a two-seater. As for your comment on the need for something cheap to ensure its appeal to the DHSS, this makes sorry reading for disabled pensioners who see the Treasury able to afford huge sums to help subsidise ventures in various Marxist countries in direct competition with Britain's economic interest.

Your article gives rightful prominence to the need for recognition by local authorities for house adaptations to suit the disabled, but, although stressing the need for such changes to be "high on any list of sociomedical priorities," it makes no mention of the rights of disabled persons to have the validity of their opinion acknowledged at governmental and local authority levels.

J S CARSLAW

Crediton, Devon

*A four-wheeled two- or four-seater car might be ideal for some disabled drivers, but there is also a need for a form of solo vehicle for people to get about in their immediate neighbourhood. Not everyone is capable of driving a car, and a slower, safer, more stable vehicle would fulfil an undoubted need. Furthermore, there is no possibility of providing expensive and advanced equipment on demand for large numbers of disabled people; cheapness does not imply second rate, but it does mean that the vehicle can be made available to all who need it.—ED, *BMJ*.

Cervical "erosion"

SIR,—Dr M J Goldacre and his colleagues (25 March, p 748) have by careful study confirmed what many gynaecologists have long taken for granted, that the common cervical erosion is not a lesion. The term is also, as they said, misleading and I wish they had gone further and suggested an alternative—what about "mucosal ectopy"? The only things to be said for "erosion" are that it is established by usage and there seems to be no alternative as brief. Unfortunately, prolonged usage has not altered the common meaning of the word for patients, nor indeed for many doctors. Thus patients are frightened by what sounds like an ulcer (and some are aware of the euphemistic use of *that* term) and doctors still frequently refer women without relevant or even any symptoms to gynaecologists because of cervical "erosion."

Some may argue that they would not want their alertness to the possibility of cervical carcinoma dulled by complacency towards any apparent lesion. However, even if they feel the need for a specialist opinion there is no need to frighten their patients by spurious terminology. In any case, in the absence of

bleeding and given normal cytology the possibility of cervical carcinoma can be ignored.

Although endocervical mucosa is exposed owing to ectropion, it is much more commonly exposed owing to differential growth of the underlying parenchyma, which effectively rolls the epithelium outwards from the external os. This is presumably hormone mediated, being commonest and most florid in pregnancy, and also common during oral contraception, as Dr Goldacre and his colleagues showed. "Mucosal ectopy" would seem an appropriate term to describe this condition, but perhaps someone could suggest a better and simpler alternative.

M G R HULL

University Department of Obstetrics
and Gynaecology,
Bristol Maternity Hospital,
Bristol

Complications of radiculography

SIR,—We were interested to read the report by Dr J B Eastwood and his colleagues of bilateral central fracture dislocation of the hips following radiculography using meglumine iocarmate (Dimer-X) (18 March, p 692). Although we have never seen such a dramatic complication as this, the very high percentage of minor but distressing complications, in particular persistent headache, has led us largely to abandon the procedure in favour of ascending lumbar venography.¹ This is a simple and reliable technique which does not involve thecal puncture and which can be used on outpatients. It also gives information about lesions that are some distance from the dura and will, for instance, help in the diagnosis of lateral disc protrusions as well as deciding whether an observed congenital or traumatic abnormality is, in fact, responsible for the symptoms.

The major drawback is gaining familiarity with the radiological appearance, but this can be rapidly achieved, especially as it is not necessary to admit the patient and hence can be contemplated earlier in the management of patients with back pain.

A K CLARKE

Royal National Hospital for
Rheumatic Diseases,
Bath

JOHN P BEAVIS
DUNCAN IRVING
B E W MACE

Lewisham Hospital,
London SE3

¹ Clarke, A K, *et al*, *Rheumatology and Rehabilitation*, 1977, **16**, 83.

Do people smoke for nicotine?

SIR,—Mr R Stepney (8 April, p 922) should beware of trying to use secular changes in cigarette consumption to give information about changes in individual smoking habits. The annual consumption per adult smoker over the 25 years 1950-75 does not describe changes in smoking habits in the same smokers but just a composite effect of all sorts of changes in a radically altering population of smokers. During this period older female smokers were dying off and being replaced by younger, heavier smoking women. At the same time many men were stopping smoking, but these tended to be lighter smokers. Thus the average consumption per smoker inevitably increased, regardless of whether or not the

persisting heavier smokers increased their consumption. Maybe they did. We just don't know.

Observation of small numbers of smokers exposed without their knowledge to changes in nicotine intake does show that abrupt, large reductions of nicotine are followed by an increased consumption, but smaller changes, as from 1.4 mg to 1.0 mg of nicotine per cigarette, are not.¹

C M FLETCHER

London SW10

¹ Freedman, S, and Fletcher, C M, *British Medical Journal*, 1976, **1**, 1427.

Hazard of chemical sympathectomy

SIR,—We feel it would be unfortunate if those who wished to perform chemical sympathectomies were led to believe that the procedure should only be done under radiological control, as suggested by Drs A P Rubin and B R Master (25 March, p 790). Not only is it frequently difficult to organise the use of an image intensifier, but we feel it is unnecessary. Using a method similar to that described by Mehta¹ we have done 287 cases and have not so far encountered any significant sequelae apart from two patients who developed neuritis of the thigh. Whatever technique is employed, it is important for the needle to be deep enough to avoid injecting near somatic nerves or into the substance of the psoas muscle.

Employing Mehta's technique it is virtually impossible to be sufficiently posterior to enter the intervertebral foramen or puncture the dural cuff. Even if one is doubtful about the position of the needle tip many radiologists would feel that Conray 280 is not without hazard if injected into the subarachnoid space.

JAMES M B BURN
L LANGDON

Department of Anaesthetics,
General Hospital,
Southampton

¹ Mehta, M, *Intractable Pain*, p 213. Philadelphia, Saunders, 1973.

Place of ultrasound and radiography in obstetrics

SIR,—I am afraid that Dr H B Meire and his colleagues (8 April, p 882) are not getting the proper benefit from radiographing fetuses if they merely look on the result as a means of measuring the conceptional age of the fetus.

Ultrasonic and radiographic fetal maturity assessments are measuring different aspects of the baby's development, one the body or head size, and the other the bone age. The relationship of these parameters of gestation to each other and to the other parameters has been discussed; the variations in their relationship to each other is an important indication of the presence and severity of fetal disease.¹ Indeed a considerable portion of antenatal care is given over to searching for differences in the various assessments of fetal age. Merely because they are measured in the same units—weeks of gestation—does not indicate that they are measuring the same entity. Radiological and ultrasonic assessments are therefore complementary to each other, and should not be regarded as supplementary.

It was interesting to note that the ratio of ultrasonic scans to radiography in this series

was 10 to 1. This is the same figure as is found in our hospital.

J G B RUSSELL

Saint Mary's Hospital,
Manchester

¹ Russell, J G B, *Radiology in Obstetrics and Antenatal Paediatrics*, p 35. London, Butterworths, 1973.

Non-autoclavable lignocaine laryngeal sprays

SIR,—We were surprised, but gladdened, to read (11 February, p 369) that Penlon Ltd in conjunction with Professor A C Forrester had in fact anticipated by some years our suggestions (10 September, p 680) for the modification of Forrester laryngeal sprays.

However, it should be noted that other manufacturers—for example, British Oxygen Company Ltd, Medishield—still distribute non-autoclavable laryngeal sprays which may well constitute the infection hazard described in our original paper.

MARK CASEWELL
MAURICE DALTON

Department of Microbiology,
St Thomas's Hospital Medical
School,
London SE1

Sarcoid heart disease

SIR,—In 1973¹ you were kind enough to publish a letter from me reporting my interest in collecting cases of suspected cardiac sarcoidosis. As a result of this and other efforts a report² was made of 50 cases. The collection of cases has continued and all surviving patients are followed up by an annual inquiry. It is felt that by this means it may be possible to gain a better knowledge of the natural history of this uncommon and often bizarre form of cardiac disease.

I would like to appeal for further cases to be reported to me. To date a total of over 150 cases have been collected. The diagnosis is certain in over 50, almost certain in another 35. There is necropsy confirmation in 37. It should be emphasised that the diagnosis is to be suspected in any bizarre form of heart disease, especially with difficult and changing dysrhythmias or congestive cardiomyopathy. Florid clinical sarcoidosis in other organs is frequently absent, but some histological evidence of sarcoidosis is essential to the diagnosis.

HUGH A FLEMING

Regional Cardiac Unit,
Papworth Hospital,
Cambridge

¹ Fleming, H A, *British Medical Journal*, 1973, **1**, 174.
² Fleming, H A, *British Heart Journal*, 1974, **36**, 54.

Rubella vaccination and pregnancy

SIR,—We are concerned about the unthinking administration of vaccine to women who wish to be immunised against rubella before they become pregnant. From time to time patients are referred for consideration of termination of pregnancy because of inadvertent administration of vaccine in the second part of the menstrual cycle when, unknowingly, they are pregnant. The decision is always difficult for the gynaecologist because of the relatively slight risk of the fetus being affected. Giles¹ reported that, in 83 women given rubella

vaccine before planned termination of pregnancy, the virus was isolated in three of the fetuses and ten of the placentas. If one assumes that 85% of women are seropositive anyway, and therefore immune, the risk of fetal infection is slight. However, for the woman who finds she is pregnant and knows that the vaccine was given shortly after conception it is impossible to view the matter so objectively, and nearly always she persists in her request for termination.

We would like to make a plea that the immune status of the patient is determined before rubella vaccine is given. If she is seronegative the vaccine should only be given if it is certain that pregnancy can be avoided by adequate contraception for at least two months.² Gringras *et al*³ have shown that this system is feasible in general practice.

FIONA STRANG
R W BEARD
D J JEFFRIES

Department of Obstetrics and
Gynaecology,
St Mary's Hospital Medical School,
London W2

¹ Giles, P F H, Mackay-Scollay, E M, Christiansen, K, *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 1973, **13**, 65.
² Department of Health and Social Security, *Rubella Vaccination*, CMO 76/4. London, DHSS, 1976.
³ Gringras, M, *et al*, *British Medical Journal*, 1977, **2**, 245.

Herpes simplex virus excretion in febrile immunosuppressed patients

SIR,—Herpes simplex virus (HSV) as well as other herpes viruses, such as cytomegalovirus (CMV) and varicella-zoster virus, are known to cause serious infections in the immunosuppressed host. Moreover, asymptomatic excretion of CMV and of another herpes virus, the Epstein-Barr virus, is more frequent in these patients than in patients without an immune deficiency. We have found that this may also be true of HSV.

We cultured urine and saliva on human diploid fibroblasts searching for CMV in 60 patients with unexplained fever. Of these 18 were considered to be immunosuppressed (14 kidney transplant recipients, 2 patients with treated systemic lupus erythematosus, and 1 each with lymphoma and chronic lymphatic leukaemia). Their mean age was 37.5 years. The remaining 42 patients, with a mean age of 35.8 years, did not have a history suggesting an immune deficiency. CMV was isolated in 6 immunosuppressed and in 10 non-immunosuppressed patients. To our surprise HSV was isolated in 7 (39%) of the former and in 2 (5%) of the latter ($\chi^2=45$, $P<0.001$) (see table). Seven of 15 (47%) saliva specimens of the immunosuppressed patients were positive. All these isolates were identified as HSV-1 by microneutralisation using specific antisera, except for one HSV-2. None of these patients had oral lesions.

Asymptomatic HSV excretion has been described in the same proportion (5%) of non-immunosuppressed patients with respiratory infections as of our patients with

unexplained fever.¹ Isolation of HSV in the immunosuppressed is probably much more frequent but has in general been associated with herpetic lesions.^{2,3} However, the high recovery rate of HSV in our febrile immunosuppressed patients without orolabial lesions was unexpected.

H STALDER

Department of Medicine,
Hôpital Cantonal,
Geneva

¹ Lindgren, K M, Douglas, R G, Jun, and Couch, R B, *New England Journal of Medicine*, 1968, **278**, 517.

² Ruffault, A, and Fauconnier, B, *Pathologie et biologie* (Paris), 1976, **24**, 37.

³ Aston, D L, Cohen, A, and Spindler, M A, *British Medical Journal*, 1972, **4**, 462.

Parenteral nutrition

SIR,—After many years of development and evaluation of total parenteral nutrition most clinicians in this country now accept that intravenous 5% dextrose is a completely inadequate source of energy. Even our medical students appreciate that 2 litres of 5% dextrose per day only provide about 1.7 MJ (400 kcal), whereas the daily energy requirements are at least 6.3–8.4 MJ (1500–2000 kcal) with considerably increased requirements in seriously ill catabolic patients.¹ It was therefore with more than a little concern that we read, in the middle of the otherwise excellent article by Mr M E J Beard and his colleagues on acute marrow folate deficiency during intensive care (11 March, p 624), that for 10 days the patient's "energy intake was maintained [our italics] with 5% dextrose; he took no food by mouth." That this kind of thinking still exists in authors (as well as in referees) demonstrates the requirement for further education in nutritional requirements of seriously ill patients. Paradoxically, this report was printed next to an article on central venous catheterisation to permit infusion of hypertonic nutrient solutions.

ALAN SHENKIN
FRANCES J DRYBURGH

Department of Biochemistry,
Royal Infirmary,
Glasgow

¹ Lee, H A, *Parenteral Nutrition in Acute Metabolic Illness*. New York, Academic Press, 1975.

Skateboard injuries

SIR,—There is considerable interest in the "dangerous sport" of skateboarding and, while not wishing to minimise the problems of epiphyseal injury, etc, there appears to be an overemphasis on the dangers.

This accident and emergency department has a population at risk of 350 000 and in the nine-month period 1 June 1977–28 February 1978 there were 43 777 new patients, of whom 116 had sustained skateboard injuries. Fractures constituted 33.6% of the skateboard injuries; 23 of these 39 fractures involved the

Isolation of herpes simplex virus in febrile patients with and without immune deficiency

	No (%) of patients					
	Immunosuppressed			Non-immunosuppressed		
	Total	No with saliva culture	No with urine culture	Total	No with saliva culture	No with urine culture
No who had cultures ..	18	15	16	42	39	41
CMV positive ..	6 (33)	4 (27)	5 (31)	10 (24)	7 (18)	8 (20)
HSV negative ..	7 (39)	7 (47)	0 (0)	2 (5)	2 (5)	0 (0)