Training role of the non-teaching hospital

SIR,—The letter from Professor J Parkhouse (22 January, p 235) seems likely to provoke a good deal of resentment. He appears to misunderstand the role of non-teaching hospitals, particularly so far as senior registrars are concerned. I am at present a senior registrar in a non-teaching hospital and have previously held posts both in undergraduate teaching hospitals and in non-teaching hospitals as senior house officer, registrar, and senior registrar, so perhaps I may be permitted to comment.

Professor Parkhouse appears to consider that until a trainee specialist has reached senior registrar grade he is a liability to patients and staff. Conversely he implies that a senior registrar has much to offer and presumably he can then go as a missionary to the rude mechanicals of the service hospital. I had thought that one achievement of the NHS was to eradicate this type of condescension. It is to be hoped that as a senior registrar a trainee will have special knowledge of some subjects and thus can make a particular contribution, but in a hospital such as this there is little likelihood of a one-way traffic. A high academic standard is quite common in non-teaching hospital consultants, who often accept a very substantial teaching commitment at all levels. Even research is not unknown here in the backwoods—for example, the development of total hip replacement has been almost entirely the work of peripheral hospitals.

It is true that there is a danger that a heavy service load may swamp a trainee who is under-supervised, just as in some "centres of excellence" the medical staff may so outnumber the patients that no single doctor gets much experience. A great achievement of the past 20 years has been the increasing interchange of trainees at all levels between different types of hospital and a recognition of the importance of teaching in every hospital. It is by improving further the teaching and facilities provided in hospitals such as this one that standards will rise, and not by retreating into ivory towers where academics can keep themselves clean from "service," formerly known as patients.

J V PARKIN

Norfolk and Norwich Hospital,
Norwich

Points from Letters

Recruitment in geriatrics

Dr W FINE (Liverpool) writes: . . . No, "Charles Hastings," (15 January, p 182) the problems of recruitment in geriatrics stems not from lack of intellectual challenge but from the subconscious dislike by many of the young for the old, the attitude of teaching hospital physicians to the specialty, and above all the existence of too many job opportunities in other sections of our medically overstaffed hospital service. Junior staff has expanded in some areas manifold since 1948, especially in general medicine and surgery and kindred specialties. Consultant staff has expanded so much in some cities there is one consultant physician per 30 000 population. Recruitment to geriatrics will be improved greatly by one simple method. A 10% reduction of establishments of all medical posts, junior and senior,

across the board. Once high-calibre junior doctors enter geriatric units they are usually happy to continue working there. Training for general practice in many parts of the country has resulted in the inclusion of geriatric posts within the rotational schemes. Such training should be obligatory in the training of any general practitioner. It is surprising that in some areas of this country geriatrics is not included in such rotational training schemes. If one looks at the time that general practitioners spend among different groups of patients, more hours are spent with elderly patients than with surgical problems, yet in the training of the medical student a surprisingly small amount of time is spent on geriatric units. . . .

Kidneys from cadavers

Dr W F JACKSON (Deputy Editor, *Hospital Update*, 33/34 Alfred Place, London WC1E 7DP) writes: I read your recent leading article (22 January, p 188) with interest. As you say, public attitudes to transplantation appear to be favourable and I am convinced that the major problem in the supply of donors lies within the medical profession. . . . Although many are convinced of the value of renal transplantation, most doctors outside the field lack the necessary information on the suitability of their patients as potential donors. This has resulted in an under-reporting of suitable donors to transplant centres. In the hope of widening knowledge of the criteria for acceptance of kidneys we recently published a chart, drawn up with the aid of Professor P J Morris and the National Organ Matching and Distribution Service, which summarised the features for "ideal" and "acceptable" kidney donors.¹ Also included in the chart was a list of telephone numbers of regional transplant units. Our hope was that this chart would serve to remind "non-renal" hospital doctors of the possibility that their patients might be potential kidney donors. I understand that the chart has proved helpful in many centres, and I would be happy to provide copies of the chart to any doctors who missed it when it was originally published.

¹ *Hospital Update*, 1976, 2, 405.

Self-poisoning with drugs

Dr M A HARPER (Department of Psychological Medicine, Welsh National School of Medicine, Cardiff) writes: Many clinicians will share Dr D I R Jones's concern and perplexity about the mounting incidence of admissions due to self-poisoning (1 January, p 28), behaviour which in 1972 was estimated to cost the Health Service a minimum of £2m annually.¹ . . . I suggest that patients who secure admission to hospital following a deliberate drug overdose should be charged for the facilities they have monopolised, perhaps on a trial basis. To be effective as a deterrent the likelihood of receiving an appropriate bill should be widely known and publicised. The receipt of a bill for damages promptly terminated the destructive behaviour of a ward window smasher under my care in the past, who not only paid up promptly but even appeared to appreciate being regarded as a member of society responsible for her actions. The imposition of charges might be a more effective and appropriate negative reinforcer

than the reintroduction of attempted suicide as a criminal offence or the hostile reaction patients sometimes face from the staff involved in their resuscitation. No one would wish to charge those whose overdose was determined by a clearcut morbid process and there is often considerable uncertainty about the diagnosis in an individual patient, but a recommendation for further psychiatric care as an inpatient or outpatient might well constitute a practical test of a theoretically thorny point.

¹ Smith, A J, *British Medical Journal*, 1972, 4, 157.

Words

Dr D L GULLICK (Stevenage, Herts) writes: . . . Dr H W Ashworth (22 January, p 226) ascribes to the Mad Hatter that truly glorious observation, "When I use a word, it means just what I choose it to mean—neither more nor less." It was of course Humpty Dumpty who vanquished Alice with this "nice knock-down argument." Dr Ashworth is right, of course; Humpty Dumpty has a lot of disciples nowadays.

Iodine content of bread

Dr N FISHER (Flour Milling and Baking Research Association, Chorleywood, Herts) writes: Some concern has been aroused as a result of the article by Drs J C Stewart and G I Vidor on thyrotoxicosis induced by iodine contamination of food (14 February 1976, p 372) regarding the iodine content of British bread. Iodates, the source of the iodine implicated in the article, are not legally permitted additives in the UK and are not used. Iodized salt is not used in commercial breadmaking in this country either.

An IUCD record?

Mr A M SMITH (Wolverhampton) writes: In these days, when one reads and learns of the various complications of the intrauterine contraceptive device, I was a little surprised to find that one of my patients had had an IUCD with no trouble whatsoever for 10 years and 4 months. Is this a record?

Is golf enough?

Dr S B BALLANTINE (Dunbar, E Lothian) writes: I heartily endorse Mr A W Fowler's letter (15 January, p 172). To my mind golf is useless as an exercise to prevent coronaries. After all, most of the game consists in gentle putting, a little walking, a few swings, and a lot of standing about. . . . The Americans with their marathon running (Dr T J Bassler, 22 January, p 229) have got the right idea. Golf if you wish, it's a fine game of skill, but, as Mr Fowler says, let's do some jogging first.

Dr I S M SMART (Bonnyrigg, Midlothian) writes: I was incensed to read Mr A W Fowler's letter (15 January, p 172) asking, "Is golf enough?" As you never give precise causes of death in your obituary column how does he know that lack of exercise was a major factor in these six deaths with an average age of 61 years? Has he no sensitivity to the feelings of these doctors' relatives? . . .