recently. The original ulcer had healed. He was transfused with a total of six units of blood and made an uneventful recovery. The patient's claim, the witness of his relatives, and a return tablet count support compliance with therapy. His claim to abstinence from alcohol is supported by his relatives and the fact that his liver function tests had returned to normal. We were satisfied that no further gastric irritants had been ingested.

This case history emphasises the point that duodenal erosions leading to severe bleeding can develop even during treatment with cimetidine. Some authorities have recommended cimetidine in the treatment of bleeding from gastric erosions arising alone or in association with liver disease.¹ It has not, however, been shown to heal erosions and their pathogenesis is still not understood.^{2,3,4} The dangers of gastrointestinal bleeding are considerable and complacency must be avoided. Cimetidine is not a panacea for gastrointestinal bleeding and its advent should not decrease our vigilance in dealing with such patients, especially when they have associated liver disease.

B D LINAKER
S HUGHES

Gastroenterology Unit,
Broadgreen Hospital,
Liverpool

¹ MacDougall, B R D, *et al*, in *Cimetidine*, ed W L Burland and M A Simkins, p 329. International Congress Series No 46. Amsterdam, Excerpta Medica, 1977.

² Silen, W, *et al*, *Advances in Internal Medicine*, **19**, 195.

³ Mann, N G, *American Journal of Digestive Diseases*, **1976**, **21**, 89.

⁴ Morley, F G, *et al*, *American Journal of Digestive Diseases*, **1976**, **21**, 146.

Antibacterial activity of cefuroxime

SIR,—With the recent introduction of the new cephalosporin cefuroxime, which is reported to exhibit marked stability to most of the bacterial β -lactamases,¹ we would like to recount the outcome of a recent survey of bacteriological sensitivities which we have carried out in our 450-bedded district general hospital.

During three months we isolated 2537 bacteria from clinical specimens and 2511 of these were tested against cefuroxime using 30- μ g discs. Isolation and identification were carried out by standard laboratory methods supplemented in some instances by the API 20E and 20A systems. Sensitivity tests were performed by the usual method.

Cefuroxime was found to be active against the vast majority of Gram-positive and Gram-negative aerobic and anaerobic bacteria from whatever source they had been isolated—for example, sputum, pus, or urine. In many instances over 90% of each genus were recorded as sensitive—for example, all strains of *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Str mitis* (viridans groups), β -haemolytic streptococci (groups A, B, C, G), *Haemophilus influenzae*, *Neisseria gonorrhoeae*, and 97.2% of *Escherichia coli*, 96.7% of *Staph epidermidis*, 94.3% of *Proteus mirabilis*, and 93.8% of *Klebsiella* spp. Grouping all the enterobacteriaceae together, 93.5% were sensitive to cefuroxime, but none of the pseudomonade were sensitive. Most anaerobes were sensitive (91.1%), although if just *Bacillus fragilis* is considered only 80.9% were sensitive. Comparison of our data with those from a larger survey in the USA² shows close similarities.

Overall, 91.7% of the isolates in our study were sensitive to cefuroxime and this wide

antibacterial cover should make it a useful antibiotic for the treatment of infections requiring prompt parenteral therapy before the causative pathogen has been identified.

It appears that cefuroxime can be added to the treatment of various infections—for example, vaginal, ear, nose, and throat, eye, chest, etc.

L POTASCHMACHER
K A JEFFERSON

Microbiology Department,
Southend General Hospital,
Essex

¹ O'Callaghan, C H, *et al*, *Antimicrobial Agents and Chemotherapy*, **1976**, **9**, 511.

² Jones, R N, *et al*, *Antimicrobial Agents and Chemotherapy*, **1977**, **12**, 47.

Contracting in or contracting out?

SIR,—I have held a kidney donor card almost since they first became available and indeed my card is almost indecipherable. Surely a sensible solution to the problems and doubts raised in your leading article (15 April, p 944) is tattooing.

In any fatal incident, notoriously those occasioned by motor accidents, singed or blood-soaked clothing becomes a barrier to the card and its information. A rapid inspection for a tattoo placed, for example, under the foot of the victim should become standard practice. A central computer could record the result of the tissue and the blood grouping determined at the time of the tattoo. I leave it to your more expert readers to judge whether more donors would be lost at the thought of a painful (?) tattoo than are lost through the current system.

D C BISHOP

Turnford,
Broxbourne,
Herts

The postcoital test: what is normal?

SIR,—Mr G T Kovacs and others (1 April, p 818) are to be congratulated on their research into what is normal in postcoital tests. However, they have not stated what the period of continence was for each couple before they performed on the night before the test. Men who have intercourse every night may produce few or no sperm, or even blood, when asked to produce a specimen. Therefore a period of continence should precede the test, and it should be of the same length as for a full sperm analysis. This period of continence may lead Dr J Slome (15 April, p 992) to change his mind about the finding of sperm in the test.

HAYDN DIDIER

Bedford

Who discovered the circulation of the blood?

SIR,—I strongly support the claim made by Dr S Al-Dabbagh (15 April, p 992) that Ibn al-Nafis, a 13th-century court physician to Sultan Baybars of Egypt, discovered the pulmonary circulation three centuries before the Portuguese Servetus¹ and the English Harvey. Not unlike Harvey, Ibn al-Nafis met with great opposition for criticising Galen and departing from traditional medicine—so much so that some historians declined to include him in their biographies. It would not therefore be amiss to rechristen the discovery

as "Nafiso-Harveian," for academic precision.

Luckily, I have access to five manuscripts by Ibn al-Nafis in the Bodleian Library, one of which is a holograph written by Ibn al-Nafis himself shortly before his death in 1288. I am quite willing to provide parallel texts from Ibn al-Nafis and Harvey to demonstrate the striking resemblance between the results arrived at by the two court physicians. Perhaps this will be the best present we can offer Harvey on his 400th birthday, as he was a great seeker of the truth.

S A KHULUSI

Clarendon Building,
Oxford

¹ Hitti, P K, *History of the Arabs*, p 685. London, Macmillan, 1964.

Hepatitis B in the school environment

SIR,—It is well established that viruses are readily transmitted in a school environment. A previous study¹ in two Glasgow schools showed that a significant number of children of African, Asian, or Chinese parents carry hepatitis B surface antigen (HBsAg) whereas native Scottish children carrying HBsAg are extremely rare. As carriers of HBsAg are potential sources of infection with hepatitis B virus we investigated whether the presence of carriers in these schools led to significant transmission of infection to their classmates.

Presence of antibody to HBsAg (anti-HBs) was used as an indicator of prior exposure to HBsAg. Sera from groups of children of African, Asian, Chinese, and Scottish parentage were examined for anti-HBs (Ausab, Abbott Laboratories). Anti-HBs was detected in some of the non-Scottish children but none of the native Scottish children (see table below).

Results of tests of schoolchildren's serum for HBsAg and anti-HBs

Group	No examined	No positive for antigen	No positive for antibody
African	88	3	4
Asian	142	2	3
Chinese	95	4	13
Scottish	78	0	0

It is unlikely that this higher prevalence of anti-HBs in African, Asian, and particularly Chinese children indicates that HBsAg is circulating in these schools preferentially among these children. What is more probable is that this higher prevalence merely reflects a much higher level of HBsAg carriage among their parentage than among the native Scottish population. In the previous study¹ sera were obtained from the parents of five carrier children—all parents' sera were positive for either HBsAg or anti-HBs.

The finding of no antibody among the native Scottish children was particularly interesting. To substantiate this finding we have obtained blood specimens from a further 146 native Scottish children through the co-operation of parents, school staff, and health visitors. Drops of blood were taken by finger puncture on to absorbent card and the extracts tested for anti-HBs. Tests on known positives showed this method to be less sensitive than testing whole serum but still capable of detecting all but very low titres of antibody. All 146 specimens were negative for anti-HBs.

Blood has long been known to transmit hepatitis B infection and recently saliva has been