destroy the identity of individual journals is more an indictment of the journal than a sensible excuse. No journal should have to rely on editorial style to make its mark. One hopes that there will be uniform and rapid support for the Declaration of Vancouver by all concerned in journal editing. We could then all concentrate more on the subject matter of papers and worry less about irritating minutiae.

K G M M ALBERTI
Editor-in-Chief,
Diabetologia

Southampton

SIR,—The citing of references is a valuable way of drawing attention to additional information or discussion without lengthening a text. But it is not unknown, even in journals of the highest repute, for authors to misinterpret or even falsify data taken from their cited references. When reference is made to work "in press" or awaiting acceptance the reader cannot check or follow up this at all.

Surely the International Steering Committee of Medical Editors (20 May, p 1334) should acknowledge that their readers take precedence over their authors' convenience and should refuse to accept work citing references which readers cannot easily consult. After all, if the information or discussion alluded to in an unpublished reference is at all relevant it can be incorporated in the text of the article, where the reader can assess its worth.

ROGER HOLE

Department of Urology,
North Ormsby Hospital,
Middlesbrough, Cleveland

Interrelated factors in hypertensive patients

SIR,—The summary and conclusions of the paper by Dr D Ballantyne and others (8 April, p 880) may suggest that findings in a hospital sample of hypertensive patients could challenge the validity of epidemiological relations established in properly drawn samples of the general population.¹ Apart from what must be an error in the standard deviation (SD) for systolic blood pressure in female non-smokers (and possibly smokers), the SDs are larger rather than smaller than those in samples of

people at work in the general population and this slightly questions the quality of the data. Bias due to selection of the sample is perhaps indicated by differences of +6% and -19% in the numbers of male and female smokers respectively compared with employed members of the general population.

Table I shows significantly lower levels of diastolic blood pressure (phase V, seated, mean of two readings) in relation to the aggregated values of the 80th and 95th percentiles of the age and sex distributions by five-year intervals for men and women examined at work in the west of Scotland, for male and female smokers. In our study cigarette smoking is also correlated strongly with alcohol consumption and high alcohol consumption in turn to high blood pressure, a relationship noted by other workers.^{2,3}

Table II shows the relation of blood pressure to obesity measured by Quetelet index ((weight × 100) ÷ (height × height)) at the 20th and 80th centiles of the QI distribution. Obesity is also related to alcohol, but the interaction of all three even after removal of the age effect is another question and one to which we have not yet the answer.

Characterisation of disease rather than patients can be determined more certainly in the general population.

V M HAWTHORNE
MARY SMALLS

Mass Health Examination Unit,
Glasgow

¹ Hawthorne, V M, *et al*, *British Medical Journal*, 1969, 4, 651.

² Ramsay, L E, *Lancet*, 1977, 2, 111.

³ Beevers, D G, *Lancet*, 1977, 2, 114.

SIR,—We were interested to read the report by Dr D Ballantyne and his colleagues (8 April, p 880) showing no relation between adiposity and blood pressure, except in male non-smokers, in a study of untreated hypertensive patients with diastolic blood pressures greater than 90 mm Hg.

In the Whitehall study of cardiorespiratory disease in 18 403 London male civil servants aged 40-64 years¹ blood pressure was measured on a single occasion and smoking habits recorded. As in the Glasgow study, there was no significant difference in mean blood pressure

(systolic or diastolic) between current cigarette smokers and non-smokers in those subjects with diastolic blood pressures greater than 90 mm Hg. However, contrary to the Glasgow findings, in this hypertensive subgroup there was a significant positive relationship between blood pressure and body mass index (weight/height)² regardless of smoking status.

It is possible that the negative Glasgow results might be due to differences between our studies in the length of time elapsed between the smoking of the last cigarette and the measurement of blood pressure. Alternatively their patients' smoking habits might have changed during the period of the three visits over which blood pressure was averaged.

JOHN H FULLER
M J SHIPLEY
GEOFFREY ROSE

Department of Medical Statistics
and Epidemiology,
London School of Hygiene and
Tropical Medicine,
London WC1

¹ Reid, D D, *et al*, *Lancet*, 1974, 1, 469.

The Stomach Cancer Group

SIR,—Stomach cancer is the third most common cancer in Britain. At present only one in six of those in whom the growth is radically resected will be alive in five years' time and, overall, the five-year survival is less than 10%. Appalled by the outlook for such patients, this group of interested clinicians has been meeting in an attempt to improve the treatment of stomach cancer.

For the past 18 months we have entered patients after gastrectomy to a multicentre controlled trial of cytotoxic chemotherapy. Drug therapy is given for two years after operation. The regimen of 5-fluorouracil and mitomycin-C given once every three weeks is well tolerated by the patients and the incidence of side effects is low. All patients are being closely monitored, in particular for biochemical and haematological changes.

So far eight centres are involved and 137 cases have been entered into the study. We would welcome to this group others keen to study this disease and its treatment. The next meeting of the group will be held in June 1978. All communications should be addressed to the secretary.

PETER F M WRIGLEY
Secretary

Department of Medical Oncology,
St Bartholomew's Hospital,
London EC1

VICTOR BROOKES
Chairman,
The Stomach Cancer Group

Queen Elizabeth Hospital,
Birmingham

Television medicine

SIR,—Sir Michael Swann's bedside manner deserves a tribute, and his imperturbability in a controversial situation should neutralise some circulating antibodies. Now that he has joined in the discussion (13 May, p 1274) I feel I can follow without any imputations.

While I was a member of its General Advisory Council (GAC) I came increasingly to respect the BBC, a body that is widely recognised as among the most praiseworthy of all British institutions, and my criticisms are those of a sincere and friendly admirer.

TABLE I—Cigarette smoking related to 80th and 95th percentiles of age-standardised diastolic blood pressure distributions in 5680 men and 987 women

Smoking category	Men				Women			
	n	80th	80th and 95th	95th	n	80th	80th and 95th	95th
Smoker	2794	2370 (84.8)	315 (11.3)	109 (3.9)	571	487 (85.3)	61 (10.7)	23 (4.0)
Non-smoker	1259	1039 (82.5)	160 (12.7)	60 (4.8)	343	267 (77.8)	48 (14.0)	28 (8.2)
Ex-smoker	1627	1259 (77.4)	293 (18.0)	75 (4.6)	73	58 (79.5)	12 (16.4)	3 (4.1)

Figures in parentheses represent percentages of totals

TABLE II—Blood pressure (mm Hg) related to obesity in 5686 men and 986 women

Obesity level by QI distribution	Men					Women				
	n	Systolic		Diastolic		n	Systolic		Diastolic	
		Mean	SD	Mean	SD		Mean	SD	Mean	SD
Low (<20%)	1057	128.5	16.2	79.1	10.1	268	130.2	16.7	78.2	9.6
Medium (≥20%)	3467	133.1	17.2	83.3	9.8	535	137.5	19.5	82.7	10.9
High (≥80%)	1162	139.1	18.7	88.6	11.2	183	146.7	20.8	89.1	12.2

Two years ago, in a study ("The BBC's medical programmes and their effects on lay audiences") for the GAC by the department of science and features, it was stated that "the responsibilities of this department, where medicine is concerned, are no different from those which apply to other scientific topics dealt with in our programmes." This tidy façade I thought to be a product of superficial reasoning. It takes no account of the connotations and imponderables in medicine or of the unique influence and immediacy of matters medical on the attitudes and lives of the lay public.

The approach to medicine could not have been thought out. How else to explain "It is programmes that make policy, not policy that makes programmes"?—a remarkable utterance by the head of science and features department at the Royal Society of Medicine in 1976. This slick thought-dampener, far from being concealed, was flaunted later before the GAC and was accepted. Presumably it is still dogma in medical broadcasting.

My respect and admiration for the BBC are undimmed by the arguments you have published, which tend to focus on irritating superficialities. It is deeper, structural flaws that should be examined and corrected, preferably by the medical profession and BBC together.

JOHN APLEY

Bath

Treatment of tuberculosis

SIR,—I in turn am surprised at the different emphases in treating tuberculosis in the Liverpool School of Tropical Medicine (Dr D Robinson, 22 April, p 1053).

(1) I believe that where possible treatment should be based on the results of controlled trials and that inspiration, eccentricity, or years of experience are insufficient as alternative bases.

(2) I did not assume that the effects of *Mycobacterium tuberculosis* and the syndrome of mycobacterial diseases are "the same thing." Indeed, the brief reference to opportunist mycobacteria simply drew attention to the fact that they behave differently. To discuss their treatment more fully would have required more space than the Editor would allow.

(3) I have never before heard tuberculosis referred to as a "subdisease." It must rank as the most important and fatal non-disease in history. Anyone who feels that the laboratory plays the major role in the management of this subdisease must have little practical experience of dealing with patients suffering from it. Perhaps that is why Dr Robinson is not aware that most doctors and indeed many laymen actually regard it as a fully fledged disease. Sensible clinicians do, however, work in close contact with sensible bacteriologists in managing the condition.

(4) The confusion does not appear to be in the paragraph but in the mind of Dr Robinson. I feel, and others more authoritative and experienced than myself seem to agree, that contacts should be investigated. Not to do so would allow cases to go undetected in the community.

(5) The side effects of the drugs are listed in my article and further information is available and referred to. Isoniazid-induced hepatitis is, as I stated, uncommon in Britain. My article makes clear that follow-up of patients is essential and that drug side effects are a problem to be aware of. Did Dr Robinson read it?

(6) Rifampicin has already shortened the standard regimen and the evidence is quoted. There are few contraindications to its use and it has now been used widely in pregnancy. Dr Robinson is badly out of date.

(7) Practice in Britain has now relegated strepto-

mycin to a somewhat less important role, though, as I state, it remains a first-line drug. Ethionamide and pyrazinamide are generally used as second-line drugs and both are mentioned in my article.

(8) Sputum culture is mentioned as a keystone in the diagnosis and is an important aspect of follow-up. The primary role in the management of tuberculosis, and with it the responsibility, should be that of the physician. A nurse or health worker is an invaluable aid in this but is surely not able to take primary responsibility.

I reply to all Dr Robinson's points because I believe he has over-reacted to my article and is almost entirely wrong in what he says. I do not wish your readers to be misled by his letter and would refer them back to my article and the suggested further reading to judge for themselves.

ANTHONY SEATON

Institute of Occupational Medicine,
Edinburgh

Diuretics in the elderly

SIR,—Your timely reminder of the dangers of prescribing diuretics to elderly patients (29 April, p 1092) fails to mention the problems of magnesium deficiency. That increased excretion of magnesium occurs has been known for some time¹ and symptomatic hypomagnesaemia has also been described.² The symptoms encountered in magnesium depletion include confusion, ataxia, tremor, nausea, and palpitations, all of which commonly occur in the elderly. Recently Ibrahim and Sutcliffe³ showed a high incidence of symptomatic magnesium deficiency in the elderly which responded well to replacement therapy.

As a number of elderly patients receiving diuretics may also be having digoxin it is important to remember that hypomagnesaemia is associated with an increased sensitivity to digoxin⁴ as well as with primary cardiac arrhythmias.⁵

I would therefore suggest that magnesium depletion should always be considered in elderly patients who become unwell, especially if on diuretics.

S WALDEK

University Department of Medicine,
Royal Hospital,
Sheffield

¹ Smith, W O, *et al*, *Clinical Research*, 1959, 7, 162.

² Wacker, W E C, and Parisi, A F, *New England Journal of Medicine*, 1968, 278, 772.

³ Ibrahim, I K, and Sutcliffe, R L G, *Modern Geriatrics*, October 1977, p 6.

⁴ Sellar, R H, *American Heart Journal*, 1971, 82, 551.

⁵ Chadda, K D, Edgar, L, and Gupta, P, *American Journal of Cardiology*, 1973, 31, 98.

SIR,—Once again Slow-K is quoted as a cause of ulceration of the small intestine (leading article, 29 April, p 1092). This is surely an extremely rare event and is almost always associated with pre-existing gut stricture or slow intestinal transit time, as in the reference cited by you.¹ Bearing in mind that approximately 600 million tablets of Slow-K are dispensed in the United Kingdom every year, Slow-K must be just about the best tolerated preparation on the market. It must not be confused with enteric-coated tablets.

DENIS BURLEY

Head of Medical Services,
Ciba Laboratories

Horsham, W Sussex

¹ Farquharson-Roberts, M A, Giddings, A E B, and Nunn, A J, *British Medical Journal*, 1975, 3, 206.

Sulphinpyrazone after myocardial infarction

SIR,—In response to your leading article (15 April, p 941) concerning rhythm abnormalities in the sulphinpyrazone cardiac infarction trial I should like to quote from our original paper¹:

"Although, as shown in table 2, more patients receiving placebo (111) than patients receiving sulphinpyrazone (83) had cardiac rhythm abnormalities at the first visit, deaths among patients with rhythm abnormalities (six in the placebo, and two in the sulphinpyrazone group) represented only 12% of all deaths; furthermore, the presence of abnormalities of rhythm at entry into the study did not appear to be a significant risk factor since 5.4% of placebo-treated patients with such abnormalities died as compared to 6.0% of placebo patients without them."

In addition, we have also calculated the death rates per 28-day exposure period $\times 1000$. The findings are shown in the table:

Treatment group	Rhythm abnormalities	
	No	Yes
Placebo	7.4 (38)*	6.6 (6)
Sulphinpyrazone	4.0 (23)	2.9 (2)
Combined	5.7 (61)	5.0 (8)

*Values in parentheses are actual numbers of deaths.

The statistical analyses employed in this trial (Cox model) permit adjustment in balances of entry criteria. The possible imbalance of rhythm abnormalities was thus adjusted for and did not account for any significant difference between unadjusted and adjusted P values.

The final report of this trial will include details of all aspects of electrocardiograms in so far as our data base will permit.

SYDNEY H KANE
for Policy Committee,
Anturane Reinfarction Trial

Pharmaceuticals Division,
Ciba-Geigy Corporation,
Summit, New Jersey

¹ Anturane Reinfarction Trial Research Group, *New England Journal of Medicine*, 1978, 298, 289.

Plasma urate changes in pre-eclampsia

SIR,—Mr W Dunlop and Mr J M Davison (25 March, p 786) query our use of plasma urate increments (25 February, p 467) to define grades of severity of pre-eclampsia. While we accept their general point that small increases of plasma urate cannot be used clinically to diagnose patients with pre-eclampsia, their claim that these changes may be "physiological" is not necessarily valid. That plasma urate levels above 350 $\mu\text{mol/l}$ (6 mg/100 ml) indicate significant pre-eclampsia has always implied that antenatal increases of plasma urate have already occurred. This is also clearly demonstrated in our recent paper. A priori in the development of pre-eclampsia there must be a time when plasma urate levels show small rises above an earlier baseline reading. The morbidity of this prodromal stage would not be expected to be high because in many women the plasma urate changes either coincide with or precede the elevation of the blood pressure. When elevation of the plasma urate occurs at or near term pregnancy-induced hypertension may not arise as the process will normally be cut short by spontaneous or induced delivery.