an hour, and was accompanied by agitation and sweating but not by palpitation or dyspnoea. Similar symptoms followed most subsequent doses of nifedipine; they ceased as soon as the drug was discontinued two weeks later. An electrocardiogram recorded since was normal.

Our patient also complained of a fine tremor involving particularly the upper limbs. It occurred within 30 min of a dose of nifedipine and lasted up to two hours. It developed after he had been on nifedipine 30 mg daily for a month, was unchanged on 60 mg daily, and resolved immediately the drug was stopped. This possible side effect of nifedipine has not apparently been reported previously.

CHRISTINE RODGER
A STEWART

Monklands District General Hospital,
Airdrie, Lanarkshire

SI, moles, and drugs

SIR,—We wholeheartedly support the plea made by Dr P Crome and others (13 May, p 1277) for the preservation of mass units for the expression of drug concentrations. As they point out, the use of several different units is a particular hazard in clinical toxicology and we could provide many examples such as those quoted in their letter. We too have experienced major problems with inquiries to the Scottish Poisons Information Bureau and our toxicology laboratory when the units used by different hospitals differ. The caller is usually unaware of the units used locally.

The introduction of molar SI units for a few drugs has exacerbated a pre-existing problem with units. Barbiturate, salicylate, and paracetamol concentrations have previously been reported in different mass units, generally $\mu\text{g/ml}$ or $\text{mg}/100\text{ ml}$, and now and again mistakes in interpretation have occurred because of this. The addition of molar units in the east of Scotland has now resulted in the use of both $\mu\text{mol/l}$ and mmol/l for paracetamol. Other laboratories and the National Quality Control Scheme for this drug continue to use mass units. With such commonly encountered drugs confusion over units remains a problem despite assurances that increasing familiarity with the new units would obviate the need for conversion.

In the case of less commonly measured drugs there are no reference values for molar units and no comprehensive list of conversion factors is available. In order to interpret a drug concentration quoted in a poisoning inquiry it is necessary to find a reference book containing the chemical formula or the molecular weight before the conversion can be calculated. This may not be too difficult for laboratory staff, but it is asking a lot of the secretaries, doctors, and nurses who staff our poisons information bureau. The dangers are well illustrated by the serious mistakes recently made by both your editorial office (18 March, p 716) and Dr S G Ball (8 April, p 920) in attempting to convert chlormethiazole concentrations.

It must also be noted that drug concentrations in most of the clinical toxicology and clinical pharmacology literature nationally and internationally are expressed in mass rather than molar units. A change is unlikely since drugs are manufactured, prescribed, and dispensed in mass units and the use of different units for dosage and concentrations in clinical pharmacokinetics is nonsensical.

If there were any practical advantages in using molar units the difficulties and hazards involved might be acceptable. But there are no obvious clinical advantages. In defence of SI it is sometimes claimed that the response to drugs is related more closely to molar than to mass concentrations at receptors. While this may be true in some circumstances it is largely irrelevant in clinical practice. For example, there may be up to a 500-fold range in mass (or molar) dosage of thiazide-type diuretics required to produce a similar clinical effect, which cannot be related to differences in plasma concentrations.

The present unacceptable situation has been brought about by those who devote only a small part of their total work load to drug analysis without consideration of the wider implications. The practical problems of clinical toxicologists, clinical pharmacologists, pharmacists, and those working in the pharmaceutical industry, all of whom are directly concerned with drugs, have not been taken into account. Unless some responsible body undertakes to secure general agreement on the use of mass or molar units for clinical drug concentrations the present muddle will continue indefinitely.

L F PRESCOTT
M J STEWART
A T PROUDFOOT

Regional Poisoning Treatment Centre
and Toxicology Unit, University Department
of Clinical Chemistry,
Royal Infirmary,
Edinburgh

Intravenous infusion of salbutamol in severe acute asthma

SIR,—The apologetic tone of the discussion at the end of the paper by Dr A J Johnson and his colleagues (22 April, p 1013) clearly betrays a lack of confidence both in the design of their study and in their rather lame conclusion that salbutamol given by intravenous infusion for 36 h does not "appear to have any advantages" over aminophylline administered in the same way.

It is, of course, true that there are ethical problems in conducting clinical trials on patients with severe acute asthma. One can, however, hardly imagine that patients who were already receiving a "standard supportive regimen comprising nebulised salbutamol (5 mg) given four times a day by intermittent positive-pressure breathing, corticosteroids, supplementary oxygen, and physiotherapy" combined with antibiotics when necessary (a generous therapeutic regimen by any standards) could have come to much harm, even if the initial response to this treatment was "unsatisfactory," as a result of the physician responsible for their care not knowing whether the supplementary (trial) treatment consisted of an intravenous infusion of salbutamol or one of aminophylline. Most hospital ethical committees would accept that view, provided the patients were all being expertly monitored, and permit both a double-blind allocation to the two types of infusion and the inclusion of a placebo infusion group.

In the way the study reported by Dr Johnson and his colleagues was, or perhaps had to be, conducted its results are virtually of no value. This is, in effect, conceded by the authors themselves when they admit (1) that "any possible additional bronchodilator effect from the intravenous administration of salbutamol may have been reduced by the large

regular doses of nebulised salbutamol causing saturation of the adrenoceptor sites"; (2) that "physician bias against the new infusion drug" (salbutamol) may have influenced the results; and (3) that "possibly, if a placebo infusion group had been included it would have yielded results similar to those of the bronchodilator infusion groups."

Indeed, the only conclusion a critical reader can draw from the paper is that patients varied in the rapidity of their response to the authors' "standard supportive treatment." They have certainly not proved, nor indeed have they claimed, that the prolonged intravenous infusion of either salbutamol or aminophylline had any significant effect on their patients' rate of recovery. In this connection the allusion at the end of the summary to a "rate [of recovery] that would occur *naturally*" (my italics) in the context of patients who were receiving what would usually be regarded as very intensive treatment for severe acute asthma seems a little strange.

If the article proves anything at all it is perhaps that some hospital ethical committees are less enlightened than others.

IAN W B GRANT

Department of Medicine,
National University of Malaysia,
Kuala Lumpur

SIR,—The paper by Dr A J Johnson and others (22 April, p 1013) concludes that in acute severe asthma not responding to an initial intensive regimen the intravenous infusion of salbutamol "has no particular advantage over aminophylline." In reaching this conclusion it seems that the authors place little weight on the fact that 6 out of 20 salbutamol patients were withdrawn because their response was "considered to be unsatisfactory" compared with 0 out of 19 in the aminophylline group ($P=0.02$, Fisher's exact test). Furthermore, two of the aminophylline group were removed late in the trial because their response was *satisfactory*.

The finding of an absence of significant differences between the lung function of the two groups at any time during the trial appears to be based on the data of those patients who were not withdrawn at any stage. If so, this has almost certainly biased the result against aminophylline since four of the salbutamol withdrawals had lung function values well below the mean response of their group. It appears that, although it may be correct to state that salbutamol has no particular advantage over aminophylline, it would be more correct to conclude that aminophylline may have some advantages over salbutamol.

H R ANDERSON

Department of Clinical Epidemiology
and Social Medicine,
St George's Hospital Medical School,
London SW17

* * * We sent copies of these letters to Dr Johnson and his colleagues. A reply from two of them, sent with the concurrence of the others, is printed below.—Ed, *BMJ*.

SIR,—In response to Professor Grant's comments we must reiterate that the purpose of our study was to assess the effect of infusing the two bronchodilator drugs in asthmatics whose attacks were severe and had responded poorly to a standard initial regimen. This is a quite different situation from the previous