elapse between vaccination at school and first pregnancy and we cannot be sure that vaccination alone, in the absence of immunological stimuli through contact with the infection, will maintain enduring immunity.

The vaccination of schoolgirls appears to be slowly taking effect: a large sample of haemagglutination inhibition tests on adult women gave a susceptibility rate of 17% in 1974; in 1978 it had fallen to 12.7%. In a survey of Essex school-leavers of average age 18 years in 1976 4.3% of the girls were susceptible. Taking only those who claimed to have been vaccinated, there were only 1.3% susceptible. Thus in 10-15 years we may expect a susceptibility rate of about 5%; if the acceptance rate is increased from 68% to nearer 100% it should fall to 1%; if retesting is carried out after vaccination then the incidence should almost vanish. It may soon be preferable to screen schoolgirls before vaccination in order to concentrate on the susceptibles.

The situation of women who were not vaccinated at school is more serious. Very little propaganda appears to emanate from central sources and no amount of local pressure appears to improve the appallingly low incidence of screening of non-pregnant women (below 10% and mainly nurses). The first visit to the antenatal clinic is the traditional occasion for screening, but the timing could hardly be worse. Preferable occasions are during visits to a family planning clinic or when submitting specimens for pregnancy tests. Certain groups of women must be screened before taking up employment—nurses, teachers, social workers, etc.—and failure of the employer to make the necessary arrangements might invoke action under the Health and Safety at Work Act.

However, the most important aspect is the education of women in this matter. They have come to understand the danger of rubella infection during pregnancy; the medical profession should co-operate with the media to persuade them to demand rubella screening as a routine procedure before pregnancy is contemplated. This would avoid situations that today cause great distress to a large number of women.

R PILSWORTH
B J PILSWORTH

Public Health Laboratory,
Chelmsford, Essex

Epidemic myalgic encephalomyelitis

SIR,—I disagree with the view you express in your leading article (3 June, p 1436) that the many epidemics you cite compose a single nosological entity. Outbreaks in hospital staff apart, these miscellaneous episodes and others described at the symposium at the Royal Society of Medicine may well have been as diverse aetiologically as they were geographically.

Nor do I agree that the Royal Free Hospital outbreak has now been proved to have been a form of encephalitis. Much play was made at the symposium on the occurrence of extensor plantar responses and of other evidence of organic disease. In fact, however, only two of the 292 patients had a frank extensor response¹ and one of these was later found to have multiple sclerosis. The other evidence adduced was equally unconvincing.

As I have previously explained,² it is relevant to consider the management of infectious mononucleosis in the Royal Free

Hospital at the time and the results obtained³—which bore no comparison with those described in a standard textbook.⁴ There seems to have been failure to realise how suggestible people are when they are ill or anxious.^{2, 5}

In a letter to *The Times*⁶ entitled "Vapour or virus" Sir Francis Walshe stated: "Epidemics of mass hysteria have punctuated history and doubtless a few centuries ago were attributed to witchcraft or demoniacal possession on evidence not less than that now evoking the mystic chant of the word 'virus.'" Though the Royal Free outbreak was a sorry affair compared with dancing madness, the classic form of mass hysteria,⁷ ataxia, vertigo, fits, and prostration were common to both. Nowadays the vapour or virus, or whatever the devil it is, seems to be most at home in nurses' homes (in "centres of excellence" particularly). Any one who wants to know if Walshe's statement could be pertinent to the Royal Free epidemic should not be content with your leading article: he should study the evidence for himself.

H G EASTON

Clinical Department of Infectious
Diseases,
Ruchill Hospital,
Glasgow

¹ *British Medical Journal*, 1957, 2, 895.

² Easton, H G, *World Medicine*, 1977, 13, 13.

³ Dunnet, W N, *British Medical Journal*, 1963, 1, 1187.

⁴ Christie, A B, *Infectious Diseases: Epidemiology and Clinical Practice*, 2nd edn. Edinburgh, Churchill Livingstone, 1974.

⁵ Easton, H G, *British Medical Journal*, 1963, 1, 1473.

⁶ Walshe, F M R, *The Times*, 19 October 1965.

⁷ Dodds, E R, *The Greeks and the Irrational*. London, Cambridge University Press, 1963.

Improvement of perinatal care

SIR,—In your leading article (27 May, p 1372) on perinatal mortality you are guilty of an inaccuracy and a non sequitur when you refer to *The Place of Birth*¹ as advocating a move "back" to domiciliary midwifery and imply that this would necessarily lead to an increase in perinatal mortality or morbidity, or both.

It would seem that you have not read the book, for if you had it would have become clear that the introduction of 100% hospital delivery in England still leaves us a long way behind Holland, where the rate is in the region of 50%. What we advocate is a move "forward" to domiciliary midwifery (the distinction is nice but not precious); and we fail to appreciate why the concentration of abnormal midwifery in larger units properly staffed and equipped to cope with predictable emergencies should be incompatible with the conduct of predictably normal deliveries at home. Admittedly, a considerable proportion of emergencies are at present not predicted; but this may be partly the result of inadequate antenatal care and partly due to the present orientation of research in obstetrics. It is also important to point out that what matters is the final outcome and that there is a difference between the conditions in which obstetricians on the one hand and women on the other feel safe and those in which babies are safe.

What *The Place of Birth* argues is that both obstetric and paediatric establishments are becoming obsessed with our place in the perinatal league table at the expense of other considerations that we think are equally important and that the means that are proposed to achieve this end will not necessarily serve the purpose. For instance, it is by no means certain that special care units as opposed to

intensive care units are really necessary and they are certainly harmful. The joint committee² should have been more radical in their thinking and in their recommendations.

JOHN DAVIS

University Department of Child
Health,
St Mary's Hospital,
Manchester

¹ Kitzinger, S. and Davis, J., *The Place of Birth*. London, Oxford University Press, 1978.

² BPA RCOG Liaison Committee, *Recommendations for the Improvement of Infant Care during the Perinatal Period in the United Kingdom*, Discussion Document. London, British Paediatric Association and Royal College of Obstetricians and Gynaecologists, 1978.

How much physical therapy for patients with stroke?

SIR,—We welcome the attempt by Professor J C Brocklehurst and his colleagues (20 May, p 1307) to draw attention to the objectives of physical therapy for strokes. We fear that the conclusion they draw could result in the withdrawal of physiotherapy from patients who continue to require it and an increase in the permanent long-stay hospital population.

In their study patients were admitted according to bed availability to one or other of six different hospitals. Four hospitals provided no occupational therapy and three provided no speech therapy. When combined with the 24% incidence of pressure sores in the stroke patients this suggests that conditions for rehabilitation were below optimal. Physiotherapy in the article is measured in days, but in most hospitals the amount a patient receives is measured in minutes. It is important to know whether physiotherapy was carried out in isolation within a physiotherapy department or in the ward, supplemented by training of the nursing staff to reinforce the physiotherapy, together with integration with occupational therapy and speech therapy in the therapeutic team. It would be of interest too to know whether there was a between-hospital variation and it is necessary to know whether physiotherapy was of the traditional type or of the facilitative type such as Bobath or proprioceptive neuromuscular facilitation.

The results in this Manchester survey confirm that mobility continues to improve in 21% after 6 months and that 6% of the study group were restored to independence after 6 months. Withdrawal of trained therapists before this period will result in an increase in the study's current 16% of stroke patients remaining as inpatients at one year.

There also remains a real need to compare the performance of a fully staffed rehabilitation team working in the wards of a general hospital ward with the results of a stroke unit.

JOHN KNOX
PETER HORROCKS

Geriatric Service, Hull Area,
Kingston General Hospital,
Hull, North Humberside

* * * We sent a copy of this letter to Professor Brocklehurst and his colleagues, whose reply is printed below.—Ed, *BMJ*.

SIR,—Our study (20 May, p 1307) was a description of the management of patients with stroke as it happens at the present time and we agree with Drs Knox and Horrocks that the present situation leaves much to be desired. Our hope is that by indicating where