Nissen (20 May, p 1354) taxes us with not considering an affection of the median nerve, but this suggestion is contraverted completely by the clinical information available and therefore not worth discussion. After localisation, we proceeded in classical neurological fashion to consider possible causes of the lesion. Professor Walker scorns the idea that piano playing might be the cause of a hand injury producing chronic partial paralysis; he does not pause to consider our argument, based on personal anatomical and surgical observation, that if piano playing was related to the hand disability, then there must have been an associated anatomical variation. This caveat was an essential point in our discussion because of the absence of other reported cases among professional pianists. Incidentally, we know many rowing men and only one has developed a posterior interosseous palsy in relation to this pastime.

Lastly, Professor Walker takes us to task for our direct, acknowledged, quotation from Dr Sams, to whom we refer him, on the subject of Schumann and the finger strengthener he is said to have ordered. As a final blow, Professor Walker implies that we entertained the idea that the unproved use of such a machine was the cause of a condition present for seven previous years. Nothing we wrote supported this ridiculous interpretation.

It might be helpful if Professor Walker were to discuss the neurological problems of this tragic but interesting case with our colleagues in the department of neurology in the Medical School of McMaster University. Alternatively, we would be pleased to pursue the matter when he is next in London.

R A HENSON

Neurological Department,
London Hospital,
London E1

*.*This correspondence is now closed.—ED,
BMJ.

Kidney failure in liver disease

SIR,—With reference to your leading article (27 May, p 1375), since I was the first in this country¹ to emphasise the finding of endotoxin in the plasma of patients with hepatorenal failure and the larger series that have followed have simply confirmed this fact, may I be allowed to observe that it is time that further discriminatory work be done? The original observation followed the introduction of the limulus assay but stemmed from radio-fibrinogen studies which showed continuing fibrinogen catabolism in such patients.² This particular type of renal failure does not have the features of a Schwartzman reaction (glomerular and peritubular thrombosis) that is the standard response to endotoxin. In part this may be due to the very active fibrinolysis that is found in cirrhosis. That there is true disseminated intravascular clotting is shown by the improvement with heparin.³

Another anomaly is that these patients have high antibody titres to O antigens of Gram-negative organisms. Thus the endotoxin is likely to be present in plasma as endotoxin-antibody complexes. Yet there is no nephritis. Four factors that require exploration come to mind. Firstly, the antibodies may be of broad specificity and low affinity: such complexes are not easily precipitated by ammonium sulphate in the Farr test. Secondly, the low level of serum esterases in chronic

liver disease⁴ may imply an inability to inactivate endotoxin that has shunted past the normally active Kupffer cells of the cirrhotic liver. Thirdly, bile acid retention in cirrhotic plasma may mean that endotoxin is dissociated into subunits that are poorly antigenic and that react in a biologically different manner. Fourthly, for this previous reason, the true antitoxic antilipid A titres may actually be low in cirrhosis.

E N WARDLE

¹ Wardle, E N, *Clinical Science and Molecular Medicine*, 1974, **46**, 17P.

² Wardle, E N, *Archives of Surgery*, 1974, **109**, 741.

³ Coleman, M, et al, *Annals of Internal Medicine*, 1975, **83**, 79.

⁴ Waterlow, J, *Lancet*, 1950, **1**, 908.

Generalised urticaria precipitated by change to highly purified porcine insulin

SIR,—Allergic reactions to unpurified insulins are well recognised.^{1,2} Recently an allergic reaction to a porcine monocomponent insulin (Actrapid MC) has been reported.³ We wish to describe another case in which generalised urticaria developed after changing to highly purified porcine insulins and which has persisted despite the subsequent use of monocomponent preparations.

A 54-year-old man who has had diabetes for 18 years was restabilised on porcine insulin. He had received continuous treatment with bovine insulins from the onset of his diabetes and for many years had been taking a twice-daily admixture of soluble and isophane insulins (Weddel). Diabetic control had been far from adequate despite increasing doses of insulin (120 U daily) and following discussion it was felt that he might benefit from a more purified insulin preparation. Leo Neutral and Leo Retard insulins (Nordisk) were started and good diabetic control was achieved with a total of 84 U daily.

Five days after starting the new regimen he began to develop intermittent generalised urticarial reactions throughout the day and night which gradually increased in severity. No other drugs had been introduced during this period and he was admitted for further assessment. Systematic investigation failed to reveal any other underlying cause for the urticaria; the peripheral blood eosinophil count was normal. It is of interest that he gave a history of hay fever, but he has had no other particular allergies.

Insulin treatment was withdrawn and no new urticarial reactions developed during a 25-h period of observation. Treatment was recommenced with Actrapid MC and Semitard MC insulins (Novo), with a prompt return of urticaria, which has persisted despite trying different regimens, including a return to the original bovine preparations.

Intradermal skin tests were performed using 2 U (0.05 ml) of the following insulins: bovine soluble and isophane (Weddel); highly purified porcine Leo Neutral and Retard (Nordisk), and monocomponent porcine Actrapid and Semitard (Novo). In addition, 0.1 ml of Nordisk protamine and "medium" and of Novo methylparaben and "carrier" was injected. Equal volumes of saline were used for controls. At 30 min a 17-mm weal had developed in response to the Nordisk "medium." In addition, 8-mm weals occurred at the sites of the Nordisk Neutral and Retard insulins, protamine, and the bovine insulins. Minimal reactions to Novo carrier, methylparaben, Actrapid MC, and Semitard MC were observed, but these were not significantly different from the controls. All reactions, with the exception of the Nordisk "medium" and bovine isophane insulin, had faded within 4 h.

This patient's urticaria developed within five days of changing from conventional to

purified insulins and has persisted despite using different insulin regimens. That it was due to the insulin treatment was demonstrated by the absence of urticaria during insulin withdrawal. The possibility exists that the urticarial reaction reflects a delayed-type hypersensitivity, but immediate reactions could be demonstrated following intradermal skin tests. The occurrence of urticaria appeared to be triggered by the change to highly purified porcine insulins and it is speculated that a particular component of one of the preparations (? Nordisk "medium") acted as an adjuvant in this particular case. Fortunately, this patient's symptoms are reasonably controlled by chlorpheniramine tablets, and despite these difficulties his diabetic control has improved. Such reactions are rare complications of insulin treatment but both this report and that of Leslie⁴ indicate that they may occur in response to porcine insulins, and in both cases there was a strong history of allergy or hay fever.

We thank Mr F A P Henesey for his kind co-operation in providing the Nordisk protamine and "medium," and Dr D Leslie for providing the Novo methylparaben and "carrier."

C REISNER
D J MOUL
A G CUDWORTH

Diabetic Department,
St Bartholomew's Hospital,
London EC1

¹ Joslin, E P, Gray, H, and Root, A F, *Journal of Metabolic Research*, 1922, **2**, 651.

² Hanauer, L, and Batson, J M, *Diabetes*, 1961, **10**, 105.

³ Leslie, D, *British Medical Journal*, 1977, **2**, 736.

Legalisation of cannabis

SIR,—I am gravely disturbed by a further attempt to legalise the use of cannabis in Britain. Whatever argument is put forward by those interested in or merely indifferent to such a measure, there is universal agreement among the comparatively few pot smokers (both habitual and occasional) I have encountered as individuals that a person under the influence of cannabis is disinclined to go to work or look after his or her children and more than likely to fail to do this or any other useful work. Even more important, none of them would care to be driven in a car by such a person. These views are amply confirmed by such examinations as I have been able to make on persons so affected; they were certainly unfit to drive.

Have we not enough misery on these scores caused by alcohol that we must now be invited to add to it? The legalisation of cannabis cannot do good. It can only do harm. I look for a unanimous reiteration of this view from the profession.

A LEWIS

London W1

Confidentiality of medical records

SIR,—I was recently informed by a patient that, while attending a school medical examination with her son, she noticed that the school medical officer had in his possession copies of letters sent to me by a paediatrician at a local hospital. I made further inquiries and discovered that copies of letters from paediatricians concerning schoolchildren are sent automatically to area specialists in child health