ordinarily good prognosis raises the possibility that their patients were not typical of most with myocardial infarction and therefore the question whether the confident conclusion that heparin should be given to non-smokers after myocardial infarction is justified.

The point of great interest in their work is the apparent difference between smokers and non-smokers. Our critical comments about the need for heparin in no way detract from the interest which their work stimulates.

PETER CARSON
HELEN TUCKER

Cardiology Department,
City General Hospital,
Stoke-on-Trent, Staffs

¹ Tucker, H H, *et al*, *British Medical Journal*, 1973, **1**, 10.

*.*We sent copies of the above letters to Dr Emerson, whose reply is printed below.—ED, *BMJ*.

SIR,—I am grateful to Dr Richards for giving me the opportunity to make good the omission of the dosage of heparin that we used in our study. The prophylactic dose of heparin was 5000 U intravenously and 7500 U subcutaneously as soon as possible after diagnosis and then 7500 U subcutaneously every 12 h for 10 days.

Drs Carson and Tucker rightly point out the importance of early mobilisation in the prevention of thromboembolic complications. Early mobilisation, by the fifth day, was the policy in both our control and treated groups. While not disputing the desirability of early mobilisation, it is worth pointing out that most leg vein thromboses, as recognised by the fibrinogen uptake test, start in the first few days after the myocardial infarction so that early mobilisation can be only a limited answer to the problem.

We were most interested to hear from Dr J K Clayton and his colleagues (29 January, p 290) that in their study designed to predict postoperative deep vein thrombosis they found that smokers again appear less likely to develop postoperative deep vein thrombosis than non-smokers. If their observations are confirmed it will certainly give no support to our hypothetical explanation of the decreased incidence of postmyocardial and postoperative leg vein thrombosis in smokers. Perhaps smokers are more fidgety people and keep moving more—perhaps they provide their own very early mobilisation.

PETER EMERSON

Westminster Hospital,
London SW1

Multiple authorship

SIR,—I feel that papers published by as many as 17 authors¹ should be discouraged because of the inconvenience caused to those who may quote their work.

R E G SLOAN

Cheltenham

¹ Willis, A T, *et al* (Willis, A T, Ferguson, I R, Jones, P H, Phillips, K D, Tearle, P V, Fiddian, R V, Graham, D F, Harland, D H C, Hughes, D F R, Knight, D, Mee, W M, Pashby, N, Rothwell, Jackson, R L, Sachdeva, A K, Sutch, I, Kilbey, C, and Edwards, D), *British Medical Journal*, 1976, **1**, 607.

*.*The trial reported in the paper referred to by Dr Sloan was a collaborative effort to

which every one of the 17 authors contributed. In citation it is now common practice to follow the example of the *Index Medicus* and use the term "*et al*" when the number of authors exceeds three.—ED, *BMJ*.

Penicillamine therapy and teratogenesis

SIR,—Minerva's comments (26 February, p 581) on the prescribing of drugs to pregnant readers of middle-class, middle-brow newspapers will have evoked understanding smiles from many obstetricians. Her reference to a recent report¹ of 15 normal babies from 10 mothers with Wilson's disease treated with the chelating agent penicillamine will also be noted with considerable interest. This powerful drug is capable of producing a number of significant toxic effects and there has been concern about its possible teratogenicity, particularly from interference with collagen synthesis and maturation in the fetus as a result of the removal of metal ions.² Walshe's recent series¹ and a collected series³ of 29 normal infants from 18 mothers with Wilson's disease appears to indicate that penicillamine therapy in this condition carries no risk to the fetus.

It seems pertinent, however, to point out that in the treatment of cystinuria the safety of penicillamine therapy is not so definite and there has been one report⁴ of a generalised congenital connective-tissue defect in a baby born to a cystinuric mother treated with this drug. It has been suggested⁵ that this teratogenic effect may have resulted from the increased dosage of penicillamine used in this disease, but doubts have been raised about this conclusion by other reports^{5,6} of normal babies being born to cystinuric mothers taking penicillamine.

R CORCORAN
W J B CASTLES

St Catherine's Hospital,
Birkenhead, Merseyside

¹ Walshe, J M, *Quarterly Journal of Medicine*, 1977, **181**, 73.
² Marsh, L, and Fraser, F C, *Lancet*, 1973, **2**, 846.
³ Scheinberg, I H, and Sternlieb, I, *New England Journal of Medicine*, 1975, **293**, 1300.
⁴ Mjølnerod, O K, *et al*, *Lancet*, 1971, **1**, 673.
⁵ Laver, M, and Fairley, K F, *Lancet*, 1971, **1**, 1019.
⁶ Crawhall, J C, *et al*, *British Medical Journal*, 1967, **2**, 216.

Pemphigus erythematosus-like rash in a patient on penicillamine

SIR,—Penicillamine therapy may be considered an established regimen in cases of rheumatoid arthritis. As a consequence several unusual side effects have been reported over the years—for example, drug-induced lupus erythematosus,¹ myasthenia gravis,² glomerulonephritis with immune complex deposition,³ and pemphigus of the vulgaris and foliaceus varieties,⁴ though not other types of pemphigus. We wish to report the association of penicillamine treatment with the development of a pemphigus erythematosus-like rash in a case recently observed.⁵

A 49-year-old woman with seropositive rheumatoid arthritis has been under treatment since 1969; initially she received indomethacin and prednisolone (5 mg/day), then chloroquine for 15 months followed by azathioprine for six months. In January 1973 treatment with penicillamine was initiated, starting with 250 mg/day and increasing

to 750 mg/day over a period of three months. In addition she continued to receive 200 mg indomethacin/day. After 12 months this regimen was terminated because of the appearance of scaly and crusted lesions in the so-called seborrhoeic locations—that is, the midline regions of the trunk and scalp. At some places individual flaccid blisters could also be observed between the scaly manifestations. Penicillamine therapy was discontinued immediately. One week later skin biopsy showed subcorneal cleft formation and acantholytic cells within the cavities.

Direct immunofluorescence revealed deposition of IgG both in the intracellular area and along the basement membrane zone of lesional and perilesional skin. C3 could be detected intercellularly within areas of epidermal destruction only. (Antisera of goat origin labelled with fluorescein isothiocyanate were used. Anti-IgG conjugate was kindly provided by Dr E H Beutner, Buffalo, New York.) LE cell preparations, antinuclear antibodies and anti-double-stranded DNA assays (Farr) were negative at this time, as was the test for anti-epithelial antibodies.

Treatment with azathioprine (200 mg/day) and betamethasone (3 mg/day) was then started. Two months later we detected circulating antiepithelial antibodies (titre 1/10 using rhesus monkey oesophagus as substrate) and antinuclear antibodies (titre 1/40) which disappeared after two weeks. During this immunosuppressive treatment the skin lesions cleared within four months and have not recurred since.

We want to emphasise that skin reactions resembling not only pemphigus vulgaris and foliaceus but also pemphigus erythematosus may occur in patients under treatment with penicillamine.

O SCHERAK
G KOLARZ
K HOLUBAR

Departments of Medicine (II) and
Dermatology (I),
University of Vienna

¹ Camus, J P, *et al*, *Annales de Médecine Interne*, 1974, **125**, 9.

² Bucknall, R C, *et al*, *British Medical Journal*, 1975, **1**, 600.

³ Jaffe, I A, *et al*, *Annals of Internal Medicine*, 1968, **69**, 549.

⁴ Marsden, R A, *et al*, *British Medical Journal*, 1976, **2**, 1423.

⁵ Kolarz, G, *et al*, *Acta Medica Austriaca*, 1975, **2**, 130.

Vitamins and squamous metaplasia

SIR,—Your leading article (3 July 1976, p 2) about the association between vitamin A and squamous-cell carcinoma does not mention the important part that riboflavin plays in the maintenance of the integrity of squamous epithelia, particularly of the oesophagus.

Eight adolescent male baboons were fed a completely balanced and adequate synthetic diet containing proteins in the form of casein with all the essential amino-acids (20 % energy), carbohydrates (70 % energy), fats (10 % energy), and all the vitamins and mineral salts except riboflavin. After 160-300 days of vitamin B₂ deprivation they developed profound changes in the skin of the face, hands, legs, and feet, as well as more sinister ones in the oesophagus.^{1,2} This diet contained no analogues of any vitamins. All the baboons received 400 IU of vitamin A and 4000 IU of vitamin D. Two pair-fed and two animals on their natural diet were used as controls and had no abnormalities.

The squamous epithelial changes in these B₂-deprived baboons included hyperkeratosis, gross derangements of the normally highly organised process of keratinisation with acanthosis, extensive and impressive pseudocarcinomatous hyperplasia, thickened intradermal extensions of widely branching rete pegs, and aberrant individual cell keratinisation with intraepidermal epithelial pearl formation. Nuclear anaplasia was absent despite the strikingly increased epidermal nuclear labelling