spontaneous remission of the nephrotic syndrome after a course of 6 months in a Sephardic Jewish boy affected with renal amyloidosis associated with familial Mediterranean fever. The patient then remained free of any renal symptom, including proteinuria, for the next 5½ years until he experienced a dramatic relapse with a full-blown nephrotic syndrome and severe renal failure.¹ A renal biopsy performed after 3½ years of remission showed the persistence of amyloid deposits with a distribution and intensity similar to those seen when the first biopsy was performed, when the nephrotic syndrome was present.

The importance of such spontaneous remissions must be stressed in view of several recent articles ascribing to various drugs the disappearance of the nephrotic syndrome in renal amyloidosis.²⁻⁵

JEAN-PHILIPPE MÉRY

Service de Néphrologie,
Centre Pasteur-Vallery-Radot,
Paris

¹ Méry, J-P, and Mostefa, S, *Annals of Internal Medicine*, 1975, **83**, 581.

² Cohen, H J, et al, *Annals of Internal Medicine*, 1975, **82**, 466.

³ Horne, M K, *Annals of Internal Medicine*, 1975, **83**, 281.

⁴ Zemer, D, et al, *New England Journal of Medicine*, 1976, **294**, 170.

⁵ Ravid, M, Robson, M, and Kedar, I, *Annals of Internal Medicine*, 1977, **87**, 568.

Snakes and snake bite

SIR,—I was interested in the review by Dr David A Warrell of the book *Snakes and Snakebite* by John Visser and David Chapman (13 May, p 1269). As I have a special interest in this subject I would like to make one or two comments on the review which may be pertinent.

Dr Warrell has already made a number of observations about the book and its contents. However, the implication of the "futility of local interference at the site of the bite" is that pressure-relieving incisions play no part in the treatment of snake bite. I believe it has been amply demonstrated in several disciplines of medicine that pressure-relieving fascial incisions can prevent necrosis. A particularly good example of this is the anterior tibial compartment syndromes. The use of tourniquets is only controversial in as far as the dangers in the hands of lay people are concerned. If they are not applied within a few minutes of the bite they are unlikely to do any good at any time.

Dr Warrell correctly points out that puff adder bites can cause significant delayed hypotension and concealed blood loss, but his suggestion that the antivenom can help in preventing local necrosis appears unfounded in view of the fact that the antivenom does not contain any local cytotoxin-effective antibodies. The point to be emphasised is that antiserum need never be given for puff adder bites except in controlled circumstances and after admission to hospital. In my own experience the incidence of reaction to intravenous antivenom is low except in snake handlers who have previously been bitten and treated. Neither the authors nor the reviewer make any mention of prevention of serum sensitivity by the use of high-dosage cortisone. Although adrenaline is highly effective in the treatment of serum sensitivity, it is in itself a dangerous drug which should not be handled by lay persons, and if treatment with antivenom is always preceded by a bulk injection of a pharmaco-

logical dose of cortisone antiserum sensitivity can be expected to disappear completely; this has been our experience. Finally I would like to re-emphasise Dr Warrell's point that the doses of antivenom administered in most hospitals are quite inadequate and should not be used routinely.

J-J BROSSY

Department of Surgery,
Somerset Hospital,
Green Point, S Africa

SIR,—It would be interesting to know the evidence on which Dr D A Warrell (13 May, p 1269) makes the assertion that venom from a spitting cobra spat into the eye "frequently causes corneal ulceration, and may be absorbed . . . causing hypopyon and anterior uveitis."

There is no doubt that in travellers' tales the severity of such incidents has been repeatedly asserted, but my own limited experience¹ suggests that such sequelae are rare and that the consequences of such an event are relatively mild and unlikely to occasion permanent damage. Duke-Elder's comprehensive *System of Ophthalmology*² makes only limited reference to this particular problem. This again would tend to confirm one's impression that this particular snake's venom in the eye is perhaps no worse a mishap than accidental contamination with, say, neat washing-up liquid or one of the quaternary ammonium compounds—painful and discomfiting but not the major event suggested by the quotation.

It is a little difficult to envisage in a case of a simple "spit" a more effective treatment than local lavage and analgesia. One surely would be very reluctant in such a case to add the potential risk of necrosis by use of antivenom.

M J GILKES

Sussex Eye Hospital,
Brighton

¹ Gilkes, M J, *British Journal of Ophthalmology*, 1959, **43**, 638.

² *System of Ophthalmology*, ed S Duke-Elder, vol 14, part 2, p 1206. London, Kimpton, 1972.

Alcohol and the blood

SIR,—Your leading article (10 June, p 1504) provides a fine historical review of the blood changes that may be related to excess alcohol consumption. However, it gives little information about which of these abnormalities are common and which are rare, and no guidance in their use for the detection of alcoholism. Most of the studies referred to were made on "skid-row" alcoholics, who are not typical of patients with alcohol-related diseases seen in the UK.

The majority of our alcoholic patients with macrocytosis are well nourished,^{1,2} are in regular employment, and have normal vitamin B₁₂ and folate stores. Unger and Johnson³ found that 3% of an insurance company's employees were macrocytic on routine blood count and, of 17 examined, 16 were probably consuming excessive amounts of alcohol. A low serum folate concentration was found in only one subject, and although the remainder were treated with large doses of first folic acid and then B₁₂, their mean cell volume (MCV) did not fall. In addition, there have been several other recent reports on macrocytosis of alcoholism from the UK and France which give a similar picture.⁴⁻⁸

Further evidence that the rise in MCV may

be unrelated to folate deficiency has been provided by the deoxyuridine (dU) suppression test,⁹ which is a sensitive biochemical test for folate or B₁₂ deficiency performed on short-term bone-marrow culture. All eight of the alcoholic patients with macrocytosis studied by Wickramasinghe and Longland¹⁰ gave normal results in this test, and their erythrocyte folate and serum B₁₂ levels were subsequently found to be normal. This implies that the alcohol effect could not be attributed to an interference with B₁₂ or folate metabolism and we believe that alcohol has some other direct effect on the developing erythroblast. When folate deficiency does occur in alcoholics it is almost always due to associated nutritional folate deficiency, a feature universally present in the "skid-row" alcoholic but seen in only one-third of our patients, who are more likely to be spirit than beer drinkers.

Finally we would suggest that British practitioners are unlikely to see many alcoholic patients with sideroblastic anaemia, thrombocytopenia, or Zieve's syndrome but will often see patients with a normal haemoglobin concentration, a normoblastic bone marrow, normal B₁₂ and folate levels, and a raised MCV that returns to normal after three or four months of abstinence from alcohol.

D M CHALMERS

I CHANARIN

A J LEVI

Divisions of Haematology and
Gastroenterology,
Northwick Park Hospital and Clinical
Research Centre,
Harrow, Middx

¹ Wu, A, Chanarin, I, and Levi, A J, *Lancet*, 1974, **1**, 829.

² Wu, A, et al, *British Journal of Haematology*, 1975, **29**, 469.

³ Unger, K W, and Johnson, D, *American Journal of the Medical Sciences*, 1974, **267**, 281.

⁴ Buffet, C, *Revue de l'Alcoolisme*, 1976, **22**, 15.

⁵ Wright, S G, and Ree, G H, *Lancet*, 1978, **1**, 49.

⁶ Khaund, R R, *Lancet*, 1978, **1**, 327.

⁷ Davidson, R J L, and Hamilton, P J, *Journal of Clinical Pathology*, 1978, **31**, 493.

⁸ Whitehead, T P, Clarke, C A, and Whitfield, A G W, *Lancet*, 1978, **1**, 978.

⁹ Metz, J, et al, *British Journal of Haematology*, 1968, **14**, 575.

¹⁰ Wickramasinghe, S N, and Longland, J E, *Acta Haematologica*, 1974, **52**, 14.

Nutrition and the brain

SIR,—Your leading article (17 June, p 1569) presents an informative analysis of the present state of knowledge and uncertainty as to the relationship between nutrition and cerebral development. The general approach is factually based, with a brief review of some of the relevant data. In sharp contrast is your curt dismissal of "unscrupulous" attempts to link malnutrition to "mental retardation in its usual sense."

Severer degrees of mental handicap (profound, severe, and moderate) occur at about 4 per 1000 in populations aged 10-16 years in Britain, as shown by a number of careful surveys. While it seems improbable that such severe mental handicap can be specifically or commonly due to malnutrition in a relatively developed country, the sources which you quote, and others, still leave open the possibility in cases of prolonged maternal or severe infant malnutrition elsewhere. However, the majority of the mentally retarded singled out for help from the special services in Britain or the USA are the mildly retarded, as defined by the International Classification of Diseases or the American

Association on Mental Deficiency (AAMD). The concept of mild retardation is quantitative and embraces, using conventional standards, some 2.2% of the population in addition to the 0.4% who are even less gifted. Not all of this considerable population is in receipt of special services, even at school age; but prospects for training for skilled employment and independent social adaptation tend to be limited. Fortunately, the AAMD has omitted from the revised classification of mental handicap¹ the term "borderline mental retardation," but this is still in common use and embraces much of the "dull normal" stream.

It seems contrary to the general sense of the leading article thus arbitrarily to exclude any causal role in mental retardation to malnutrition when this can admittedly cause some deficiency in the developing brain. It would seem more reasonable in the present state of science to keep an open mind on this question and to view it as one of the many possible modes of operation of the impact of adverse social and economic factors among underprivileged groups and disadvantaged classes.

BRIAN KIRMAN

Queen Mary's Hospital for Children,
Carshalton, Surrey

¹ *Manual on Terminology and Classification in Mental Retardation*, ed K J Grossman. Washington, DC, American Association on Mental Deficiency, 1973.

Factors affecting length of hospital stay

SIR,—May I comment on Mr H B J Chishick's second letter (17 June, p 1622) as his first (29 April, p 1145) referred to a letter of mine (25 March, p 789)?

He seems to share the common delusion that the only obstacle to reducing the waiting time for cold surgery is "unnecessary delay in discharging patients." This is almost never true because the real limiting factor nowadays is the work capacity of the operating theatres and anaesthetic services (perhaps not the latter in cataract surgery). To dwell on figures of bed occupancy, turnover intervals, mean duration of patient stay, or cost per patient is a great waste of many well-meaning people's time, and generates a vast amount of totally unproductive heat.

ROGER HOLE

Department of Urology,
North Ormesby Hospital,
Middlesbrough, Cleveland

Effects on fetus of stilboestrol in pregnancy

SIR,—With reference to the paper by Mr J M Monaghan and Mr L A W Sirisena (17 June, p 1588) your readers may be interested to know of research carried out by Henderson *et al*¹ in the USA into whether there was a higher incidence of abnormalities in the sons of women who had taken diethylstilboestrol (DES) during their pregnancies as compared with the sons of controls.

They found that the only significant difference was in the incidence of diseases of the urogenital tract—55 out of 225 exposed males (25%) compared with 17 out of 111 controls (15%). Their most striking finding was the difference in incidence of abnormalities of the penile urethra—10 of the exposed males compared with none of the unexposed males. These abnormalities con-

sisted of one case of hypospadias (the number which could be expected among the total of 306 males surveyed) and nine cases of urethral stenosis, a finding which was highly significant.

No cases of malignancy have been reported, but this is presumably not yet ruled out as a possibility owing to the youth of the subjects involved. No systematic follow-up of such patients is possible in the USA since there is on centralised record keeping. This could perhaps be undertaken in the UK since it appears that many women who took DES are already known and could therefore be contacted for information concerning their sons.

VALERIE HARTLEY-BREWER

London N3

¹ Henderson, B E, *et al*, *Pediatrics*, 1976, **58**, 505.

Inappropriate use of tricyclic antidepressants

SIR,—The short report by Dr J C Little and others (17 June, p 1593) pointing out their difficulty in obtaining depressed patients for a clinical trial raises an important question. They requested general practitioners to refer those patients whom the GP thought suitable for tricyclic antidepressant therapy. The table in the report shows that 12 of the 35 patients referred (33%) were either unsuitable for treatment with tricyclics or were suffering from conditions in which tricyclic treatment was actually contraindicated. If these patients had not been referred for these specific investigations their GPs would presumably have initiated tricyclic therapy and this suggests that there may be a disturbingly high inappropriate use of these drugs in general practice.

JOHN R KIRWAN

Harefield Hospital,
Uxbridge, Middx

"Innovation in the Pharmaceutical Industry"

SIR,—Dr B W Cromie (17 June, p 1618) corroborates the views expressed at a Council for International Organisations of Medical Sciences/World Health Organisation meeting six months ago on "Trends and prospects in drug research," at which speakers with a knowledge of the pharmaceutical industry were uniformly pessimistic.¹

As he indicates, there are many contributory factors besides the drug control authorities. However well intentioned individually, the constraints have collectively created a paradoxical situation in which the industry is no longer able to develop the drugs that society needs. The cost, delays, and uncertainties make drug research progressively less attractive and the large multidivisional enterprises that make up the research-based industry now see other more promising areas for investment than pharmaceuticals. Once they start to run down their investment the industry will rapidly decline and would take years to rebuild.

In the past the industry has attracted a great deal of criticism, but it has largely been responsible for the tremendous therapeutic advances of the past 30 years. It still offers the best prospect for further progress, but if the profession, politicians, and public want progress to continue they must promote or at

least permit a climate in which a vigorous industry can operate. This is the fundamental problem and it is urgent.

T B BINNS

Department of Pharmacology and
Therapeutics,
London Hospital Medical College,
London E1

¹ *Lancet*, 1977, **2**, 1335.

Utility of needle aspiration of tumours

SIR,—One statement bothers me in your excellent leading article on this subject (10 June, p 1507) although I admit that from the surgeon's point of view it is probably true that "when challenged, most of those who seek cytological specimens would agree that a histopathological diagnosis is necessary before ablative surgery or other radical treatment, such as chemotherapy or radiotherapy." However, from the cytopathologist's point of view when surgical biopsy fails to confirm a positive diagnosis of carcinoma made on needle aspiration cytology biopsy of the tumour must be repeated in order to find the malignant tissue. False-positive cytological diagnoses of cancer are very rare in my experience, giving this procedure a specificity of almost 100%. The drawbacks of aspiration cytology compared with surgical biopsy are (1) its lower sensitivity—that is, more false-negative reports—and (2) its narrower diagnostic facility—for example, adenocarcinoma cannot be so readily differentiated from squamous carcinoma. On the other hand it needs no anaesthetic and it can be done on outpatients at negligible cost.

P A TROTT

Department of Pathology,
Royal Marsden Hospital,
London SW3

Factors influencing the incidence of neonatal jaundice

SIR,—Dr B S B Wood (3 June, p 1488) criticises our study on neonatal jaundice (13 May, p 1235) because it was retrospective. However, while accepting that a prospective study is preferable, the large number of cases included strengthens the value of our conclusions and a prospective study of 12 000 neonates would be expensive and time consuming to carry out. Our arbitrary allocation of an average normal bilirubin level to those infants in whom the measurement was not considered necessary is said to be "not acceptable." This procedure merely allowed us further to quantify the importance of the different factors found to be independently associated with changes in the jaundice rate and it did not bias the identification of these factors.

Dr Wood reports that breast-feeding produces a twofold increase in jaundice and we did not study the influence of breast-feeding. We have now been able to add this factor to our analysis for 9300 babies born between 1971 and 1975. The breast-feeding rate was 55% and significant jaundice ($\geq 205 \mu\text{mol/l}$ ($\geq 12 \text{ mg/100 ml}$)) was detected in 13.2% of breast-fed and 11.4% of bottle-fed infants. Multivariate analysis confirmed breast-feeding as an independent factor associated with increased jaundice, but the estimated average increase in peak bilirubin concentration due to it was only $5 \mu\text{mol/l}$ (0.3 mg/100 ml). This