sound recommendations may be modified by recent events in the United States.

The US Senate Select Committee on Nutrition and Human Needs reported last February.¹ It had been taking evidence from doctors at public sessions since July 1976.² The committee has recommended many dietary changes. In order of priority these changes are: (1) increase consumption of starchy carbohydrates, called complex carbohydrates, from 22% of energy intake at present to 40-45% and increase consumption of whole grain products and also of fruit and vegetables, for the fibre from both is "important in nutrition"; (2) decrease sugar consumption from 24% to 15% of energy intake; (3) decrease consumption of total fats from 42% to 30%, saturated fats from 16% to 10%, and polyunsaturated and monounsaturated from 26% to 20%; (4) protein no change at 12%.

The evidence²⁻⁴ to support this enormous and difficult dietary change has been implemented by legislation on 31 March to create a new Bureau of Human Development.⁵ It has been funded by hundreds of millions of dollars. The committee is continuing to hold sessions and another report, this time on dietary fibre and health, has been promised.⁶

HUGH TROWELL

Woodgreen,
Fordingbridge, Hants

¹ US Senate Select Committee on Nutrition and Human Needs. *Dietary Goals for the United States*. US Government Printing Office, Washington, 1977.

² US Senate Select Committee on Nutrition and Human Needs. *Diet and Killer Diseases with Press Reaction and Additional Information*. US Government Printing Office, Washington, 1977.

³ *Hearings before the US Senate Select Committee on Nutrition and Human Needs. Part 1. Cardiovascular Disease*. US Government Printing Office, Washington, 1977.

⁴ *Hearings before the US Senate Select Committee on Nutrition and Human Needs. Part 2. Obesity*. US Government Printing Office, Washington, 1977.

⁵ A Bill to amend the Public Health Service Act to establish a Bureau of Human Development. *US Congressional Record*, vol 123, No 57, 31 March, 1977.

⁶ Senator George McGovern, personal communication, 1977.

Mycoplasma and encephalitis

SIR,—I was interested in the report of meningo-encephalitis associated with *Mycoplasma pneumoniae* infection (by Dr P Kopelman (2 April, p 881)) and the author's remarks concerning pathogenesis. It is relevant that sera of patients infected with *M. pneumoniae* show IgM antibodies that react with brain, since there are common glycolipid antigens.¹

E N WARDLE

Newcastle upon Tyne

¹ Kaplan, M H, and Rapaport, F T, *Proceedings of the First International Congress of Immunology*, p 1153. London, Academic Press, 1971.

Management of childhood epilepsy

SIR,—To manage means to control. As every schoolboy knows, drugs do not control childhood epilepsy, so one hoped that your leading article under the above title (16 April, p 986) would have been about something other than them. The control of the household, for instance, and of going to school, of climbing trees and swimming and of lighting fires and going camping; and of the prospect of marriage and pregnancy and rearing a family and

eugenics. All this was left out, and instead we had a little cliché about sympathetic advice.

That epileptics fall in the fire and char their limbs away; that by even lying on hot piping for an hour they can irretrievably damage them; that they drown in a bath three inches deep; that it is impossible, when swimming out of your depth, to support a child above water level for the duration of a major fit and that inhalation of river water is fatal: none of these was mentioned. No fire guards, no protection of staircases, no removal of bathroom locks, but only sadly ineffective, and sadly poisonous, drugs were discussed. It is as if your expert on epilepsy saw patients only in a hospital clinic, left the clinic only to go to the laboratory, and never saw the results in the home, in plastic surgery wards, and in the cemeteries, of such neglect of management.

H WYKEHAM BALME

London EC2

Methadone: evidence of accumulation

SIR,—Arising from the letter of Mr D M Rutherford and Dr K Raymond (19th March, p 774) we would firstly like to apologise for our inadvertent solecism in misspelling Mr Rutherford's name on two occasions in our previous letter (5 February, p 375).

We were most interested in their finding that the half life of methadone was 12-30 h. This would appear to bear out our comments on the variability which occurs in the published figures for this value.

It is of course well known that if a drug is administered repeatedly at constant time intervals it will accumulate until a "plateau level" is reached, when the concentrations fluctuate between a maximum and minimum level which can be calculated with precision by the method of Dost.¹ We would therefore agree with their introductory comments. However, we cannot accept their comments on the methadone blood levels which may originate from a misinterpretation of the graph published in our letter: If the whole-blood methadone concentrations of the three subjects are considered from the time of peaking over the longer 12-h period assessed on day 1 of the study the figures shown in the table are

Whole-blood methadone levels (ng/ml) after administration of one controlled-release tablet containing 10 mg of methadone

Time (hours)	Subject 1	Subject 2	Subject 3
1	—	—	58
2	54	89	34
3	48	60	27
6	44	55	25
8	41	50	23
12	32	33	18

obtained. Although the idealised flat blood concentration curves were not attained, as indeed they rarely are in practice, the results after peaking were very close to this theoretical concept. The release mechanism of the tablet was in fact designed to produce rapid analgesic activity through a measured loading dose and also to complement these initial levels by a further controlled release of methadone.

However, our main object in replying is not to argue the minutiae of the bioavailability pattern of our controlled-release methadone tablet but to highlight the problems of

accumulation occurring with methadone whether in a controlled, non-controlled, or liquid form. We found that the accumulation that occurred even during our short study gave rise to considerable side effects in the young, healthy volunteers. This accumulation, even if not indefinitely progressive, can thus give rise to problems even before a steady state is reached. Obviously this could be dangerous in elderly patients—as exemplified by Dr Paul Symonds's patient (19 February, p 512) who became unconscious with symptoms of narcotic overdose after the administration of 5 mg of methadone three times a day for two days. Such patients, often additionally exhausted by unrelieved pain or severe coughing, are vastly different from either healthy volunteers or young addicts, whom they also vastly exceed in number. In these elderly patients the risk of methadone accumulation and toxicity represents a substantial hazard.

We are aware that other physicians have encountered problems with methadone accumulation and may wish to comment in your columns.

S T LESLIE
R B MILLER
C BORODA

Research Division,
Napp Laboratories Ltd,
Watford, Herts

¹ Dost, F H, *Grundlagen der Pharmakokinetik*, Stuttgart, Georg Thieme, 1968.

Hospitable Trinidad

SIR,—Recently we three, all final-year medical students, spent elective periods attached to the University of the West Indies based in Port of Spain, Trinidad. We were all able to choose the departments in the hospital, health service, or research centre in which we wished to study. The many members of the medical services we met spent a great deal of time and effort teaching us and seeing to our general welfare. We are very grateful to them all. Our time in Trinidad proved an extremely valuable experience.

Arranging such elective studies is an obvious financial strain on any medical student. Our problems were greatly lessened by the fact that we were provided with rooms and meals at Port of Spain General Hospital free of charge. We would like to urge all British medical schools to extend similar hospitality (as we believe some already do) to University of the West Indies students and others on electives so that more medical students may have the chance of an experience such as ours. Our thanks to all in Britain and Trinidad who made our electives possible.

VALERIE A HARPIN

Middlesex Hospital Medical School

NIGEL ZOLTIE

Bristol University Medical School

BRIAN FERNANDES

Edinburgh University Medical School

Visual and manual examination of whole stools in plastic bags

SIR,—We should like to draw attention to a simple method for visual as well as tactile examination of faeces which we developed several years ago in order to detect remainders from oxytetracycline capsule medication. The