CORRESPONDENCE

Rubella vaccination Catherine S Peckham, MD, and others....	1278	Publicity and Huntington's chorea Kathleen Evans	1282	Inactivation of penicillins and cephalosporins by enzymes R W Lacey, MD.....	1285
Drug compliance in hypertensive patients A J Marshall, MRCP, and D W Barritt, FRCP	1278	Concentration of thyroxine-binding globulin R Mardell, BSC.....	1282	Morale and management in general practice J S Berkeley, FRCGP	1285
Haematemesis and melaena D J Allison, FRCR.....	1279	Localised airway obstruction P Lockwood, MSC.....	1283	Bed occupancy: misleading figures T L Barclay, FRCSed, and G A Craig, FRCOG	1285
Biliary atresia in childhood A P Mowat, FRCP, and E R Howard, FRCS; K J Foster, MRCP, and S J Karran, FRCS...	1279	Prevention priorities R S Savidge, FRCP.....	1283	Differentials in payment of hospital staff A Watson, FRCS.....	1286
Commitment to oncology K E Halnan, FRCP.....	1280	Breast size and cancer A J Robertson, MB, and others.....	1283	Reduction of waiting lists G T Watts, FRCS.....	1286
Speed limits and the public health J L T Birley, MRCP; A K Clarke, MRCP; L W Lauste, FRCS.....	1280	The bedpan round E Mary Grant, SCM.....	1283	Hospital practitioner anaesthetists: requirements for appointment W J M Hood, MB, DA; M Marshall, FFARCS; Jennifer R Jenkins, FFARCS.....	1286
Staffing in the medical laboratory service A R Haxton, FIMLS; A M Grant, MRCP....	1280	Colour blindness and arithmetic L H G Moore, MB.....	1283	Whole-time consultants and the contract M M Burrows, FFARCS.....	1286
Laparoscopic sterilisation with Silastic bands A M Smith, FRCOG.....	1281	Diet and coronary heart disease Rev H C Trowell, FRCP.....	1283	Professional indemnity G Russell, MRCP.....	1287
Swimming goggles causing severe eye injuries Flight Lieutenant R J Lamb, MB.....	1281	Mycoplasma and encephalitis E N Wardle, MD.....	1284	Salaried doctors in Europe A J Rowe, MRCGP.....	1287
Disseminated histoplasmosis in Britain D W R Mackenzie, PHD.....	1281	Management of childhood epilepsy H W Balme, FRCP.....	1284	Points from Letters Diet and coronary artery disease (P E Moffitt; H Watson); Haematemesis and melaena (R G Potter); Seat-belt injury? (P K P Harvey); Orthopaedic services in Lincoln (P J Ellis); An IUCD record? (C M Flood); Impeccable syllable (Jean Wilson).....	1287
"Experience not essential" S S H Wasty, LAH.....	1282	Methadone: evidence of accumulation S T Leslie, MPS, and others.....	1284		
Mobility for the disabled T Marshall, MSC.....	1282	Hospitable Trinidad Valerie A Harpin, and others.....	1284		
		Visual and manual examination of whole stools in plastic bags L van Wijk, MD, and J Wieriks, MD.....	1284		
		Partial floral suppression D O Lewis, MRCPed.....	1285		

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Rubella vaccination

SIR,—The suggestion made by Drs C G D Brook and E M Ross (16 April, p 1028) that rubella vaccination in the United Kingdom should be switched to conform with the mass vaccination policy as practised in the United States "forthwith" rather than putting increased effort into the present policy of selective immunisation must not pass unchallenged. Rather than being as simple as they suggest, the whole problem of prevention of congenital rubella by immunisation is exceedingly complex.

The immunisation programme which they advocate is designed to produce herd immunity and thereby prevent the acquisition of rubella by susceptible pregnant women. Attempts to produce herd immunity for rubella with vaccines have been unconvincing and there is no epidemiological evidence that this has been or can be achieved.¹⁻³ It should also be made clear that the original recommendation for the use of rubella vaccine in the United States has now been modified by recommending that the programme of vaccination of adolescent girls and adult women should be extended. Thus they are also using selective immunisation of women of childbearing age similar to the practice in this country.

Before the incidence of congenital rubella in the United States can be assessed the adequacy of the method of surveillance of

congenital rubella must be carefully evaluated. The fact that relatively little congenital rubella has been recognised in the United States since the widespread vaccination of children began in 1972 could also be accounted for by the infrequency of epidemics of rubella, declining birthrates, and the increased availability of induced abortions.⁴

Immunisation of girls aged 11-14 years has certain advantages: (1) Approximately half the girls will already have acquired natural infection and will therefore have solid protection for the future. (2) Another reason for delaying immunisation to the 11-14-year-old group is that we still do not know how long vaccine-induced immunity will last in comparison with natural immunity from rubella. The need for a booster dose when vaccine is given in very early childhood cannot be discounted. (3) Clearly there is greater interest among the girls and their parents to protect them against rubella than would be the case for boys. (4) Selective immunisation is far less costly.

It is not yet possible to evaluate fully either the British policy of selective immunisation or the United States policy of mass immunisation. However, for the above reasons we believe that the present immunisation programme in the United Kingdom is the correct one and every effort should be made by members of

the medical profession and others concerned with immunisation to promote the use of rubella vaccine for all groups for whom it is currently recommended. The national acceptance rate is at present far below the desired level.

CATHERINE S PECKHAM
W C MARSHALL
J A DUDGEON

Department of Microbiology,
Institute of Child Health,
London WC1

¹ Klock, L E, and Rachelefsky, G S, *New England Journal of Medicine*, 1973, **288**, 69.

² Guyer, B, et al, *Journal of the American Medical Association*, 1974, **227**, 1298.

³ Bolognese, R J, and Corson, S L, *Obstetrics and Gynecology*, 1973, **42**, 851.

⁴ Schoenbaum, S C, *Journal of the American Medical Association*, 1976, **235**, 248.

Drug compliance in hypertensive patients

SIR,—Your leading article on adherence to prescribed drug therapy (26 March, p 793) is timely. The realisation that the treatment of asymptomatic hypertension improves prognosis has created another large group of patients who will face years of swallowing tablets.

We studied 24 hypertensive patients, treated with one of the simplest of hypotensive regimens, over a 26-week period. Their ages ranged from 28 to 65 years. They were entered into a double-blind comparison of atenolol with placebo, given as matching tablets, firstly twice a day and then once daily. We used all the aids to improve drug compliance described in your article, assessed the patients at two-

weekly intervals, and reinforced the importance of treatment. The tablets were dispensed in a tin which contained 12 tablets more than required for each two weeks. A treatment period was designated as compliant if 80% or more of the tablets had been removed. All tablet counts were made at the end of the trial. No patient defaulted from attendance and all the tablet containers were returned as requested. The tables show the number of times the 80% value was achieved. Adherence to therapy was marginally better with atenolol and with once-daily treatment, but the differences were not statistically significant by the χ^2 test. No patient consistently failed to comply but only one complied throughout all periods of the trial. Side effects to atenolol were inconspicuous and were not associated with periods of drug-defaulting. Despite our attempts we failed to reach our goal for consistent tablet-taking one-third of the time. Tablet counts are indeed likely to lead to an overestimate of compliance.

TABLE I—Compliance on active and placebo tablets (twice-daily treatment regimen)

	Total No of analyses	Proportion (and No) of analyses designated compliant
Atenolol ..	112	66% (74)
Placebo ..	58	60% (35)

TABLE II—Compliance on once- and twice-daily treatment

	Total No of analyses	Proportion (and No) of analyses designated compliant
Twice daily ..	170	64% (109)
Once daily ..	35	72% (24)

Hypertension does present special problems as it is a chronic disease requiring treatment in the absence of symptoms. Clearly there is little point in developing more sophisticated pharmacological methods of treatment if patients cannot be persuaded to take medication reliably. Our findings emphasise the need for caution when using hypotensive agents, such as clonidine, known to be sometimes associated with rebound hypertension when stopped suddenly.¹ Compliance has also to be considered in the clinical evaluation of any hypotensive agent and in the interpretation of treatment and mortality statistics.

More attention needs to be given to long-term tablet-taking. It is particularly the case in the treatment of hypertension that failure of the patient to take his tablets regularly leads not only to waste but also to loss of confidence of the doctor in his chosen drug. Drug treatment of hypertension has become simpler² and it is now appropriate to test carefully whether patients are more likely to take tablets once or twice daily and whether calendar packing is feasible and preferable.

ANDREW MARSHALL
D W BARRITT

Bristol Royal Infirmary,
Bristol

¹ Hunyor, S N, et al, *British Medical Journal*, 1973, 2, 209.

² Marshall, A J, et al, *Lancet*, 1977, 1, 271.

Haematemesis and melaena

SIR,—It was disappointing to note that your recent leading article on upper gastrointestinal bleeding (16 April, p 989) contained no reference to any of the radiological techniques now available for controlling such bleeding. These include the infusion of vasoconstrictor agents selectively into the coeliac or superior mesenteric arteries,¹ the injection of emboli into the left gastric or gastroduodenal arteries through percutaneous arterial catheters,²⁻⁴ and the injection of vasoconstrictors, sclerosants, or emboli into bleeding varices via percutaneous transhepatic portal catheters.^{5, 6}

At the Hammersmith Hospital we have recently successfully embolised the gastroduodenal artery in two patients, both shown at endoscopy to be bleeding from a duodenal ulcer. These patients were considered by our clinical colleagues to be "poor operative risks" and it was preferable to arrest their bleeding by means of a relatively short and painless procedure under local anaesthesia rather than a potentially hazardous laparotomy under general anaesthesia.

The use of such radiological techniques is now almost routine in some centres abroad, particularly in North America and Scandinavia, and I suggest that your proposed "specialised team" for the management of patients presenting with haematemesis and melaena should include a radiologist.

D J ALLISON

Department of Diagnostic
Radiology,
Royal Postgraduate Medical School
and Hammersmith Hospital,
London W12

¹ Rossi, P, and Taverna-Giardina, L, *Rendiconti di Gastroenterologia*, 1975, 7, 70.

² Reuter, S R, Chuang, V P, and Bree, R I, *American Journal of Roentgenology*, 1975, 125, 119.

³ Goldman, M L, et al, *Radiology*, 1976, 120, 513.

⁴ McCloy, R F, and Allison, D J, *Lancet*, 1977, 1, 647.

⁵ Scott, J, et al, *Lancet*, 1976, 2, 53.

⁶ Dick, R, *British Journal of Radiology*, 1977, 50, 241.

Biliary atresia in childhood

SIR,—We would like to express our concern at the misleading title to the interesting case report by Dr K J Foster and Mr S J Karran (26 March, p 815).

Atresia (Greek, *a*, negative + *trésis*, a hole), is usually applied to a congenital absence or closure of a normal opening, passage, or cavity. In biliary atresia there is complete obstruction, destruction, or absence of part or all of the bile ducts. The operative cholangiogram illustrating this case report, however, does not show obstruction, although the common bile duct is narrowed. This finding correlates with the preoperative bilirubin level of only 22 μ mol/l (1.3 mg/100 ml).

The state of the bile ducts at the initial laparotomy is more difficult to assess from the report, but if bile was obtained when the common bile duct was opened and a biliary fistula developed the proximal bile ducts must have been in continuity with the intrahepatic ducts. The spontaneous closure of the biliary fistula and the subsequent clinical course suggest that the distal common bile duct was also patent. Unfortunately operative cholangiography, which would have settled this point, was not performed or not reported. We would suggest that liver biopsy at that time may have shown an intrahepatic cause for jaundice with reduced bile flow and narrow bile ducts. We would hesitate, therefore, to

accept that this child had, at 5 months of age, biliary atresia which subsequently underwent spontaneous recanalisation.

The reports of Holder¹ and Krovetz,² cited in the article as evidence for spontaneous recanalisation in biliary atresia, provide unconvincing evidence. Holder's article refers to an earlier report by Clatworthy and Macdonald,³ who in fact made no claim for recanalisation, while the case of Krovetz was eventually considered by that author to be one of intrahepatic biliary hypoplasia, a different entity.

We appreciate that the correct characterisation and nomenclature of biliary tract disorders are confusing but recommend that wherever possible the internationally agreed nomenclature be used.⁴ A more appropriate title for this case report would have been: "Cholecystojejunostomy: an unconventional but successful treatment for pruritus complicating cirrhosis (morphology and aetiology undetermined)."

It would be very unfortunate if this case report were to cause unnecessary delay in the investigation of infants with suspected extrahepatic biliary atresia. The majority of patients with confirmed extrahepatic biliary atresia still die before 2 years of age, although the exceptional case may survive until late childhood with severe jaundice and cirrhosis.⁵ Where the atresia is distal and the proximal bile duct patent long-term survival is still exceptional, but the success rate is improved by early surgery.⁶ The operation of hepatic portoenterostomy has been followed by long-term survival in patients with atresia of the proximal extrahepatic bile ducts, but the frequency of successful bile drainage falls if surgery is delayed beyond 60 days of age.⁷

In any infant with conjugated hyperbilirubinaemia and acholic stools biliary atresia must be excluded as rapidly as possible in a unit suitably equipped and staffed to investigate such infants.⁸

ALEX P MOWAT
EDWARD R HOWARD

Department of Child Health,
King's College Hospital Medical
School,
London SE5

¹ Holder, T, *American Journal of Surgery*, 1964, 107, 458.

² Krovetz, J, *Surgery*, 1960, 17, 453.

³ Clatworthy, H W, and MacDonald, V G, *Surgical Clinics of North America*, 1965, 36, 1545.

⁴ *Diseases of the Liver and Biliary Tract. Standardisation of Nomenclature. Diagnostic criteria and diagnostic methodology.* DHEW Publication No (NIH)76-725. Bethesda, Maryland, National Institutes of Health, 1976.

⁵ Mowat, A P, Psacharopoulos, H T, and Williams, R, *Archives of Disease in Childhood*, 1976, 51, 763.

⁶ Borenson, M M, Garde, A R, and Moody, F G, *Gastroenterology*, 1974, 66, 260.

⁷ Kasai, M, Watanabe, I, and Ohno, R, *Journal of Pediatric Surgery*, 1975, 10, 173.

* * * We sent a copy of this letter to Dr Foster and Mr Karran, whose reply is printed below.
—ED, BMJ.

SIR,—Thank you for the opportunity to comment on the letter from Dr Mowat and Mr Howard. We were interested in their comments.

We felt that the case presented had sufficiently unusual features to warrant publication in a journal of general interest. Unfortunately some information was incomplete so we elected to present the case as a short report. Accordingly discussion was limited and the title chosen to be instantly recognisable to the general reader.