defined where the population of points was densest to the very sparsely populated area concerned. A further test was obviously required: the triiodothyronine (T3) level. The T3—Thyopac-3 pairs plotted on a T3 versus Thyopac-3 map fell in each case clearly in the euthyroid region, though again some extrapolation was necessary because of the high Thyopac-3 values. The resolution of the difficulty here reflects the well-known superiority of the level of T3 (which should be related correctly in some manner to Thyopac-3 or TBG for maximum discrimination) over that of T4 as a discriminator of high thyroid hormone output states. Only one of the sets of results seems likely to have been caused by familial high TBG levels; the others were caused by identifiable factors (pregnancy or oestrogen administration).

RICHARD MARDELL

Wessex Regional Department of
Nuclear Medicine,
Southampton General Hospital,
Southampton

Localised airway obstruction

SIR,—Your leading article entitled "Localised airway obstruction" (12 March, p 669) should strictly have been called "Upper airway obstruction" in accordance with established usage and as shown by Harrison, whom you quote.¹ The use of the term "localised airway obstruction" should be restricted to describe an obstruction in an airway at any level distal to the carina. Thus the term implies that as well as the localised obstruction, causing localised reduction in airflow, there is also some other airway without a reduced airflow. This situation is most commonly encountered in carcinoma of the bronchus.

Since generalised airways obstruction is accepted as the greatest single cause of post-operative cardiopulmonary complications in surgery for carcinoma of the bronchus the assessment of the functional state of the lungs prior to thoracotomy in this condition resolves into displaying whether or not there is significant generalised airways obstruction attributable to bronchitis as well as the localised airway obstruction due to the tumour. The situation was first put in those terms in 1964 by Simonsson and Malmberg,² and investigations since then have been directed at improving methods of discriminating between the contributions to lung dysfunction of the two disease processes.

P LOCKWOOD

Respiratory Physiology Department,
Harefield Hospital,
Middlesex

¹ Harrison, B D W, *Quarterly Journal of Medicine*, 1976, 45, 625.

² Simonsson, B G, and Malmberg, R, *Thorax*, 1964, 19, 416.

Prevention priorities

SIR,—Your leading article (16 April, p 989) supports the false hope of the House of Commons Expenditure Committee that "preventive medicine might save money." Surely "investment in prevention" will never "pay off in financial terms"—not even in "15 or 20 years."

It is likely that man will always die sometime, and if in old age the more will he be subject to illness, the more drugs will he

consume, and the more appliances will he require, and time spent in a long-stay geriatric ward will cost more than a pneumonectomy.

Prevention certainly, and the necessary finance must be found, for we are a caring society, but in the interest of its members in 20 years' time the BMA should warn MPs now that adequate health care of an older population will cost more. The Government will also need to finance more pensions⁷

RONALD SAVIDGE

Woodbridge, Suffolk

Breast size and cancer

SIR,—We were most interested to read the letter from Professor B N Catchpole (19 March, p 776) indicating the present lack of information on breast volume in relation to cancer.

As part of a study on breast carcinoma we have been recording breast volume and computed tumour volume,¹ in all stage I-III (TNM classification) patients undergoing mastectomy. Our present results, from 14 cases, show a breast volume range of 200-1150 (mean 562) cm³, a tumour volume range of 0.36-23.4 (mean 7.95) cm³, and a tumour : breast volume ratio, expressed as a percentage, of 0.08-6.78 (mean 2.18).

There has been no evidence so far of a correlation between any of the values recorded, the clinical stage, or the aggressiveness of the tumour as assessed by the percentage lymph node involvement by metastases. It remains to be seen whether long-term studies will reveal breast volume to be of importance in the eventual outcome of the disease.

ALISTAIR J ROBERTSON

J CLARK

A W HALL

J SWANSON BECK

A CUSCHIERI

Departments of Pathology and Surgery,
Ninewells Hospital and Medical School,
Dundee

¹ Pathology Working Group, Breast Cancer Task Force, *American Journal of Clinical Pathology*, 1973, 60, 789.

The bedpan round

SIR,—With Telstar in the sky the wasteful manner in which nurse manpower is deployed in the wards of Britain's nationalised hospitals is surely a blot on management. In an age when children are carrying goldfish swimming in water in polythene bags nurses in the wards of Britain's nationalised hospitals are being made to use metal bedpans weighing 4 lb each to transport a few ounces of bulk or liquid excreta in a manner which no employee of a firm in industry would tolerate.

This is not all. The unhygienic traditional manner in which bedpans are stored in between use in ward sluices is still in use. Because these bedpans after use cannot be guaranteed to have been made bacteria-free before being placed for use between the sheets of other patients it does not require much imagination to conclude that the procedure provides a channel whereby infection is transferred from one patient to another.

In no other situation would taxpayers be content to foot the bill for the increasing incidence of postoperative cross-infection

which is not only causing pain and distress to patients and their relatives but is responsible for the prolonged use of hospital beds, the extra time spent on care by medical men, nurses, and auxiliary staff, as well as the extra cost of hospital supplies.

In the year 1961, when I first had the idea of introducing a polypropylene bedpan with a detachable polythene liner which would take the place of the bedpan, I discussed the matter with the matrons of the main London hospitals. They all said they would never allow such a thing as bedside storage of bedpans. These inflexible attitudes have persisted and have led to all of the costly new hospitals being provided with ward sluices which are equipped with the old-fashioned metal bedpans weighing 4 lb, metal racks for storing them in between use, as well as bedpan washers and machines for macerating papier maché bedpan inserts, both of which use millions of gallons of water as well as thousands of pounds worth of electricity every day in the nationalised hospitals.

I wonder if your readers will think as I do that innovators have been under a delusion in thinking they would get any encouragement to invent new equipment and methods which could bring increased productivity, reduce unpleasantness, and obviate the danger of cross-infection for both nurses and patients.

E M GRANT

Albury, Surrey

Colour blindness and arithmetic

SIR,—I recently saw a case which I thought would be of interest to many of your readers. A bright young boy of 6 was found to be backward at arithmetic. The headmaster of the school explained to the parents that the system that they were using to teach the children was the Fletcher system, which uses colours, and he wondered whether the boy had a colour vision defect. On examination I confirmed that he was incompletely red/green blind; I also got a family history that he had an uncle with the same complaint.

Various authorities give different figures for the incidence of colour blindness, but I would say that the most generally accepted one is that 8% of males and 2% of females do have some form of inherited colour defect. If, therefore, a system like this is going to be used widely to teach arithmetic to youngsters then 5% of them are going to be backward through no fault of their own. I wrote to the specialist in community medicine drawing his attention to this point and had a most helpful letter in return. He is certainly taking action at a local level, but I thought that the issue raised was of such importance that it might be as well if more of my colleagues knew of this problem.

L H G MOORE

Dudley, W Midlands

Diet and coronary heart disease

SIR,—Professor A G Shaper and Miss Jean W Marr (2 April, p 867) have recommended a dietary change to postpone coronary heart disease in the British community. They recommend reduced intakes of foods high in saturated fats and replacement with foods relatively high in polyunsaturated fats. These

sound recommendations may be modified by recent events in the United States.

The US Senate Select Committee on Nutrition and Human Needs reported last February.¹ It had been taking evidence from doctors at public sessions since July 1976.² The committee has recommended many dietary changes. In order of priority these changes are: (1) increase consumption of starchy carbohydrates, called complex carbohydrates, from 22% of energy intake at present to 40-45% and increase consumption of whole grain products and also of fruit and vegetables, for the fibre from both is "important in nutrition"; (2) decrease sugar consumption from 24% to 15% of energy intake; (3) decrease consumption of total fats from 42% to 30%, saturated fats from 16% to 10%, and polyunsaturated and monounsaturated from 26% to 20%; (4) protein no change at 12%.

The evidence²⁻⁴ to support this enormous and difficult dietary change has been implemented by legislation on 31 March to create a new Bureau of Human Development.⁵ It has been funded by hundreds of millions of dollars. The committee is continuing to hold sessions and another report, this time on dietary fibre and health, has been promised.⁶

HUGH TROWELL

Woodgreen,
Fordingbridge, Hants

¹ US Senate Select Committee on Nutrition and Human Needs. *Dietary Goals for the United States*. US Government Printing Office, Washington, 1977.

² US Senate Select Committee on Nutrition and Human Needs. *Diet and Killer Diseases with Press Reaction and Additional Information*. US Government Printing Office, Washington, 1977.

³ *Hearings before the US Senate Select Committee on Nutrition and Human Needs. Part 1. Cardiovascular Disease*. US Government Printing Office, Washington, 1977.

⁴ *Hearings before the US Senate Select Committee on Nutrition and Human Needs. Part 2. Obesity*. US Government Printing Office, Washington, 1977.

⁵ A Bill to amend the Public Health Service Act to establish a Bureau of Human Development. *US Congressional Record*, vol 123, No 57, 31 March, 1977.

⁶ Senator George McGovern, personal communication, 1977.

Mycoplasma and encephalitis

SIR,—I was interested in the report of meningo-encephalitis associated with *Mycoplasma pneumoniae* infection (by Dr P Kopelman (2 April, p 881)) and the author's remarks concerning pathogenesis. It is relevant that sera of patients infected with *M. pneumoniae* show IgM antibodies that react with brain, since there are common glycolipid antigens.¹

E N WARDLE

Newcastle upon Tyne

¹ Kaplan, M H, and Rapaport, F T, *Proceedings of the First International Congress of Immunology*, p 1153. London, Academic Press, 1971.

Management of childhood epilepsy

SIR,—To manage means to control. As every schoolboy knows, drugs do not control childhood epilepsy, so one hoped that your leading article under the above title (16 April, p 986) would have been about something other than them. The control of the household, for instance, and of going to school, of climbing trees and swimming and of lighting fires and going camping; and of the prospect of marriage and pregnancy and rearing a family and

eugenics. All this was left out, and instead we had a little cliché about sympathetic advice.

That epileptics fall in the fire and char their limbs away; that by even lying on hot piping for an hour they can irretrievably damage them; that they drown in a bath three inches deep; that it is impossible, when swimming out of your depth, to support a child above water level for the duration of a major fit and that inhalation of river water is fatal: none of these was mentioned. No fire guards, no protection of staircases, no removal of bathroom locks, but only sadly ineffective, and sadly poisonous, drugs were discussed. It is as if your expert on epilepsy saw patients only in a hospital clinic, left the clinic only to go to the laboratory, and never saw the results in the home, in plastic surgery wards, and in the cemeteries, of such neglect of management.

H WYKEHAM BALME

London EC2

Methadone: evidence of accumulation

SIR,—Arising from the letter of Mr D M Rutherford and Dr K Raymond (19th March, p 774) we would firstly like to apologise for our inadvertent solecism in misspelling Mr Rutherford's name on two occasions in our previous letter (5 February, p 375).

We were most interested in their finding that the half life of methadone was 12-30 h. This would appear to bear out our comments on the variability which occurs in the published figures for this value.

It is of course well known that if a drug is administered repeatedly at constant time intervals it will accumulate until a "plateau level" is reached, when the concentrations fluctuate between a maximum and minimum level which can be calculated with precision by the method of Dost.¹ We would therefore agree with their introductory comments. However, we cannot accept their comments on the methadone blood levels which may originate from a misinterpretation of the graph published in our letter: If the whole-blood methadone concentrations of the three subjects are considered from the time of peaking over the longer 12-h period assessed on day 1 of the study the figures shown in the table are

Whole-blood methadone levels (ng/ml) after administration of one controlled-release tablet containing 10 mg of methadone

Time (hours)	Subject 1	Subject 2	Subject 3
1	—	—	58
2	54	89	34
3	48	60	27
6	44	55	25
8	41	50	23
12	32	33	18

obtained. Although the idealised flat blood concentration curves were not attained, as indeed they rarely are in practice, the results after peaking were very close to this theoretical concept. The release mechanism of the tablet was in fact designed to produce rapid analgesic activity through a measured loading dose and also to complement these initial levels by a further controlled release of methadone.

However, our main object in replying is not to argue the minutiae of the bioavailability pattern of our controlled-release methadone tablet but to highlight the problems of

accumulation occurring with methadone whether in a controlled, non-controlled, or liquid form. We found that the accumulation that occurred even during our short study gave rise to considerable side effects in the young, healthy volunteers. This accumulation, even if not indefinitely progressive, can thus give rise to problems even before a steady state is reached. Obviously this could be dangerous in elderly patients—as exemplified by Dr Paul Symonds's patient (19 February, p 512) who became unconscious with symptoms of narcotic overdose after the administration of 5 mg of methadone three times a day for two days. Such patients, often additionally exhausted by unrelieved pain or severe coughing, are vastly different from either healthy volunteers or young addicts, whom they also vastly exceed in number. In these elderly patients the risk of methadone accumulation and toxicity represents a substantial hazard.

We are aware that other physicians have encountered problems with methadone accumulation and may wish to comment in your columns.

S T LESLIE
R B MILLER
C BORODA

Research Division,
Napp Laboratories Ltd,
Watford, Herts

¹ Dost, F H, *Grundlagen der Pharmakokinetik*, Stuttgart, Georg Thieme, 1968.

Hospitable Trinidad

SIR,—Recently we three, all final-year medical students, spent elective periods attached to the University of the West Indies based in Port of Spain, Trinidad. We were all able to choose the departments in the hospital, health service, or research centre in which we wished to study. The many members of the medical services we met spent a great deal of time and effort teaching us and seeing to our general welfare. We are very grateful to them all. Our time in Trinidad proved an extremely valuable experience.

Arranging such elective studies is an obvious financial strain on any medical student. Our problems were greatly lessened by the fact that we were provided with rooms and meals at Port of Spain General Hospital free of charge. We would like to urge all British medical schools to extend similar hospitality (as we believe some already do) to University of the West Indies students and others on electives so that more medical students may have the chance of an experience such as ours. Our thanks to all in Britain and Trinidad who made our electives possible.

VALERIE A HARPIN

Middlesex Hospital Medical School

NIGEL ZOLTIE

Bristol University Medical School

BRIAN FERNANDES

Edinburgh University Medical School

Visual and manual examination of whole stools in plastic bags

SIR,—We should like to draw attention to a simple method for visual as well as tactile examination of faeces which we developed several years ago in order to detect remainders from oxytetracycline capsule medication. The