eye injuries was in players aged 11 to 15 years. Largely as a result of this study and others like it protective headwear and facewear is now mandatory in almost all junior ice hockey in Canada.

There have also been recent articles regarding eye injuries in badminton² and squash.³ I am not aware of such an article on cricket eye injuries. This is just the right time for such a survey. I write from the land of ice hockey to the land of cricket in the hope of stimulating somebody to tackle the sticky wicket.

PAUL M H CHERRY

Department of Ophthalmology,
University of Toronto,
Canada

¹ Pashby, T J, *et al*, *Canadian Medical Association Journal*, 1975, **113**, 663.

² Chandran, S, *British Journal of Ophthalmology*, 1974, **58**, 757.

³ Easterbrook, M, *Canadian Medical Association Journal*, 1978, **118**, 298.

Antipodean summer

SIR,—Dr Malcolm Schonell (18 March, p 711) states that the bite of the blue ringed octopus is always fatal as there is no known antidote. Certainly there is no known antidote, but the bite should never be fatal if there is a second person present. The toxin, related to tetrodotoxin, produces paralysis. The patient dies from anoxia due to his inability to breathe. If he is given artificial respiration he recovers. Dr Schonell will find his beach-walking hours less worrying if he purchases a copy of *Dangerous Marine Animals of the Indo-Pacific Region* by Dr Carl Edmonds, published by Wedneil Publications, Newport, Victoria, Australia, and available from any diving shop or good bookshop in Australia.

JOHN KNIGHT

Melbourne,
Australia

Bell's palsy and herpes simplex

SIR,—Last year Dr P Grout described in these columns a case of Bell's palsy associated with herpetic stomatitis (24 September 1977, p 829). Dr Grout's claim that the Bell's palsy was probably caused by herpes simplex virus was, however, subsequently questioned by Dr B E Juel-Jensen (1 October 1977, p 1086).

In the four months following the publication of these two letters four cases of Bell's palsy occurred in my practice and the opportunity was taken to determine the herpes simplex antibody titre in each case (see table). The rising antibody titres in cases 1 and 2 indicate current antigen stimulation, though the speed with which antibody was produced in case 1 suggests that the infection was not a primary one.

Case 1 has already been the subject of comment (12 November 1977, p 1292), for 12 months previously the patient had suffered

an attack of Bell's palsy on the opposite side. In case 2, though there was a significant rise in antibody titre to herpes simplex virus, the patient had, 10 days before the onset of the Bell's palsy, developed a typical herpeticiform rash around the auricle, spreading into the neck below the ear. The varicella-zoster antibody titre in both samples of blood was elevated to 256.

These results demonstrate a definite relationship between Bell's palsy and herpes simplex infection or reinfection in two of these cases and support the concept that Bell's palsy is a disease of multifactorial aetiology.

JOHN B WILSON

Lochmaben,
Lockerbie, Dumfriesshire

Relative activity of beta-blockers

SIR,—Dr J D Harry and Anne G Shields (8 July, p 128) report that in a healthy volunteer study atenolol is significantly more effective 24 hours after ingestion than metoprolol in reducing heart rate. Professor J H Barber (29 July, p 357) reports that in a double-blind general practice study he has shown metoprolol and atenolol equipotent three and 23 hours after ingestion in the treatment of hypertensive patients. He further comments that, in view of these results, metoprolol is preferable to grounds of comparative cost.

Professor Barber suggests that studies of real patients in general practice are of greater value than investigations in volunteers. Shaw,¹ writing on the use of once daily sotalol in general practice, comments that the drug gives effective 24-hour control of lying, standing, and postexercise blood pressures and pulse rates.

Lehtonen and Sundquist² have recently published their findings of a double-blind comparative study of sotalol and metoprolol in hypertensive patients, medication being administered once daily. Sotalol (240 mg) and metoprolol (200 mg) were shown to have similar hypotensive effects 24 hours after ingestion, but the effect of sotalol on pulse rate reduction was seen to be greater than that of metoprolol and lasted for at least 48 hours after a single dose. This suggests that sotalol may have a better cardioprotective effect than metoprolol when both drugs are given once daily.

There is a growing tendency towards using single daily doses of beta-blockers in the management of cardiovascular disease. In the light of these clinical findings, perhaps Professor Barber would now consider sotalol. The economic comparison is also favourable.

B M GUYER
Medical director

Bristol-Myers Co Ltd,
Langley,
Slough

¹ Shaw, H L, *Journal of the Royal College of General Practitioners*, 1977, **27**, 742.

² Lehtonen, A, and Sundquist, H, *Current Therapeutic Research*, 1978, **23**, 131.

Drugs acting on the bladder and urethra

SIR,—In the recent article by Mr Stuart L Stanton entitled "Drugs acting on the bladder and urethra" (17 June, p 1607) a number of statements were made which we found surprising.

(1) It is difficult to equate urethral sphincter dysfunction with genuine stress incontinence, since there are many cases of sphincter overactivity which do not result in "stress." The author is aware of this, as he recommends the use of baclofen in external sphincter spasm later in the text.

(2) It was pointed out that since the prostaglandins (PG) stimulate bladder muscle in vitro they might be of some consequence in the treatment of "decreased bladder activity." The author added, however, that he had been unable to initiate bladder contractions after intravesical administration of PGE₂. By contrast our studies¹ have shown that PGE₂ administered in this way will initiate bladder contractions in atonic bladders and we now use PGE₂ routinely to treat idiopathic chronic retention in women and also in men who remain in chronic retention after relief of the causative outflow obstruction.

(3) The author rightly suggested the use of carbachol or bethanechol for treatment of the atonic bladder but went on to say that its action is due to an ability to release acetylcholine from parasympathetic nerve terminals. Surely a direct action at muscarinic receptors on the smooth muscle is the major effect, with increased transmitter release being only a contributory factor.

(4) In the section on side effects attention is given to diphenylhydantoin in that it may aggravate incontinence because it is an α -blocking agent. We do not know any evidence to support this suggested mode of action.

N G BOWERY

Department of Pharmacology,
St Thomas's Hospital Medical School,
London

N H HILLS

Frimley Park Hospital and
St Peter's Hospital,
Chertsey

¹ Bultitude, M I, Hills, N H, and Shuttleworth, K E D, *British Journal of Urology*, 1976, **48**, 631.

*. *We sent a copy of this letter to Mr Stanton, whose reply is printed below.—Ed, BMJ.

SIR,—I should like to make these points in reply to the letter of Dr Bowery and Mr Hills.

(1) Urethral sphincter dysfunction is a term I have used for several years to denote incompetence or decreased activity of the urethral sphincter mechanism. It is a parallel term to genuine stress incontinence. I never use it to describe sphincter overactivity or spasm, which is very different and which does not result in "stress incontinence."

(2) I think Bowery and Hills's experience with PGE₂ is unique and would welcome evidence of its efficacy from other centres.

(3) I would not dispute that the major direct action of carbachol and bethanechol is at muscarinic receptors on smooth muscle. In endeavouring to be succinct, I have perhaps oversimplified the property of choline esters by describing their action as one of stimulation of the parasympathetic nervous system to release acetylcholine.

(4) I would refer Bowery and Hills to a

Cases of Bell's palsy and reciprocal herpes simplex antibody titres

Case	Age	Sex	Date of onset	First specimen		Second specimen	
				Date	Titre	Date	Titre
1	62	F	11.9.77	14.9.77	32	3.10.77	256
2	92	F	16.9.77	26.9.77	32	11.10.77	256
3	48	M	7.1.78	7.1.78	128	15.2.78	64
4	25	F	1.2.78	1.2.78	128	14.2.78	64