CORRESPONDENCE

Syringe-transmitted hepatitis A J Zuckerman, FRCPATH..... 696	Self-poisoning by tobacco L M Johnston, MB..... 700	Drugs acting on the bladder and urethra N G Bowery, PHD, and N H Hills, FRCS; S L R Stanton, FRCS..... 704
Undergraduate and postgraduate clinical training J P Quilliam, FRCP..... 697	Health of King Henry VIII M Skoblo, MB, and J Leslau..... 700	Epilepsy in coeliac disease T H Hughes-Davies, FRCP..... 705
Distribution of registrars L P Le Quesne, FRCS..... 697	Haemodialysis-triggered asthma R H Jones, MRCP, and others..... 701	Length of hospital stay M G Jacoby, MB, and D T Greaves, MD.... 705
Survey of notifications of tuberculosis in England and Wales K M Citron, FRCP, and W Fox, FRCP..... 697	Sexual pressures on children J S Bradshaw, MB; Mary Kingston; Elizabeth M Elliott, MB; J Phelan..... 701	Treatment of ulcerative colitis R J Dickinson, MRCP, and others..... 705
Efficacy of rubella vaccination D S Freestone, MB..... 698	Controlled prescribing C R Howlett, MRCP..... 702	Infant feeding practices P D Manuel, MRCP, and J A Walker-Smith, FRCP..... 705
Rubella susceptibility in women M Gringras, MB, and C H Bowker, MB.... 698	Proprietary names of drugs Dorothy C Moir, MD..... 703	Audiometry in industry D W Robinson, DSC, and W Burns, FRCP.. 706
Rubella immunisation J Frampton, FRCOG..... 698	Prevalence of multiple sclerosis in north-east Scotland A J Silman, MRCP..... 703	Amphi- F W Hanford, MRCS..... 706
Rubella embryopathy R Pilsworth, MD, and Barbara J Pilsworth, FIMLT..... 698	Lorazepam unsuitable for day surgery P J Simpson, FFARCS..... 703	Maltese medical students J C Cameron, FRCP..... 706
Relative chloroquine resistance of Plasmodium falciparum in Zambia H De Glanville, MB; M J Maguire, MRCP.. 698	Health education in schools Rosalind Cole..... 703	Radiology work load M Lea Thomas, FRCP; B Eyes, MB, and A F Evans, FRCP..... 706
Whole gut irrigation K D Vellacott, FRCS, and T W Balfour, FRCS; J Hurdley, FFARCS..... 699	PUVA treatment of psoriasis E J Land, PHD..... 703	Money for old rope W Sagar, MB..... 707
If I had... a sigmoidoscopy Major-General W H Hargreaves, FRCP.... 699	Stroke after acute myocardial infarction N P Quinn, MRCP..... 703	X-ray services for GPs J D Wiggdahl, MRCP..... 707
Psychiatric ward rounds H D Beckett, MRCPsych..... 699	Injuries to cricketers P M H Cherry, FRCS..... 703	Farical overtime rates A R Rogers, MB..... 707
Breathing other people's smoke S L Stock, BSC; R A McCartney; M C S Kennedy, MRCS..... 699	Antipodean summer J Knight, FFARCS..... 704	"Caithness and Sunderland Division" W H Gossip, MB, CHB..... 707
	Bell's palsy and herpes simplex J B Wilson, FRCPED..... 704	New consultant contract D E B Powell, MD..... 707
	Relative activity of beta-blockers B M Guyer, MB..... 704	Correction: Six months in South Africa Szanto..... 707

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters must be signed personally by all their authors.

Syringe-transmitted hepatitis

SIR,—A recent inquiry from an area health authority addressed to me concerned the risk of hepatitis B and the current practice of performing mass BCG vaccination using the same syringe but with a different hypodermic needle for each person. In my view this method carries a high risk of transmitting viral hepatitis type B¹ and perhaps also the more recently described third form of hepatitis (so called non-A: non-B hepatitis).

The evidence is indisputable. Over 30 years ago hepatitis among patients attending a diabetic clinic was traced to a common syringe which was used for withdrawing blood for blood sugar estimations.² Over two years 63 cases of hepatitis with four deaths occurred in patients attending this clinic. It was considered that infection resulted from contamination of the nozzle of the syringe with blood. It was subsequently shown that removal of a needle from a syringe after completion of an injection can draw blood or tissue fluid in the lumen of the needle back into the syringe nozzle.^{3,4} Other laboratory studies⁵⁻⁷ also confirmed this explanation of

outbreaks of hepatitis where the multidose method of syringe loading, but with a change of needle after each injection, was practised. Epidemiological evidence for the transmission of hepatitis B by this method is not lacking. In 1948 23 cases of hepatitis occurred in 110 soldiers inoculated with tetanus toxoid by the multidose-syringe technique.⁸ In another study significantly more cases of jaundice than would be expected were found among Royal Navy personnel who had been inoculated with typhoid-paratyphoid vaccine or tetanus toxoid, or both.⁹ In a more recent retrospective and prospective study of hepatitis in the Royal Air Force it was concluded that about 11% of 895 patients with acute hepatitis contracted the infection through syringe transmission.¹⁰ The multidose-syringe technique was universally practised at the time in the Royal Air Force.

Survival of hepatitis B virus is ensured by the reservoir of persistent carriers of this virus in the world, estimated to number between 120 and 175 million. The prevalence of carriers, particularly among blood donors, in North America and in northern Europe is

0.1% or less and in central and eastern Europe up to 5%; the frequency is higher in southern Europe and the countries bordering the Mediterranean, and in some parts of Africa and Asia as many as 20% of the population may be carriers.¹¹ The risk of transmitting hepatitis by the multidose-syringe technique is therefore considerable and it is imperative that an adequately sterilised or a disposable syringe and needle should be used for each individual patient.

A J ZUCKERMAN

WHO Collaborating Centre for Reference and Research on Viral Hepatitis, London School of Hygiene and Tropical Medicine, London

¹ Zuckerman, A J, *Virus Diseases of the Liver*, p 23-41. London, Butterworths, 1970.

² Droller, H, *British Medical Journal*, 1945, 1, 623.

³ Hughes, R R, *Journal of the Royal Army Medical Corps*, 1946, 87, 156.

⁴ Evans, R J, and Spooner, E T C, *British Medical Journal*, 1950, 2, 185.

⁵ Fleming, A, and Ogilvie, A C, *British Medical Journal*, 1951, 1, 543.

⁶ Ruys, A C, and Prakken, J, *Nederland Tijdschrift Geneesk*, 1955, 99, 1163.

⁷ Kranerdonk, O, Borst, J, and Ruys, A C, *Nederland Tijdschrift Geneesk*, 1960, 1-4, 1576.

⁸ Capps, R B, Sborov, V, and Scheiffley, C S, *Journal of the American Medical Association*, 1948, 136, 819.

⁹ Ellis, F P, *Journal of Hygiene, Cambridge*, 1953, 51, 145.

¹⁰ Zuckerman, A J, *British Journal of Social and Preventive Medicine*, 1964, 18, 183.

¹¹ Zuckerman, A J, *Bulletin of the World Health Organisation*, 1978, 58, 1.