It should perhaps be pointed out that normal melatonin levels during the night (2400) cover a very large range,² which in my own laboratory currently extends from undetectable (<7) to 161 pg/ml with one value of 269 pg/ml at 0200. Great care is therefore needed in describing a night-time plasma melatonin level as "abnormal." Inter-individual daytime levels are admittedly more stable (mean $\pm$ SEM at 1200: 13.9 $\pm$ 0.5 pg/ml, $n=50$, men; 14.6 $\pm$ 0.9 pg/ml, $n=45$, women) and the main interest of the case reported by Dr Barber and his colleagues lies in the very high daytime levels.

These data were largely collected at the University Paediatric Clinic, Geneva

JOSEPHINE ARENDT

Department of Biochemistry,
University of Surrey,
Guildford

¹ Kennaway, D J, Thesis, University of Adelaide, Australia, 1978.

² Arendt, J, *Journal of Neural Transmission*. In press.

SIR,—I read with interest the article by Dr S G Barber and his colleagues (29 July, p 328) but would like to introduce a word of caution in the interpretation of serum melatonin levels in patients with pineal tumours. Last year we reported a raised level of melatonin in the serum of a 30-year-old man who was found on biopsy to have a germinoma of the pineal region.¹ The calcified pineal gland was seen radiologically adjacent to the tumour. The preoperative level (at 10 am) of 185 pg/ml fell to 19 pg/ml four weeks after treatment by surgery and radiotherapy. Since that time melatonin has not been detectable in the serum. One year after treatment the patient has a thalamic syndrome but there is no evidence of recurrence of the tumour. It is clear, therefore, that patients with non-parenchymatous tumours of the pineal region may have raised levels of melatonin in the serum, and consequently it must not be assumed that raised serum melatonin levels necessarily indicate a melatonin-secreting tumour (pineocytoma) of the pineal gland.

In our case it is just possible that the raised levels of melatonin were the result of secretion from differentiated pineal tissue within the germ-cell tumour. It is far more likely, however, that they were the result of interference with the mechanisms controlling melatonin secretion by the pineal gland. These are, of course, incompletely understood, but it is clear that melatonin secretion in experimental animals is mediated by the autonomic nervous system and in particular by postganglionic sympathetic nerve fibres originating in the superior cervical ganglion. These in turn are influenced predominantly by the effect of light on the retina, the pathway to the superior cervical ganglion involving the inferior accessory optic tracts. Under these circumstances it is difficult to understand how Dr Barber and his colleagues explain how a tumour which presumably does not have any autonomic nervous supply can be "under at least some degree of rhythmic control." It would be most interesting to see the variations in the serum melatonin levels over a rather longer period than the 24 hours given in their article.

In our case there was a complete disappearance of melatonin from the serum at three months after treatment. If the unlikely possibility of there being pineal tissue in the germ-cell tumour is discounted, then it must

be assumed that this was due to complete ablation of the pineal gland by the treatment. It is clear, however, that serum melatonin levels in patients with secreting pineal tumours may well be useful in determining the effect of treatment and, of course, in detecting recurrence of such tumours.

Finally, with regard to the effect of the pineal gland on tumours elsewhere in the body it is worth noting that melatonin does not necessarily suppress tumours and may indeed have quite the opposite effect, the growth and malignancy of dimethylbenzanthracene-induced mammary tumours in rats being potentiated by the administration of melatonin.²

Thanks are due to Mr R A Daws for the clinical information and Dr P G Lynch for the histological report.

E TAPP

Group Laboratory,
Royal Infirmary,
Preston

¹ Tapp, E, Skinner, R G, and Phillips, V, *Proceedings of Symposium, Vienna, 1978*. Basle, Karger. In press.

² Hamilton, T, *British Journal of Surgery*, 1969, 56, 764.

Penicillamine and previous treatment with gold

SIR,—As a scientist with an interest both in statistics and in the treatment of rheumatoid arthritis I was fascinated to read the report by Drs M Webley and E N Coomes on the possible link between penicillamine side effects and previous gold therapy (8 July, p 91). The rationale for their survey seems excellent, but I was surprised and dismayed at the unsatisfactory treatment of their results.

Their statement that "rashes and marrow depression were twice as common in the [previously] gold-treated group" is incontrovertible. However, the relevance of the observation cannot be judged without knowledge of the associated statistical probabilities. Recourse to the simple and widely used χ^2 test would have easily provided this valuable piece of information. Calculation of the probabilities that these were chance observations shows them to be greater than 10% for rashes and almost 10% for marrow depression. In other words the differences are not statistically significant.

On this basis I would respectfully suggest amendment of the first sentence of the final paragraph to read, "Our results show that rashes, and possibly marrow depression, may be more common if patients have received gold. However, our patient sample was too small to allow the hypothesis to be accepted at the required level of probability."

I fear that this report confirms the worry that medical statistics still have a long way to go before their necessity is universally and fully accepted by the writers, publishers, and readers of original medical articles.

D A GOUGH

Caversham,
Reading, Berks

* * * We sent a copy of this letter to Drs Webley and Coomes, whose reply is printed below.—
Ed, *BMJ*

SIR,—In our short report we merely recorded our experiences and tabulated our findings. Like Mr Gough we know about the χ^2 test and agree that our results were not statistically

significant and never said they were. However, we feel we have made an interesting point which needs looking into further.

In fact, we have done the sums ourselves and got the same results as him, but can't one make an observation these days without statistics?

E N COOMES
MICHAEL WEBLEY

St Stephen's Hospital,
London SW10

SIR,—We would like to comment on the report by Drs M Webley and E N Coomes (8 July, p 91) which provides evidence that penicillamine-induced rashes and marrow depression are more common in patients with rheumatoid arthritis who have previously received chrysotherapy. This has been our impression and that of others.¹ They speculate that some of their early rashes might have been due to release of tissue gold. Penicillamine, which can certainly mobilise gold in man with the formation of a stable chelate,² has in fact been used to treat gold dermatitis.³

An alternative explanation of the increased prevalence of side effects might be the presence of underlying Sjögren's syndrome. In a random series of 32 patients with rheumatoid arthritis⁴ nine of 11 with keratoconjunctivitis, an important feature of Sjögren's syndrome, developed a drug reaction. Six of these were rashes due to gold, while two patients developed a rash on penicillamine. One of these two had previously experienced a gold rash and the other persistent proteinuria, again during chrysotherapy. The remaining three patients were on other drugs. By comparison there were only four with drug reactions out of 21 patients without keratoconjunctivitis. This difference is statistically significant ($\chi^2=9.33$, $P<0.005$). Other workers⁵ have also found drug allergy to be more common in patients with Sjögren's syndrome.

L FERNANDES
ERIC HAMILTON

Department of Rheumatology and
Rehabilitation,
King's College Hospital,
London SE5

IVAN WILLIAMS

Department of Rheumatology and
Rehabilitation,
Kent and Sussex Hospital,
Tunbridge Wells, Kent

¹ Day, A T, and Golding, J R, *Postgraduate Medical Journal*, 1974, 50, suppl 2, p 71.

² Eyring, E J, and Engleman, E P, *Arthritis and Rheumatism*, 1963, 6, 216.

³ Davis, C M, *American Journal of Medicine*, 1969, 46, 472.

⁴ Fernandes, L, and Hamilton, E B D, unpublished observations.

⁵ Whaley, K, et al, *Quarterly Journal of Medicine*, 1973, 42, 513.

Thyroid extract

SIR,—I was surprised that Dr W van't Hoff and his colleagues (15 July, p 200) felt so strongly about thyroid extract. They consider it to be "dangerous." If these authors now subscribe to my view that inadequate thyroid replacement leads to coronary heart disease (CHD) it might theoretically be argued that the possible varying potency of thyroid extract could be dangerous. However, it is the adequate replacement dose rather than the preparation used which is crucial in preventing CHD.

If the patient is symptom free and you do not subscribe to my view that inadequate replacement therapy leads to CHD, thyroid

extract can in no way be considered dangerous. Many wise doctors will safely continue to give thyroid extract to patients who have been well on it for years. Newly diagnosed myxoedema should be treated with thyroxine (or very rarely triiodothyronine), but elderly women on thyroid extract are not living dangerously.

P B S FOWLER

Charing Cross Hospital,
London W6

Beta-blockers in treatment of hypertension

SIR,—Dr R G Wilcox's recent comparative study of once-daily hypotensive treatment (5 August, p 383) showed that atenolol 200 mg produced significantly greater effect than the other agents in the doses tested. He rightly points out that the doses of beta-blocking drugs may not have been equipotent, illustrating a potential limitation of fixed-dose comparative studies. Little further hypotensive effect can be expected by increasing the dose of atenolol further,¹ but this is not necessarily true of the other beta-blocking drugs, and he himself suggests that higher doses of acebutolol or timolol might be more effective.

I am able to confirm these speculations in the case of acebutolol from a recent study.² In the twice-daily dose ranging part of this study ($n=11$), where the dose of acebutolol was increased above the 400 mg/day used by Dr Wilcox, further falls in blood pressure were produced. This fall in blood pressure was unchanged when each subject's total daily dose was given once daily and blood pressure measured more than 24 h after dosing. Furthermore, there was no increase in the low incidence of side effects. In addition, I found that acebutolol 400 mg produced less beta-blockade, measured as reduction in exercise tachycardia, than atenolol 200 mg (maximum $23 \pm 4\%$ v $29 \pm 3\%$, $P < 0.05$, $n=8$), confirming a lack of equipotency in beta-blockade in Dr Wilcox's study.

If useful short-term differences are to be shown between different beta-blockers in the treatment of hypertension they will relate to the magnitude of maximum effect and the side effects and compliance present at any specified level of effect with different agents. Consequently full assessment of the comparative efficacy of beta-blockers in hypertension will also require optimum-dose studies of these agents.

MICHAEL MARTIN

Department of Therapeutics,
Hallamshire Hospital,
Sheffield

¹ Zacharias, F J, *et al*, *Postgraduate Medical Journal*, 1977, **53**, suppl, p 114.

² Martin, M A, *et al*, *British Journal of Clinical Pharmacology*. In press.

SIR,—We were interested in the paper by Dr R G Wilcox (5 August, p 383), as too few studies measure exercise blood pressure during the 24-h period following drug administration. It is surely misleading, though, for the author to discuss changes in exercise blood pressure levels when the only reading taken was immediately on cessation of exercise, this presumably being at some point between 10 and 45 s of the post-exercise phase. Of more interest would be the blood pressure recorded during, say, the last 30 s of exercise. Absolute levels, in both the untreated and treated

hypertensive patient, fall sharply by as much as 40–50 mm Hg systolic within the first 10–15 s of stopping exercise¹ and therefore it appears inappropriate to use a Hawksley random zero sphygmomanometer in a very rapidly changing haemodynamic situation. Furthermore, a 1-min dynamic exercise is useless in these studies as it takes at least 3 min to achieve steady-state levels of raised blood pressure.

As none of the drugs used in the study totally prevented significant increases in the immediate post-exercise systolic blood pressure it is reasonable to assume that the same applies to the blood pressure levels occurring towards the end of exercise and would suggest that the dose used was inadequate. From work we are currently evaluating it would appear that exercise-induced "peaks" of systolic blood pressure in certain hypertensive patients of 200 mm Hg or more can be attenuated in the whole of the 24-h period by adequate doses of some beta-blocking drugs following a single daily dose, the systolic pressure being reduced to the more normal range of 140–160 mm Hg. That the "peaks" carry important risks still remains to be proved, but in the meanwhile Sokolow's findings² may well be pertinent.

Finally, we feel that to demonstrate the hypotensive effect only at rest and not during exercise is an observation of limited value when most hypertensive patients are mobile during much of the 24-h period. The correct therapeutic dose is important and can only be judged by effective control of the stress peaks.

M BRIAN COMERFORD
EDWIN BESTERMAN

Waller Cardiopulmonary Unit,
St Mary's Hospital,
London W2

¹ Millar-Craig, M, and Raftery, E B, personal communication.

² Sokolow, M, *et al*, *Circulation*, 1966, **34**, 279.

Computer confidentiality

SIR,—It is profoundly to be hoped that the medical profession as a whole does not conclude from the expressed intentions of the Department of Health and Social Security or from the views of its supporters in this matter (Drs P M Dunn and C H M Walker (22 July, p 276) and Dr P Samet (5 August, p 429)) that workers in the field of computing are collectively bent on undermining the concept of confidentiality of patients' records. To one who has worked in co-operation with doctors for any length of time it should be apparent that adequate physical security of computer files, even if it can be guaranteed (Dr Samet), is only a first prerequisite for overall effectiveness of a system which is bound to involve the human element at many levels. It should also be evident that it is equally important that a belief in this security be transmitted to the patients (that is, the general public) or their trust in their medical advisers will be lost, with all the unfortunate consequences that that would entail. I contend that at a time when stories of computers losing driving licences and producing inflated gas bills are rife, and when there is talk of totalitarian regimes using computerised records for their own ends, public faith in the integrity of the proposed scheme would with some justification be low, and that the medical professions are right in their present attitude of opposition (22 July, p 298; Dr R M Forrester, 5 August, p 429).

The situation may very well change, perhaps sooner than we realise; but it is imperative that even when the time comes when medical records can confidently be committed to computer storage on a nationwide scale each individual patient record must remain localised and under the control of the doctor or doctors responsible for that patient.

K A JOHNS

Department of Medical Statistics,
Welsh National School of Medicine,
Cardiff

Cigarette smoking and postoperative deep-vein thrombosis

SIR,—Mr J K Clayton and his colleagues (5 August, p 402) confirm a lower incidence of postoperative isotopic deep-vein thrombosis (DVT) in cigarette smokers and suggest that this may be related to body weight or may "reflect resistance to venous thrombosis as a constitutional feature which correlates with but is not due to a predilection to smoke cigarettes."

In a recent series of 166 patients undergoing emergency or elective laparotomy for benign or malignant disease or retropubic prostatectomy in whom two methods of intermittent pneumatic compression of the calves were compared we found that the total incidence of isotopically detected DVT was 31.3%. Ten out of 52 (19.2%) cigarette smokers, seven out of 17 (41.2%) pipe smokers, and 35 out of 97 (36.1%) non-smokers developed DVT. Cigarette smokers had a significantly lower incidence than the rest ($\chi^2=5.15$, $P < 0.05$).

The principal difference between non-smokers and cigarette smokers turned out to be their age, which was 68.2 ± 1.0 (SEM) years among non-smokers and 61.0 ± 1.4 among cigarette smokers. This difference is highly significant ($t=4.12$, $P < 0.001$). The mean age of pipe smokers was 66.5 years, which did not differ significantly from that of non-smokers.

Analysis of the patients' weights as percentages of those expected (*Geigy Tables*) showed such a large variance (from -31% to $+237\%$) that we could not use a parametric test. Rather we arranged the weights into five groups: $<0\%$, $0-4\%$, $5-9\%$, $10-14\%$, and $15\%+$. The rates of DVT in these groups (cigarette smokers first, non-smokers together with pipe smokers second) were: 20% and 30%, 20% and 53%, 25% and 35%, 0% and 30%, and 22% and 43%.

Although it may be true that cigarette smokers are younger and thinner than non-smokers, that is certainly not the whole explanation of the protective effect of this otherwise undesirable habit.

A V POLLOCK
MARY EVANS

Scarborough Hospital,
Scarborough, N Yorks

Investigating stroke

SIR,—Your leading article (10 June, p 1503) highlights the dilemma clinicians face now that accurate, non-invasive technical equipment with which to investigate stroke patients has become available, at least in some areas. As you say, this applies particularly to at-risk groups—diabetics, hypertensives, and the elderly in general.

While working in the department of