loss of mobility he has always displayed the euphoria which as a student I learnt was characteristic of disseminated sclerosis. On 14 August I dropped in to see him and was struck by the fact that he was depressed and quite unlike his usual self. I do not have television in my house and knew nothing of the programme. It was only when I was leaving the house that his wife told me that this change came on after he had seen the programme on the Friday night.

The programme was apparently called "Pioneers of Hope" but I gather it was what he learnt about possible bladder complications which destroyed any hope he might have had. Is it really a good thing for the general public to see and hear a layman putting questions to doctors who do not know the certain answers and have different opinions about possible theoretical solutions?

R N RUTHERFURD

Oakley, Kirkcudbright

Carcinoma of the breast in women under the age of 30

SIR,—We note with concern the comment in your leading article on "Breast lumps in adolescent girls" (4 February, p 260) that cancer of the breast under the age of 30 is extremely rare. Haagensen¹ collected 6000 cases over the period 1916-66 at the Columbia Presbyterian Medical Centre, New York, and found the incidence in his series to be 1.2% for the women under the age of 30 years.

This series should not be regarded as representing the present-day incidence, as it does not take into account the recent increase in carcinoma of the breast and changes in Western society. Our experience is rather alarming and we wish to alert readers to the fact that cancer of the breast under the age of 30 is not rare.

From July 1976 to June 1978 510 patients (6 men) had their breast lesions excised in Doncaster Royal Infirmary, a 804-bedded hospital serving a working-class population of 280 000 in south Yorkshire. There were 197 carcinomas (two of them in men), and 10 (5.1%) occurred in women aged under 30. The age distribution in women is shown in the table below.

Histological examination of the carcinoma in these 10 women showed all to be poorly differentiated carcinomas. In one case axillary lymph nodes, the liver, and the opposite breast contained tumour, and in another the axillary lymph nodes, the lungs, and the vertebrae contained tumour. The latter patient died within nine months from the beginning of symptoms while the former is now jaundiced and in a terminal condition. Nine of the 10 women were married and six already had children. In four cases the lump developed within a year after delivery and in three of these four cases lactation stopped within a few weeks after delivery and the lump became apparent.

Thus cancer of the breast below the age of 30 does appear more often than was thought, and we consider doctors should be aware of this fact. We will be pleased to hear about the experiences of other centres in this country.

A S PURANDARE
J FINBOW

Doncaster Royal Infirmary,
South Yorkshire

¹ Haagensen, C D, in *Diseases of the Breast*, p 352. Philadelphia, Saunders, 1971.

Genital chlamydial infection

SIR,—I would like to comment on the subject of chlamydial genital infections in women, raised by Drs Shirley J Richmond and J D Oriel (12 August, p 480). In my current survey of gynaecology inpatients with their first reported attack of acute pelvic inflammatory disease results so far show only three positive chlamydial cervical cultures from the 27 patients with valid specimens. The diagnostic criteria are purely clinical, similar to those proposed elsewhere.¹

Using very strict laparoscopic diagnostic criteria, Mardh² collected positive chlamydial cultures in over 30% of cases. The report³ of 9% of a series of gynaecology outpatients with positive chlamydial cultures is not directly comparable because of the wide range of conditions seen in such clinics. Reports from studies performed on patients attending clinics for sexually transmitted diseases show that rates of positive chlamydial cultures vary depending on the exact status of the groups investigated⁴⁻⁶ and also on the clinical cervical signs.^{7,8}

Selection of specific patients for chlamydial culture is likely to continue for a variety of reasons, but further work continues here on the identification of certain groups of patients whose management would benefit from a chlamydial culture service, as Dr Richmond suggested. The measurement of antichlamydial antibodies in serum or genital secretions, or both, may also assist in this identification in the future.

B H O'CONNOR

MRC Sexually Transmitted
Diseases Unit,
Middlesex Hospital Medical School,
London W1

¹ Wright, N H, and Laemmle, P, *American Journal of Obstetrics and Gynecology*, 1968, **101**, 979.

² Mardh, P, et al, *New England Journal of Medicine*, 1977, **296**, 1377.

³ Paavonen, J, et al, *British Journal of Venereal Diseases*, 1978, **54**, 257.

⁴ Dunlop, E M C, et al, *British Journal of Venereal Diseases*, 1972, **48**, 425.

⁵ Oriel, J D, et al, *British Journal of Venereal Diseases*, 1972, **48**, 429.

⁶ Oriel, J D, *British Journal of Venereal Diseases*, 1974, **50**, 11.

⁷ Rees, E, et al, in *Non-Gonococcal Urethritis and Related Infections*, ed D Hobson and K K Holmes. Washington, DC, American Society for Microbiology, 1977.

⁸ Oriel, J D, et al, *Journal of Infectious Diseases*, 1978, **137**, 443.

Thyroid function tests

SIR,—Dr P Sheridan and others (12 August, p 477) suggest the use of a "modified" free thyroxine index (FTI) to allow for a "zero error" of the Thyopac 3 assay of thyroid hormone binding capacity. They suggest that such a modified FTI may be a better discriminant than T4:thyroxine binding globulin (TBG) ratio^{1,2} for evaluating thyroid function in patients with abnormal binding proteins. Although Sheridan *et al* state that we did not study patients with low TBG concentration, we did in fact include data on five patients with absent TBG² and showed that in such euthyroid patients a serum T4 concentration of 20 nmol/l (1.6 µg/100 ml) represents the hormone bound to thyroxine binding pre-albumin and albumin. Sheridan *et al* used data from these patients to calculate their modified FTI. Before accepting that a modified FTI is a useful index for diagnosing thyroid disease, we should need to see more evidence of the ability of such an index to identify patients with thyroid disease.

Since Sheridan *et al* have used data from our patients with abnormal TBG concentrations to calculate a modified FTI, we feel justified in applying the modified FTI to our dysthyroid patients, comprising 101 with thyrotoxicosis and 36 with myxoedema (in borderline cases the diagnosis was confirmed by measurements of TSH, T3, and dynamic tests of thyroid function as well as clinical response). Using the confidence limits of Sheridan *et al*, eight patients with myxoedema were classified as euthyroid by the modified FTI, as were three with thyrotoxicosis. Using conventional FTI, four patients with myxoedema were incorrectly diagnosed as euthyroid while all patients with thyrotoxicosis were correctly diagnosed. T4:TBG ratio classified two patients with myxoedema as euthyroid, and all those with thyrotoxicosis were correctly diagnosed. We suggest that the modified FTI is probably inferior to conventional FTI in thyroid diagnosis and almost certainly inferior to the T4:TBG ratio.

We think that most clinicians would prefer to be provided with a direct measurement of thyroxine and its major binding protein, rather than a figure derived from mathematical manipulation of an empirical test. Since TBG measurement is less costly than commercial T3 uptake kits, it would seem logical to abolish the T3 uptake and replace it with direct measurement of TBG.

W A BURR

University Department of
Medicine,
Queen Elizabeth Hospital,
Birmingham

¹ Harvey, R F, *Lancet*, 1971, **2**, 230.

² Burr, W A, et al, *British Medical Journal*, 1977, **1**, 485.

SIR,—The report by Dr P Sheridan and others (12 August, p 477) suggesting a new equation for calculating free thyroxine index (FTI) is interesting but not, we feel, very helpful. The FTI, calculated using the Thyopac-3 test, may be misleading at both high and low levels of thyroid-binding protein, but since this is for different reasons separate modifications are required for high and low Thyopac-3 results.

We consider that the problem of the misleading FTI is associated with technical factors in the Thyopac-3 test, and modification of the method of assessing levels of thyroid

Age distribution of women with breast lesions

Age (years):	11-20	30	40	50	60	70	80	>80	Total
Benign	23	76	85	94	21	7	3	0	309
Malignant	0	10	16	45	43	42	31	8	195
Total	23	86	101	139	64	49	34	8	504