CORRESPONDENCE

Drug monitoring and adverse reactions J G R Howie, MD..... 1467	Vitamin D "resistance" and bioavailability of calciferol tablets A M Parfitt, MD..... 1470	Coronary artery spasm M J Butler, FRCS..... 1472
Diet and coronary heart disease Sir John McMichael, FRCP, FRCS..... 1467	Insurance companies' attitude to psychiatric illness D T Maclay, FRCPsych..... 1471	Effect of antihypertensive drugs on growth hormone secretion I Lancranjan, MD, and P Marbach, PhD... 1472
MRC treatment trial for mild hypertension W S Peart, FRCP, and W E Miall, MD.... 1468	Heparin and pulmonary embolism Marian Roden, MB..... 1471	Benign mucous-membrane pemphigoid associated with penicillamine treatment J S Pegum, FRCP, and A C Pembroke, MRCP 1473
Breath, alcohol, and the law J P Payne, FFARCS, and D W Hill, PhD.... 1468	Misdiagnosis of urinary tract infection R Hole, FRCS..... 1471	Correcting the calcium P J Phillips, MRACP, and R W Pain, FRCPA 1473
Malaria in returning travellers R G Thompson, MRCPATH, and others.... 1468	Rise in antibodies to human papova virus BK and clinical disease J v d Noordaa, MD, and P Wertheim-van Dillen, MD..... 1471	Medical hazards of air travel J G Callanan, FRCS..... 1473
Making better use of our nurses B L E C Reedy, MRCP..... 1468	Further assessment of the normal cholecystogram Margaret R Jones, FRCP, and D J T Webster, FRCS..... 1471	Confidentiality and the three wise men J B Harman, FRCP..... 1474
Intrauterine fetal transfusion J M Bowman, MD..... 1469	Management of childhood epilepsy D C Thrush, MD..... 1472	COHSE and the Royal Commission V B Whittaker, MRCPED..... 1474
Plasma exchange in severe rhesus disease E P J McGuinness, MRCP, and D J Reen, PhD..... 1469	Acute renal failure associated with carbenoxalone treatment B P Hurley, FRACP..... 1472	Related ancillary staff M E Glanvill, MRCP..... 1474
Ulcerogenic action of azapropazone R J Ancill, FRCPATH..... 1469	Transient hypotension following intravenous ethamsylate (Dicynene) L Langdon, FFARCS..... 1472	Professional indemnity M J Illingworth, MB..... 1474
Interferons and rabies Colonel E E Vella, FRCPATH..... 1470		Points from letters Charges to private patients for use of NHS facilities (J H E Bergin); Pelvic sepsis after hysterectomy (G S W Evans); Yoghurt to the rescue (E Jean Higgie); Donor insemination (C Makin; S Robinson); The doctor's place in the team (S Hagard).... 1474
Lithium carbonate in schizoaffective states W W Sargent, FRCPsych..... 1470		
Endocrine versus cytotoxic treatment in advanced breast cancer B S Mantell, FRCP..... 1470		

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Drug monitoring and adverse reactions

SIR,—I have read with interest the recent contributions to your columns regarding improved means of identifying adverse drug reactions.¹⁻⁴ I am particularly interested in the debate on the direct involvement of patients in this process and would like to report a recent experience.

In a comparison between high-dosage cotrimoxazole (three tablets twice daily for seven days) and phenoxypenicillin in the treatment of suspected streptococcal sore throat patients recorded their own progress daily on numbered, reply-paid postcards to be returned after two weeks. The cards included a blank space for recording and describing side effects. One hundred and thirteen of 139 patients (81%) returned cards. Of 58 cotrimoxazole-takers, eight (14%) reported rashes, 11 (19%) gastrointestinal side effects ranging from "painful swollen tongue" to "nausea," and four (7%) non-specific complaints. The 55 penicillin-takers acted as controls, having presented similar illness and been investigated and prescribed medicine similarly. Fifty-five penicillin-takers reported no rashes and no gastrointestinal side effects and six (11%) non-specific complaints.

The following points are relevant to the present debate:

(1) Six of the eight rashes (all apparently macular and in adults) appeared on the last day of treatment or later, beyond the time of planned clinical observation. Only five of the eight were reported to the general practitioner.

(2) One of the rashes was initially diagnosed by the doctor as rubella; however, subsequent serological examination showed this diagnosis to be untenable.

(3) An error of documentation by one doctor

resulted in the incorrect initial reporting of two rashes as being in penicillin-takers. This resulted in the trial continuing beyond the point of unacceptable incidence of side effects.

(4) A referee who advised against publication of the findings described above as a "Short Report" included in his comments the opinion that "gastro-intestinal side effects... are not usually complained of from this drug" and concluded that the study and method used were "quite unfair to" cotrimoxazole.

This is not my first experience of obtaining a high response rate and valuable information by patient reply card⁵ and I wish to support strongly the suggestion by Dollery and Rawlins¹ that this inexpensive method be used more fully. A degree of patient selection is probably necessary to exclude those unlikely to co-operate adequately and a form of random allocation between study drug and conventional drug or placebo required as a control for the method of surveillance. At the same time I must record serious reservations about the methods proposed by Crombie.⁴ Retrospective use of morbidity data is surely the method least able to compensate for under-reporting by patients and has no defence against non-recognition of events, especially apparently minor ones, by the doctor. No check of patient compliance can be included and assessment of the possible role of concurrent or self-medication cannot be adequate. Further, the matching of patients receiving a drug for one illness with those consulting for different disorders or not consulting for any illness can be justified only if there is evidence that the different groups of patients recognise and report illness with the same frequency, unless of course the event being evaluated is

of a kind unlikely to be overlooked by patient or doctor.

J G R HOWIE

Department of General Practice,
University of Aberdeen

¹ Dollery, C T, and Rawlins, M D, *British Medical Journal*, 1977, **1**, 96.

² *British Medical Journal*, 1977, **1**, 862.

³ Lawson, D H, and Henry, D A, *British Medical Journal*, 1977, **1**, 491.

⁴ Crombie, D L, *British Medical Journal*, 1977, **1**, 1217.

⁵ Howie, J G R, and Clark, G A, *Lancet*, 1970, **2**, 1009.

Diet and coronary heart disease

SIR,—Dr K P Ball (21 May, p 1346) continues to quote a large number of out-of-date reports and to ignore the recent evidence mentioned in my letter (30 April, p 1155). The US report which he quotes is "prepared by the staff" of Senator McGovern's committee, which, however, is still taking evidence. I have no quarrel with reduction of fat to maintain an optimum weight, but there is no need to change the nature of the fat we consume.

It is a sobering thought that so many in our profession have allowed themselves to be brainwashed by propaganda into a widespread acceptance of a dietetic fashion which can only be transient. The leaders in this propaganda movement are presenting their (mainly epidemiological) "research" directly to the public over the media without subjecting it to debate and criticism in a proper scientific forum. Before these permissive days this propaganda would have been regarded as "infamous conduct in a professional respect."

The matter can be resolved only by a proper assembly of informed scientists. When this was done recently in private session the lipid theory of causation was routed. Never before has it been suggested that such important scientific matters can be resolved by "Gallup polls" of doctors or even committees. Norum¹ conducted a poll among 214 doctors, only a

few of whom were in a position to give an informed opinion; 15% did not reply but those who did were obviously indulging in wishful thinking that the diet regimen would be helpful as 92% of them were already themselves on the diet! It is time that we neutralised such unacceptable "evidence."

Professor Jens Dedichen,² of Oslo, recently admitted responsibility for introducing low-fat diets following observations by himself and Strom 25 years ago on the decline of recorded coronary deaths in the war years, which might have been related to fat deprivation. From 1955 he strongly pursued this policy and stuck to a low-fat diet himself, criticising colleagues who would not support him. He now confesses that in spite of the Norwegian effort (soy bean oil consumption in Norway increased fivefold) there had been no fall in coronary mortality but rather a steady increase. He regrets the anxiety created by this advice to the population. During these 25 years "it has become increasingly clear that we are on the wrong track." Could increasing consumption of polyunsaturated fats in Norway have caused the increase in coronary mortality? We do not know.

Professor Bengt Borgstrom,³ of Lund, an outstanding leader in fat biochemistry, says that pursuit of these policies of saturated fat reduction and more unsaturated fats is "irresponsible, as it would have far-reaching economic, agricultural, and psychological effects." There might be quite unknown consequences of altering the nature of our dietetic fats. Even now we can be certain that a diet rich in polyunsaturated fats will increase the risk of gall stones.

With the absence of any proof of benefit from reduction of blood lipids we have reached the stage where we must admit that this regimen has failed when tried out on subjects at risk of coronary disease and therefore the imposition on the public of dietetic restrictions for its prevention has no scientific justification.

The fatty streaks which occur in young children are not the beginning of atheroma. The Bromley Health Authority⁴ has been advising mothers of babies about the avoidance of eggs and discarding the cream on the top of the milk. This is creating alarm in the public and it is urgent that our professional advice should get back on to an even keel and put a stop to this propaganda, which threatens to become a cruel imposition terrifying mothers of families.

JOHN McMICHAEL

London NW11

¹ Norum, K, *Tidsskrift for den Norske Laegeforening*, 1977, **97**, 363.

² Dedichen, J, *Tidsskrift for den Norske Laegeforening*, 1976, **96**, 915.

³ Borgstrom, B, *Sartryk ur Livsmedelsteknik*, 1976, **7**, 302.

⁴ Leaflet issued by Bromley Area Health Authority and produced with the support of the Flora Project for Health Education.

MRC treatment trial for mild hypertension

SIR,—On p 1437 of this week's *BMJ* an account is given of the pilot phase of the Medical Research Council's multicentre treatment trial for mild hypertension.

If it is undertaken, and this is under present consideration, the main trial would need to recruit another 150 clinics, most of which are likely to be group practices. The need to

recruit quickly so that the trial would give an answer in the shortest time means that we should now see which practices would be interested in collaborating. In the pilot stage financial assistance was provided towards the compiling of age/sex registers and for additional nursing or medical sessions required. Equipment, laboratory facilities, postal arrangements, and compensation for the use of cardiographs were also provided and no practice was out of pocket as a result of its participation. It would be expected that this pattern would continue.

Practices with total lists of at least 7500 (preferably 10 000 or over) are required and ideally those with accommodation which will allow one room to be used specifically for follow-up clinics at regular times each week (inside or outside normal surgery hours) and for storage of records, drugs, and instruments. The initial blood-pressure screening of those aged 35-64 years would be carried out by locally recruited and specially trained nurses, largely in purpose-built mobile clinics which would relieve busy surgeries of much of the additional load imposed by the trial.

Several centres currently participating are willing to be contacted or visited by others wanting to know more about what the trial involves. Their names and addresses and further information on the study can be obtained from the Co-ordinating Centre, MRC Treatment Trial for Mild Hypertension, MRC/DHSS Epidemiology and Medical Care Unit, Northwick Park Hospital, Watford Road, Harrow HA1 3UJ.

W S PEART
Chairman

Medical Unit,
St Mary's Hospital,
London W2

W E MIALL
Secretary,
MRC Working Party on Mild
to Moderate Hypertension

MRC DHSS Epidemiology and
Medical Care Unit
Northwick Park Hospital,
Harrow, Middx

Breath, alcohol, and the law

SIR,—Dr B M Wright (7 May, p 1216) is wrong; we would take issue with him on his contention that our results are "much worse than those obtained in recent years by a number of reputable observers." Dr Wright does not name these reputable observers but by implication they seem to include himself. Of the two papers he quotes in this connection, that of Jones, Wright, and Jones¹ claims that a comparison between capillary blood values and those of breath shows a line of best fit at a ratio of 2300:1, but no hard data are provided to allow an independent check. However, in a separate and earlier publication which is not quoted by Dr Wright, Jones² reported a blood:breath ratio which averaged 2361:1. What is not clear is whether Dr Wright and his colleagues have carried out a separate independent study or whether the results presented have been derived from the earlier work of Dr Jones. Perhaps Dr Wright will enlighten us on this point.

The second paper to which Dr Wright refers, that by Dr Dubowski,³ contains much that is interesting and valuable but it contains no quantitative data on the blood:breath ratio that can usefully be compared with our own results. Thus the only data available for comparison are those of Dr Jones, and these do

not differ substantially from our own figures, especially if it is taken into account that in our study breath analysis was begun immediately after drinking had ceased—that is, before equilibrium had been reached and when the effects of mouth alcohol were still evident.

Finally, we would emphasise yet again that breath analysis is not an acceptable method for accurately determining blood alcohol concentrations.

J P PAYNE
D W HILL

Research Department of
Anaesthetics,
Royal College of Surgeons
of England,
London WC2

¹ Jones, A W, Wright, B M, and Jones, T P, *Proceedings of the 6th International Conference on Alcohol, Drugs, and Traffic Safety*, p 509. Toronto, Addiction Research Foundation, 1975.

² Jones, A W, PhD Thesis, University of Wales Institute of Science and Technology, 1974.

³ Dubowski, K M, *Clinical Chemistry*, 1974, **20**, 294.

Malaria in returning travellers

SIR,—We wish to illustrate, using our recent experience, the ever-increasing trend of imported malaria in the UK mentioned by Dr Salil K Ghosh and I Nicholson (30 April, p 1136). Nineteen cases of malaria have been diagnosed in our health area since the beginning of the year; nine of them have presented since the beginning of May. All but one have been infections with *Plasmodium vivax* and have occurred in Asians returning from the Indian subcontinent; the exception was in a young soldier found to be infected with *Plasmodium falciparum* on returning from a tour of duty in Gambia.

May we remind our clinical colleagues of the possibility of malaria when considering febrile patients from abroad with possible infection, since we are now at the beginning of the tourist season? Furthermore, we would suggest that wider publicity should be given to the Asian communities so that they are made aware that on returning to Asia or Africa they may well acquire malaria there unless adequate precautions, such as antimalarian prophylaxis, are taken while overseas.

We are indebted to our colleagues in the haematology departments of our hospitals for giving us the opportunity to see blood films and acquire data on the cases quoted.

R G THOMPSON
I A HARPER

Public Health Laboratory,
New Cross Hospital,
Wolverhampton

A M PATEL
W S A ALLAN

Haematology Laboratories,
Wolverhampton Area Health Authority

Making better use of our nurses

SIR,—Your leading article (21 May, p 1306) makes it appear that it is the "attached" nurses employed by area health authorities who work only 23 hours a week. This confuses our finding¹ that two-thirds of general practices in England have one or more nurses "attached" to them with our finding that it is the nurses employed by the general practitioners themselves who work an average of 23 hours a week. In our survey¹ we did not inquire about the work of attached nurses, but since then we have