CORRESPONDENCE

Unused sphygmomanometer H A Fleming, FRCP..... 1659	Aerosols containing neomycin I Friedmann, FRCPATH..... 1662	Peritoneal fibrosis R D Rosin, FRCS..... 1664
Testing dyspeptic patients S Lipetz, FRCPG..... 1659	Side effects of sodium valproate M H Bellman, MRCP, and E M Ross, MRCP..... 1662	Tetracycline and toxoplasmosis P L Grossman, MD, and J S Remington, MD..... 1664
Anabolic steroids and anticoagulants C W Howard, MRCP, and others..... 1659	Basal body temperature R F Vollman, MD..... 1662	Output of doctors J Parkhouse, FFARCS..... 1665
Tuberculosis among immigrants in Glasgow K M Goel, MD, and others..... 1660	Guillain-Barré syndrome and influenza vaccine W Ehrengut, MD..... 1662	Shortage of anaesthetists J M Small, FRCS; C Langmaid, FRCS; C Brun, FRCS, and D B Murray, DA; R M Heddle, FRCS, and M R Lock, FRCS..... 1665
Diet and coronary heart disease Lindsay M Elliott, MB..... 1660	New strategies for drug monitoring R H B Meyboom, MD..... 1663	Peers or leaders? R E Smith, FRCP..... 1666
Risk of coronary heart disease in different populations A R P Walker, DSc, and C Isaacson, FRCPATH..... 1660	Alkaline phosphatase isoenzymes in dialysis bone disease Th Mountokalakis, MD, and others..... 1663	Risks and rewards G T Watts, FRCS..... 1666
Hazards of the sauna S E Josse, MRCPG..... 1660	Chronic urticaria after vasectomy R P Warin, FRCP..... 1663	The cowardice continues J P Anderson, FRCPED..... 1666
Suppression of intractable cough C J Stewart, MD, and T J Coady, MB..... 1660	Sexually induced headaches N R Nutt, MB..... 1664	Distinction awards A P J Ross, FRCS; H Jacobs, MRCPsych..... 1667
Randomised clinical trials S J Pocock..... 1661	Psychologically mediated abdominal pain I Taylor, FRCS..... 1664	A confidence trick M Jones, FDS; J P Beavis, FRCS, and G N Boobbyer, FRCS; P W M Copeman, FRCP... 1667
Delivery of postgraduate education P J Tyrer, MD..... 1661	Management of the stove-in chest with paradoxical movement C P de Fonseca, FRCS..... 1664	Professional indemnity P G Stableforth, FRCS..... 1667
Management of elderly demented patients J M S Pearce, FRCP..... 1661	Transient hypotension following intravenous ethamsylate (Dicynene) B Watson..... 1664	Resignation by profession R R Drury, MB..... 1667
Cholestyramine in uraemic pruritus O M Wrong, FRCP..... 1662		National Insurance contributions J Kelly, MRCPG..... 1667

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Unused sphygmomanometer

SIR,—The two papers by Dr R F Heller and Professor Geoffrey Rose (4 June, pp 1441 and 1442) are appropriately published in the same issue as the preliminary report of the MRC pilot trial on mild to moderate hypertension. It has long been my contention that we should tackle much more adequately the severe hypertension that is already known to us.¹

As Heller and Rose indicate, many opportunities to measure the blood pressure are not taken. This even occurs when it is crying out to be measured. Not rarely one finds the modern general practitioner getting an electrocardiogram done long before using a sphygmomanometer even in a patient on the contraceptive pill complaining of severe headaches.

When raised blood pressure is recorded it is frequently not acted upon. In hospital the measurement may be made but may not be communicated to the general practitioner. One sees patients with complicated hypertension in whom a search of their old hospital notes shows that the anaesthetic chart years before records gross hypertension which has never been mentioned again.

It is my belief that we as a profession have some sort of mental block about the use of a sphygmomanometer and that the public might be better served if the health centre nurse or some other suitable person routinely recorded the blood pressure of all patients over the age of 40 attending their general practitioners.

Repeat observation should be made at suitable intervals.

HUGH A FLEMING

Cambridge

¹ Fleming, H A, *Cardiology, Peterborough Symposium*, p 16. London, Pitman Medical, 1975.

Testing dyspeptic patients

SIR,—The article on the "Use of barium meal examination in dyspeptic patients under 50" (4 June, p 1460) points out that of 100 dyspeptic patients under the age of 50 who underwent barium meal investigation no abnormality was found in 76. We described¹ a study in general practice showing the importance of stool benzidine testing in the examination of alimentary ulceration and comparing the results with those found by subsequent x-ray investigation. In 36 such cases a positive stool benzidine test led to the final diagnosis although the radiological findings were repeatedly negative. Perhaps the significance of testing for occult blood in the stool is not sufficiently appreciated.

S LIPETZ

Edinburgh

¹ Lipetz, S, Sklaroff, S A, and Stein, L, *British Medical Journal*, 1955, 2, 172.

Anabolic steroids and anticoagulants

SIR,—We should like to report the possible hazard of the potentiating effect of anabolic steroids on anticoagulant therapy. This has been reported previously¹⁻³ but is not in our opinion widely appreciated. It seems possible that the use of these agents will increase, particularly in those patients with recurrent deep venous thrombosis who have not responded adequately to anticoagulant therapy.

Recently we have had experience of prolongation of thrombotest times following commencement of stanozolol treatment in patients previously well controlled on dicumarol anticoagulation. One patient, a 42-year-old lady with recurrent deep venous thrombosis, was admitted with haematuria 16 days after starting stanozolol treatment at 5 mg b.d. Thrombotest on admission was 1% (20 min). Treatment was given by infusion of fresh frozen plasma. In view of this experience a second patient, a 41-year-old man with deep vein thrombosis following hernia repair, was recalled 12 days after commencing stanozolol treatment in the same dosage. Thrombotest was 3.5% (225 s). Anticoagulation was discontinued and restarted in lower dosage.

The effect of anabolic steroids on anticoagulant tolerance seems to depend on the chemical structure of the compound, the greatest reduction in anticoagulant requirement being produced by C₁₇-alkylated derivatives of testosterone. The 17 alkyl group is introduced in the steroid molecule with the purpose of making anabolic steroids active by mouth. C₁₇-alkylated steroids are known to cause abnormalities of some liver function tests, abnormal BSP (bromsulphalein) retention, and increased serum transaminase levels.