persistent cough following radiotherapy. An inhalation of lignocaine relieved the cough within two hours and the effect continued for a week when a further inhalation was given. Since that time there has been no recurrence of the severe coughing fits.

Each of these patients obtained relief with lignocaine 400 mg in saline administered via a Bird Micronebulizer. Salbutamol 2.5 mg was given by wet inhalation immediately before each treatment. It is our impression that mucus clearance was not impaired in these three patients. From this very limited experience we would not withhold lignocaine aerosol treatment in any patient who had distressing and otherwise uncontrollable cough and was suffering a fatal illness. We would agree with you that further evaluation of this technique is necessary in other types of illness giving rise to severe cough, particularly until the possible effects on mucus clearance is better understood.

C J STEWART
T J COADY

Ipswich Hospital,
Ipswich

¹ Howard, P, *et al*, *British Journal of Diseases of the Chest*, 1977, **71**, 19.

Randomised clinical trials

SIR,—Your leading article (14 May, p 1238) presents a good case for randomised clinical trials (RCTs). Alas, I feel that many investigators will still be unconvinced, arguing that non-randomised trials are justified if no sources of bias can be identified. Thus, I would like to present some evidence that bias can exist without any apparent explanation.

In the United States cancer chemotherapy co-operative groups it is common practice to include the same control treatment in consecutive RCTs. At the statistical laboratory in Buffalo, New York, I identified 19 such pairs of trials, mostly in advanced lung cancer, and compared the annual death rates of the two groups of patients on the same treatment:

The trials were fairly large, most having over 100 patients per treatment, and there were no known sources of bias. One would have expected little change in death rates on the same treatment from one trial to the next, whereas, in fact, the changes ranged from -46% to +24%. In four instances the change was statistically significant at the 5% level (using a two-sided F-test for comparing two exponential death rates) and a further six changes were significant at the 20% level. Such marked evidence of differences between trials indicates that any comparison of treatments not within an RCT must be deemed highly suspect.

Regarding the failure to achieve sufficient patients, a current survey on the size of RCTs conducted by P Armitage, D A G Galton, and myself provides some interesting evidence. From a random sample of 50 cancer trials registered with UICC during 1972-5 our preliminary results based on 30 replies to a questionnaire indicate a median accrual rate of

34 per annum (total on all treatments) and a median of 5 years' accrual to achieve the prespecified number of patients. Evidently many will not achieve their accrual targets, so that although the design and execution of these trials were on the whole satisfactory a substantial number will prove inconclusive owing to the failure to make a realistic assessment of patient accrual.

This survey also showed that nearly all trials have repeated analyses of results, the most common interval between analyses being 6 months. Neither fixed nor sequential designs cope satisfactorily with such periodic assessments, but some recent developments in "group sequential" designs do provide a proper statistical basis for this approach.¹⁻³ A further point from the survey was that two-thirds of trials incorporated stratification for prognostic factors in the randomisation. Thus your leading article's declaration that stratification is unnecessary is not being followed, and I think this is sensible in view of the fact that retrospective adjustment in analysis requires rather sophisticated statistical methods and even these are useless in cases of severe imbalance.

Also, I think the suggestion of an unequal randomisation (say, a 2:1 ratio), whereby a greater proportion of patients receive the new treatment, is to be encouraged. This is especially true if there are suitable historical data on the old treatment, since there exist statistical methods for incorporating such extra information into the trial's design and analysis.⁴ Lastly, I think the conversion of investigators to RCTs would be enhanced considerably if medical journals took a firmer stance in refusing to publish the results of non-randomised studies when an RCT was appropriate.

STUART J POCOCK

Medical Computing and Statistics
Group,
Medical School,
Edinburgh

¹ McPherson, K, *New England Journal of Medicine*, 1974, **290**, 501.

² Pocock, S J, *Biometrika*, 1977 (in press).

³ Canner, P, *Biometrics*, 1977 (in press).

⁴ Pocock, S J, *Journal of Chronic Diseases*, 1976, **29**, 175.

Delivery of postgraduate education

SIR,—Professor A H Crisp (28 May, p 1397) has illustrated admirably the dilemma that faces many university departments in trying to strike a balance between undergraduate and postgraduate teaching responsibilities. In Wessex we were more fortunate than Professor Crisp in that a well-established postgraduate teaching programme existed long before the medical school at Southampton was opened, and to maintain the high quality of this programme academic sessions were specifically allocated for this purpose. Nevertheless, problems still persist, and in addition to the salaried sessional commitments by clinical tutors, suggested by Professor Crisp, a reappraisal of the teaching responsibilities of academic staff is necessary.

Although for historical and administrative

reasons university clinical departments are funded according to their undergraduate teaching commitments it is impossible (and inappropriate) for them not to be associated with postgraduate training. This should be recognised as a formal commitment to maintaining academic standards through their region. This commitment can to some extent be balanced by greater involvement of clinical teachers in undergraduate teaching. In our undergraduate teaching course in psychiatry we involve many consultants throughout the region, who devote considerable time and effort but who receive no recognition apart from the title of honorary clinical teacher. Formal recognition is needed of these changes in the job descriptions of both academic and clinical staff in addition to the extra funding that Professor Crisp suggests if we are to maintain a healthy integration between academic and clinical teaching responsibilities.

P J TYRER

General Hospital,
Southampton

Management of elderly demented patients

SIR,—Your leading article (21 May, p 1301) on the management of elderly demented patients is aptly coloured in pessimistic lines familiar to those who have dealt with such patients and families. Having referred to a recent conference¹ you lapse immediately into an attitude of *caring* for what seems to be assumed to be an a priori irreversible state. I wish to make two points.

(1) Although you mention "accurate diagnosis," this is quickly passed over. Recent work^{2,3} indicates that a reversible cause for dementia is present in between 10 and 20% of patients referred to both specialist units and district general hospitals provided that investigation is adequate. Age alone is no guarantee that dementia is caused by irrevocable cerebral degenerative pathology. Such facts are politically unpalatable in times of stringency, because the tests necessary to the proper study of demented patients include time-consuming and expensive metabolic and neuroradiological investigations, not least the EMI scan. It is, however, our professional duty to patients to indicate clearly that unless these methods are applied to the vast numbers involved (at least 5% of the over-65s, who number several million in the UK) many will be thrown on to the scrap heap and will require unnecessarily the expensive facilities of hospitals, hostels, and day centres your article describes so well.

(2) In discussing management you fall back on the trendy euphemism of the caring team (*sic*). You say, "the doctor's role is as a member of the caring team, *not necessarily as its leader* . . ." (my italics). Is it not time we dispelled this currently fashionable notion so widely advocated by the radical chic on TV and radio? The exclusion of the doctor from leading the team, assuming that a team is needed in all cases, leads to the omission of proper diagnostic methods, the consequent prognosis, and to the pitiful retreat implicit in the practice of subsequent care. The properly trained doctor is the only person capable of initiating the correct sorting out and selection of patients for investigation. And he is the only person capable of talking to relatives in a manner based on the full understanding of the patients' disease in its complex pathological,

clinical, and therapeutic aspects. If the disease is shown to be an irreversible one, then ancillary workers are most helpful in assisting with the difficult and important skills of subsequent psychosocial adjustment and management.

But please, Sir, when we are discussing illness (as opposed to the more vague, pure social maladjustments of contemporary society) let us not be falsely persuaded by the fashionable soft sciences into abrogating our inalienable role as leader of the clinical team.

JOHN M S PEARCE

Hull Royal Infirmary,
Hull

¹ Anderson, W F. and Carlton-Ashton, J R, eds, *Age and Ageing*, 1977, suppl.

² Harrison, M J G, and Marsden, C D, *Archives of Neurology*, 1977, **34**, 199.

³ Pearce, J, and Miller, E, *Clinical Aspects of Dementia*. London, Baillière Tindall, 1973.

Cholestyramine in uraemic pruritus

SIR,—Most renal physicians find uraemic pruritus difficult to treat, so the report of Dr D S Silverberg and his colleagues (19 March, p 752) of a favourable response to cholestyramine will excite interest and encourage others to try this treatment.

The authors realise that cholestyramine has the potential to cause acidosis but found no evidence of this in their patients. The absence of acidosis was probably the result of haemodialysis three times a week, which would return the acid-base state to normal and prevent cumulative hydrogen ion retention. But dialysed patients are less liable to this complication than uraemic non-dialysed patients, and the latter would be particularly at risk from the acidifying effect of cholestyramine, as the following argument makes clear.

Cholestyramine is a strongly basic anion-exchange resin in the chloride form, with an exchange capacity of 4.1 mEq/g. A study of several similar anion-exchange resins some years ago found that after oral ingestion the resin was excreted in the stool predominantly with organic anion and bicarbonate, and only 15% in the chloride form.¹ By extension of this finding to cholestyramine, every gram of resin taken would contribute 3.5 mEq of chloride to body fluids and remove in exchange 3.5 mEq of bicarbonate or bicarbonate-precursor from the body. Loss of bicarbonate from the body is equivalent to a gain in hydrogen ion, so the 10 g dose of cholestyramine used daily by Dr Silverberg would donate 35 mEq of hydrogen ion to body fluids, an amount which is only a little less than the daily excretion of hydrogen ion by healthy kidneys. It might be argued that this would not occur in a uraemic subject because of a change in the ionic pattern of intestinal juices; however, a study of this point has been made and showed no significant difference in the anion pattern of faeces from uraemic and normal subjects.² There are therefore strong a priori grounds for predicting progressive hydrogen ion retention in non-dialysed uraemic subjects given cholestyramine. Although Silverberg referred to *hyperchloraemic* acidosis, the resultant acidosis might not be accompanied by hyperchloraemia because of the retention of unmeasured anion, which is usual in renal failure.

This problem would not arise if the manufacturers marketed cholestyramine in acetate or lactate form rather than as chloride. The necessary conversion can easily be carried out

in a laboratory with the simplest equipment, such as exists in hospital pharmacies, but this would remove the added flavouring which is important for palatability, and the conversion would therefore be best incorporated during manufacture. Cholestyramine modified in this way would be just as effective as the present chloride form for the other therapeutic purposes for which it is used.

Incidentally, the name "cholestyramine" is misleading, as it implies a particular mode of action (binding of bile anions), whereas the approved names of most modern drugs are derived from the chemical structure of the drug alone. Anion exchange resins remove much more from the body than bile anions, which are not retained in renal failure, and any effect they may have in relieving uraemic pruritus is likely to be mediated by some other action.

O M WRONG

University College Hospital
Medical School,
London WC1

¹ Hurst, P E, *et al*, *Clinical Science*, 1963, **24**, 187.

² Wilson, D R, *et al*, *Clinical Science*, 1968, **35**, 197.

Aerosols containing neomycin

SIR,—I refer to your "Warning on Aerosols containing Neomycin" (21 May, p 1360). May I draw your attention to two illustrations, 19.1 and 19.2 (pp 454 and 455), in my textbook,¹ which show the extensive damage to Corti's organ and its innervation caused by neomycin aerosols in two children with cystic fibrosis treated in a neomycin atmosphere for several months.

I FRIEDMANN

Northwick Park Hospital,
Middlesex

¹ Friedmann, I, *The Pathology of the Ear*. Oxford, Blackwell, 1974.

Side effects of sodium valproate

SIR,—While we agree that sodium valproate (Epilim) is a most useful anticonvulsant in childhood, we should like to report an apparent side effect not mentioned in your leading article (16 April, p 986).

A 14-year-old Asian boy was born in hospital in Kenya. Delivery was apparently normal, birth weight 3 kg. From the age of 6 months he started attacks of eye rolling, rigidity, and shaking, which lasted about three minutes and occurred about every two weeks until he was 5 years old, when they became less frequent. He came to the UK in 1969 and now attends a boarding school for educationally backward children. Clinical assessment revealed bilateral optic atrophy, clumsy gait, but no obvious syndrome. Head circumference was at the 10th centile and skull x-ray was normal.

Since 1973 he has had occasional brief episodes of loss of vision and jerking of the head followed by intermittent "blackouts." Electroencephalography showed epileptic features, and he was prescribed phenobarbitone. The convulsions became more frequent, occurring two or three times a week, and treatment has included various combinations of anticonvulsant drugs, including sodium valproate, with no definite success. The regimen was changed recently to sodium valproate alone at a dose of 42 mg/kg (1600 mg) per day. After 14 days free from convul-

sions he became confused and had recurrent episodes of bizarre behaviour in which he stared vacantly, shouted gibberish, and had frightening hallucinations. He was visited by a general practitioner deputy, who stopped the drug. Two days later he was seen at hospital, when he was still moderately confused and appeared frightened though apparently had steadily improved since the sodium valproate had been stopped. Plasma valproate level at that time was 13 µg/ml. The drug was recommenced at half the previous dose (800 mg per day). Ten days later his behaviour had returned to normal and he remained free from convulsions.

The clinical course of this patient's acute episode of psychotic behaviour in relation to his medication is suggestive of a drug reaction. The plasma level two days after discontinuing sodium valproate is surprisingly high. Considering the average half life of the drug in epileptic patients to be about six hours, the level at the time of the reaction would have been far above the expected therapeutic range. Careful questioning did not suggest that an overdose had been taken (this would have been unlikely as the patient had been on anticonvulsant drugs for many years under the supervision of the excellent nursing sister and medical officer at his special school). We have found no other reports of similar side effects of sodium valproate when given on its own and suggest that a watch should be kept for psychotic changes in patients on this drug.

M H BELLMAN

EUAN M ROSS

Central Middlesex Hospital,
London NW10

Basal body temperature

SIR,—In their paper "Problems in using basal body temperature recordings in an infertility clinic" (26 March, p 803) Dr Elizabeth A Lenton and others are trying to achieve the impossible. They asked "experts" and "non-experts" to diagnose a small mixed sample of BBT curves from healthy volunteers (8), infertility patients with ovulatory cycles (14), with anovulatory cycles (24), and pregnant cycles (4). Such a hodgepodge of variables cannot be analysed statistically and meaningfully. Further, the authors did not provide definitions according to which the BBT curves should be examined, nor were the experts and non-experts asked to spell out their own diagnostic criteria. Thus, the results reflect the participants' guesses but not the diagnostic value of the BBT curve.

The elevated phase of the biphasic BBT curve indicates corpus luteum activity, and consequently an ovulatory menstrual cycle. However, the BBT curve does not allow us to time precisely the day of ovulation.

RUDOLF F VOLLMAN

Switzerland

Guillain-Barré syndrome and influenza vaccine

SIR,—Your leading article on Guillain-Barré syndrome and influenza vaccine (28 May, p 1373) stimulates me to consider the various possibilities of neural disturbances after influenza immunisation. We differentiate the