tactics summed up by Stevenson in his book *Ethics and Language*.¹ Teachers may resort to emotive techniques in order to change the attitudes of children towards various methods of behaviour.

It is my opinion, however, that any attempt at crude emotivism will fail. We should present the children with the medical facts against a background of caring, now about their parents, later about their dependants.

V L IRWIN

St Edmund's College,
Ware, Herts

¹ Stevenson, C L, *Ethics and Language*. New Haven, Yale University Press, 1944.

Flautist's chin: a companion to fiddler's neck

SIR,—A 32-year-old female amateur flautist presented with a persistent eruption on her chin which had been present intermittently for two to three years. The only symptoms were of discomfort when playing the flute, but the eruption subsided if she stopped playing for two to three weeks. Examination revealed acneiform lesions confined to the central mid-portion of the chin consisting of inflammatory pustules, papules, and some slight hyperpigmentation. Elsewhere on the face there were a few inconspicuous open comedones and there was mild seborrhoea. She had not had significant acne as a teenager. Oxytetracycline 250 mg daily produced rapid improvement and she was able to toot her flute in complete comfort.

Peachey and Matthews¹ described an acneiform eruption on the necks of violinists and viola players related to contact with and friction from their instruments. A similar situation pertains to the chins of flautists with the additional factor of constant wetting with saliva. "Flautist's chin" and "fiddler's neck" appear to be close companions in the acne orchestra demonstrating variations on the theme of acne mechanica.² Perhaps more musical diseases such as "cellist's thigh" or "bagpiper's midriff" will yet be recorded.

M G C DAHL

Dermatology Department,
Royal Victoria Infirmary,
Newcastle upon Tyne

¹ Peachey, R D G, and Matthews, C N A, *British Journal of Dermatology*, 1978, **98**, 669.

² Mills, O H, and Kligman, A, *Archives of Dermatology*, 1975, **111**, 481.

*. Examples of "guitar nipple," "guitarist's groin," and even "cello scrotum" were reported in these columns during 1974.—ED, BMJ.

Thyrotoxic Graves's disease after primary hypothyroidism

SIR,—We were interested to see the report by Dr S Olczak and others (2 September, p 666) of a patient who developed thyrotoxicosis after well-documented primary hypothyroidism, for we have recently seen a similar case.

Our patient, who is a female nursing officer aged 56, was seen in 1975 complaining of increasing tiredness for one year, a husky voice, puffiness of the eyelids, mental and physical slowness, marked intolerance to the cold, and slight ataxia. There was no significant past history and there was no family history of thyroid disorder, diabetes

mellitus, or pernicious anaemia. On examination she had a dry skin with a puffy face, typical of hypothyroidism, the pulse was 80/min, there was slow relaxation of the deep reflexes, and the thyroid gland was not palpable. The serum thyroxine was <10 mmol/l (<0.8 µg/100 ml), triiodothyronine <0.1 mmol/l (<6.5 mg/100 ml), and thyroid-stimulating hormone 80 mU/l. She was started on thyroxine and soon regained her normal health, being maintained on 0.2 mg daily.

She remained perfectly healthy for two years but in March 1978 she reported the loss of 1 stone (6.4 kg) in weight in spite of a good appetite and her colleagues had noted that she was nervous and jumpy. She was unusually tired in the hotter weather, but there was no excessive perspiration and no palpitations. She had noticed a little tremor. The serum thyroxine was 237 mmol/l (18.4 µg/100 ml) and triiodothyronine 5.3 mmol/l (6.35 µg/100 ml), so the dosage of thyroxine was reduced and ultimately stopped. Six weeks after stopping thyroxine completely the symptoms continued and she was more hyperactive; the pulse was 88/min, regular in rhythm; there was considerable swelling of the eyelids with minimal exophthalmos, and slight lid retraction with lid-lag; there was no pretibial myxoedema. The thyroid was just palpable on both sides. The serum thyroxine was 185 mmol/l (14.4 µg/100 ml) and triiodothyronine 6.0 mmol/l (0.39 µg/100 ml); thyroid autoantibodies were absent. She has been treated with carbimazole with marked improvement in her general symptoms, but there has been further aggravation of the exophthalmos. A thyroid scan, radioiodine uptake, and TRH-TSH stimulation tests were not performed. However, it is clear clinically that she has now developed Graves's disease with exophthalmos and thyrotoxicosis and it is equally clear that she had a history of hypothyroidism for at least one year before this diagnosis was established.

We have seen several patients who have had hypothyroidism and have then become euthyroid with a goitre typical of autoimmune thyroiditis, but we have not previously seen anyone develop frank hyperthyroidism and we have no further observations to explain this unusual phenomenon.

R B STOTT
J S STAFFURTH

Lewisham Hospital,
London SE13

Carcinoma of the breast in women under the age of 30

SIR,—I was very interested to see the analysis by Mr A S Purandare and Dr J Finbow of the ages of their female patients with carcinoma of the breast (9 September, p 771). In a somewhat smaller hospital than the Doncaster Royal Infirmary, serving a smaller population and sharing 60 general surgical beds, I have seen 213 female patients with the same condition. This is a very similar number to that seen by Mr Purandare and Dr Finbow in one year. The number of patients aged under 30 is the only significant difference in the age distribution in the two series (see table). Both series are small, but the incidence in this hospital of 1.4% aged under 30 agrees with the 1.2% figure given for Haagensen's very large number of cases.¹

Age distribution of women with carcinoma of the breast in two series

Age (years)	20-30		-40		-50		-60		-70		-80		>80		Total
	No	%	No	%	No	%	No	%	No	%	No	%	No	%	
Purandare and Finbow	10	5.1	16	8.2	45	23.1	43	22.1	42	21.5	31	15.9	8	4.1	195
Logan . . .	3	1.4	20	9.4	43	20.2	53	24.9	46	21.6	38	17.8	10	4.7	213

It would appear that Mr Purandare and Dr Finbow have been unfortunate to have had so many young patients, but it was worth while to bring their findings to the attention of all doctors. The three patients under 30 in my series were not suspected of having malignant tumours and this was only discovered after biopsy. None of these patients has been shown to have metastases after several years.

C J H LOGAN

Ulster Hospital,
Dundonald, Down

¹ Haagensen, C D, *Diseases of the Breast*, 2nd edn, p 352. Philadelphia, Saunders, 1971.

Hormone receptors in breast cancer

SIR,—Your leading article (8 July, p 77) gives an excellent survey of the situation with respect to hormone receptors in breast cancer. As you note, some tumours may be composed of a mixed population of receptor-positive and receptor-negative cells; continued growth of negative cells would then lead to hormonal insensitivity. In fact, direct evidence that hormone-responsive mammary tumours are heterogeneous populations of cells has been obtained in studies of mammary cancer in mice^{1,2} and rats.³ Endocrine ablation affects only the hormone-dependent cells in these tumours, leaving the hormone-independent cells intact.^{1,2} Progression from hormone dependence to autonomy in mammary tumours appears to be a manifestation of sequential clonal selection.¹⁻³ It seems likely that the hormone-responsive human breast cancers also are mixed populations of hormone-dependent and -independent cells. Thus breast cancers which have been treated with hormones, hormonal ablation, or antioestrogens practically all recur after a certain period of time and ultimately all become autonomous. This indicates that the endocrine treatment only stops growth of the receptor-positive cells and not of the receptor-negative cells. Better therapeutic results with these cancers might be achieved by combining endocrine treatment at an early stage with chemotherapy, when the autonomous cells are still only few in number. In this way perhaps only low doses of cytostatics suffice to kill all the autonomous cells present in the tumour. It would be worth while to compare this form of treatment with the conventional form in which chemotherapy is not given until autonomous cells have been given the chance to form a sizable tumour and high doses of cytostatics have to be applied to achieve regression.

Lipmann *et al*⁴ have reported that tumours with low or absent oestrogen receptor have a much higher response to chemotherapy than tumours with high oestrogen receptor values. They conclude that oestrogen receptor values are an important predictor of response to cytostatic chemotherapy. However, studies on mammary tumours in mice indicate that cytostatic treatment affects cells with different

receptor content to equal extent but has a larger effect on rapidly proliferating cells than on slowly growing cells.⁵ Recent data show that receptor-negative human breast cancers generally have a higher growth rate, as indicated by a higher labelling index and mitotic index.⁶ The conclusions by Lipmann *et al* may therefore have to be amplified to state that human breast cancers with low oestrogen receptor content usually are the more rapidly growing tumours and consequently are relatively more sensitive to cytostatic treatment.

MELS SLUYSER

Division of Endocrinology,
Antoni van Leeuwenhoek-Huis,
Netherlands Cancer Institute,
Amsterdam, Netherlands

¹ Sluysers, M., and Van Nie, R., *Cancer Research*, 1974, **34**, 3253.

² Sluysers, M., Evers, S. G., and De Goeij, C. C. J., *Nature*, 1976, **263**, 386.

³ Kim, U., and Depowski, M. J., *Cancer Research*, 1975, **35**, 2068.

⁴ Lipmann, M. E., *et al*, *New England Journal of Medicine*, 1978, **298**, 1223.

⁵ Sluysers, M., and Benckhuysen, C., *Cancer Treatment Reports*, 1977, **61**, 861.

⁶ Meyer, J. S., *et al*, *Cancer*, 1977, **40**, 2290.

Prevalence of multiple sclerosis in north-east Scotland

SIR,—We note with interest the comments of Dr A J Silman (2 September, p 703) concerning our assertion that the prevalence of multiple sclerosis in north-east Scotland shows a significant deviation from random distribution (29 July, p 314). We were not aware of any marked variation in the age distribution in the area and further statistical analysis of the 20-50-year age groups supports this. A subsequent study in 1973 (yet to be published) reveals a higher overall prevalence and higher rates for 22 out of the 28 districts. Many of the additional patients should have been included in the 1970 study but were missed at that time.

We would disagree with the second comment that the prevalence tended to increase nearer to Aberdeen. The map of the relative prevalence shows, for instance, that all Banffshire districts were above the mean despite the fact, as mentioned in the paper, that there might be a slight loss of patients from such peripheral areas to adjacent regions. The ease of access to Aberdeen therefore does not seem to be a factor, and in fact no part of the area is difficult of access except for short periods in the winter.

We would agree with the final comment that further studies of HLA status of patients in a high-risk region would be valuable and hope that such studies can be pursued locally.

DAVID I SHEPHERD

Wessex Neurological Centre,
Southampton

A W DOWNIE

Department of Medicine,
Medical School,
Aberdeen

Bacterial colonisation of small intestine in tropical malabsorption

SIR,—We enjoyed the paper by Professor G C Cook (22 July, p 238) and suggest that his demonstration of delayed small-intestinal transit in severe cases of tropical malabsorption, confirming earlier impressions from barium studies in southern India,¹ is of considerable importance. However, we do not

consider that the bacterial colonisation commonly present in such cases with² and without³ giardiasis is necessarily the result of abnormality of intestinal transit because we have shown that bacterial colonisation, albeit in lower numbers, also occurs in patients with mild tropical malabsorption without steatorrhoea⁴; such patients would have normal intestinal transit by Professor Cook's criteria. The aerobic flora of both our severe and mild cases, predominantly *Klebsiella pneumoniae*, is different from the anaerobic flora of patients with intestinal stasis.⁵

The finding of bacterial colonisation in patients with acute gastroenteritis and the demonstration that such organisms possess colonisation factors⁶ suggests that it may be characteristics of the bacteria rather than a change in intestinal transit which determine colonisation. Nevertheless, it is still far from clear which is the primary event in many intestinal diseases of the tropics and further studies are needed.

ANDREW TOMKINS
B S DRASAR
S G WRIGHT

Clinical Nutrition and Metabolism Unit,
Hospital for Tropical Diseases,
London NW1

¹ Baker, S. J., and Mathan, V. I., in *Tropical Sprue and Megaloblastic Anaemia*, ed Wellcome Trust, p 200. Edinburgh and London, Churchill Livingstone, 1971.

² Tomkins, A. M., *et al*, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1978, **72**, 33.

³ Tomkins, A. M., Drasar, B. S., and James, W. P. T., *Lancet*, 1975, **1**, 59.

⁴ Tomkins, A. M., *et al*, *Gut*, 1975, **16**, 392.

⁵ Drasar, B. S., and Shiner, M., *Gut*, 1969, **10**, 812.

⁶ Satterwhite, T. K., *et al*, *Lancet*, 1978, **2**, 181.

Nasogastric suction in acute pancreatitis

SIR,—I am sure that Dr R Naeije and his colleagues (2 September, p 659) are correct in their conclusion that nasogastric suction is unnecessary for most patients with mild or moderately severe pancreatitis, a conclusion which is probably applicable to the majority of the therapeutic advances made towards such patients.

Acute pancreatitis is a very variable condition and I feel that caution is needed before applying the results of a clinical study of this type to the routine management of the condition. A randomised study of 58 patients is perhaps rather lacking in numbers for any trend to appear when a fairly mild spectrum of disease is studied in a fairly young group of patients. Death has traditionally been the most useful end-point to use in the assessment of treatment for this disease, and in this study no patient died. It is well recognised that the clinical assessment of an attack of acute pancreatitis can be most misleading and patients with severe disease can appear to be quite mildly ill on first acquaintance. Furthermore, observations of the effect of acute pancreatitis in experimental animals indicate how common dilatation of the stomach can be, and this is also a feature of attacks in man, especially in the elderly.

While I would agree that the omission of this rather unpleasant form of treatment does not often influence the outcome of an attack, I think that it is very difficult to pick out the minority of patients in whom it is likely to be of benefit in a prospective manner. Thus we would omit nasogastric suction from routine management only in young, fit patients who appear to have very mild attacks of pancreatitis. The remainder have a nasogastric tube which

is withdrawn as soon as abdominal pain subsides (so long as aspirates are not high). In many patients the tube remains in situ for only a few hours.

M J McMAHON

University Department of Surgery,
General Infirmary,
Leeds

Toilet paper and spread of infection

SIR,—An important change in community hygiene has taken place in recent years with the introduction and increasing use of soft toilet paper. At first some people found the soft paper difficult to use owing to its porosity and lack of strength, but most were soon able to use it satisfactorily. Its superior cleansing power was demonstrated to me some years ago when I planned a long-term study of intestinal infection in a large children's home using soft paper in alternate cottages. The study had to be abandoned because the reduction in soiling of underclothing in the cottages using soft paper was so striking that the staff in the other cottages could not be persuaded to persist with the old paper.

In an intestinal infection spread largely by personal contact, such as Sonne dysentery, contamination of the clothing and of the skin of the perineum and buttocks can act as a source of continual infection of the hands, from where infection can pass to other persons. Soft toilet paper was introduced soon after the last war, but its great increase in popularity took place in the period 1957-71. Today the amount of hard paper in use is comparatively very small indeed. From 1957 onwards notifications of dysentery in England and Wales declined steadily. In the period 1957-9 the average annual notifications were 34 214. In the years 1969-71 the figure was 14 524 and in 1975-7 only 6776.

The mere fact that a reduction in dysentery and the introduction of soft paper were related in time does not, of course, prove that one is the cause of the other. The reduction in dysentery probably has a complex origin, arising from a number of different causes. It seems reasonable, however, to believe that the increased use of soft toilet paper may have been one factor in the decline.

IAN TAYLOR

Carshalton, Surrey

Hospital inquiries

SIR,—I was interested to read Dr D A Spencer's letter on hospital inquiries (16 September, p 828). I feel very strongly about the way in which the medical profession tends to be casual about the predicament in which some of our colleagues may find themselves as a consequence of such "inquiries." After one particularly well-publicised and extensively reported inquiry south of the Border the consultant concerned was publicly crucified by the chairman of the health board and the matter was given headline coverage in many national dailies. What was never adequately emphasised was that this man was doing the work of three consultants with poor junior staff and was indeed dying during this inquiry. I tried very hard to stimulate some interest in the profession at the time so that the grossly unfair verdict and the publicity received by this unfortunate consultant could be countered by the profession, but to little or no avail. The