Shortcomings of the NHS: a yawning chasm

SIR,—I am an NHS family doctor, one of a group who try to provide a competent 24-hour service to our patients. I am dispirited by the inadequacies of the service. It is not possible to practise adequately without access to x-rays, and acute admissions should certainly be available in emergency. In this area there are normally two radiologists working and they have a very heavy work load. Following the tragic death of one there has been a vacancy for most of this year which it is hoped to fill by the end of the year. Cuts had to be made and general practitioners found that no x-rays could be ordered except for chest films. This meant that patients needing x-rays had either to be referred to hospital clinics which continued to carry their normal work load or go privately 15 miles, which few can afford. Another alternative for the doctor is to order the x-ray to be performed, as there is no shortage of radiographers, and then himself attempt to interpret the films—a job in which he has had no training. Admittedly this shortage will soon be over, but one fears other shortages may not be long in developing. Apparently in some specialties suitably trained hospital staff are not keen to apply for consultant vacancies in NHS areas such as this, away from teaching centres.

Three patients of mine have died in the past three months, all in their fifties, who might have survived had there been adequate hospital facilities available for treatment. The most recent died at home of a coronary thrombosis about an hour after the only local hospital capable of providing treatment refused his admission as all their medical and intensive care beds were occupied. A few days earlier there was another fatal thrombosis in a man awaiting investigation with a view to coronary surgery. The head of the academic department concerned tells me that his waiting list for such cases is about one year. The third death was that of a man awaiting replacement of a heart valve, and his life would certainly have been prolonged had the operation been successfully performed.

It is popularly supposed that the NHS is under some stress but that only non-urgent cases are suffering. This is just not true. My patients express incredulity when the true situation is put to them. Family doctors have a contract with the NHS and it is arguable that they cannot fulfil this under the conditions described above. Resignation would, however, not help. There is likely always to be a gap between what the nation can afford and the latest technical achievements which are possible. At present it is not so much a gap as a yawning chasm and people might do well to look where they are standing lest they fall in.

G F G WOODMAN

Morpeth, Northumberland

Preventing deaths from malaria

SIR,—When I read the article by Dr A P Hall (23 September, p 877) I experienced a feeling of déjà vu. Following the Junior Members Forum of 1974, where the symposium was entitled "Medicine in the supersonic era," four motions were sent to the Annual Representative Meeting at Hull. The first, that legislation should be passed to compel airlines to issue malaria warning cards, was passed

as a reference to Council. The other three, requiring the travel industry to give adequate health information in holiday brochures, urging the DHSS to issue an annual gazetteer and establish a tropical diseases information centre manned 24 hours a day, and recommending the availability of malarial prophylactic drugs on NHS prescription, were passed by very large majorities and so became the policy of the Association.

Since that time the annual figures for cases of malaria in Britain have continued to rise at the same rate as previously and nothing is done to implement simple measures that might cut down on the number of needless deaths and time loss from illness. Dr Hall's article is a valuable reminder that the BMA has allowed the grass to grow under its feet and perhaps we should nudge the sleeping "Elephant" into activity.

MICHAEL J G THOMAS

Army Blood Supply Depot,
Aldershot, Hants

SIR,—With reference to Dr A P Hall's article on the above subject (23 September, p 877), so far as the major British airlines are concerned he is obviously misinformed. Information on malarial risk has been available to our passengers in booklet form over many years. As this method of communication had limited success, since 1977, owing to strong representations from our respective medical services, in-flight announcements have been made concerning malarial risk. These are made before landing in malarious areas or where onward travel can involve such a risk. It should be mentioned that we have been complimented by both the World Health Organisation and the Department of Health and Social Security for taking the lead in this field and it is noted that a number of foreign operators have now followed our example.

W DONALD MACKENZIE
Principal Medical Officer (Overseas),
British Airways

P J C CHAPMAN
Chief Medical Officer,
British Caledonian Airways Ltd

London (Heathrow) Airport,
Hounslow, Middx

Heat stroke in a "run for fun"

SIR,—Heat stroke is emerging as the principal medical emergency of mass participation community running. In accord with the report of Drs M R Nicholson and K W Somerville (10 June, p 1525) a mass participation 10-km run conducted in the normally temperate climate of Canada resulted in 15 people treated in hospital for heat stroke of varying severity.

The run was conducted in Waterloo, Ontario, at the beginning of the Canadian summer (11 June 1978). The starting time was 1 pm, when the temperature was 24.3°C, humidity 49%, wind velocity 16 knots, and cloud cover minimal. Approximately 1300 people participated in the run. While many people were treated for minor heat problems, the 15 admitted to hospital exceeded the proportion observed in the New Zealand run. Eight of the victims were kept at least overnight and two for four days. The age and ability levels varied considerably. Three were girls aged 14-18 and the 12 males ranged in age from 17 to 38 years. Eight of 12 respondents to a

survey conducted after treatment in hospital considered themselves to be runners and not joggers, although none were the front runners. This differs from the New Zealand run, in which the victims were active in sports other than running.

An attempt to discover whether there was a common cause of the heat stroke resulted in the following conclusions. Firstly, all victims had been able to cover the 10-km distance in training with no problem. However, no individual trained in the hottest part of the day. Secondly, many victims had drunk large volumes of fluid before the run and some had drunk fluid during the run, yet this did not provide adequate protection from the heat. Thirdly and most importantly, most victims had thought they could finish in spite of the relatively hot conditions. They therefore continued to push themselves, ignoring signals from the body concerning heat load. The result of this continued running was that eight people lapsed into a state of unconsciousness. The unconscious state was followed immediately by collapse in some, but at least four of the victims have no recollection of distances varying from 800 m to 2 km. This obviously can have serious life-threatening implications if exercise is continued when the heat-dissipating mechanisms are ineffective.

It is of interest that none of the experienced competitive runners were admitted to hospital and very few suffered even minor heat-related problems. The reason for this is probably twofold: the competitive runners are trained to a greater extent and they have also developed the art of listening to signals of fatigue and stress from the body.¹

It becomes apparent that stricter control must be placed on the organisation of mass participation runs because the participant may not be sufficiently aware of methods of preparation and of warning signs. Recommendations concerning the conduct of distance runs have been published before,²⁻⁵ but now the medical and sports science communities must petition the amateur athletic sanctioning bodies to enact restrictions which prohibit the scheduling of a distance run for potentially hot periods of the day. In addition, regulations governing minimal medical precautions should be established. No longer is it sufficient to count on chance to bring a cool day in the middle of the summer.

RICHARD L HUGHSON

Department of Kinesiology,
University of Waterloo,
Waterloo, Ontario

JOHN R SUTTON

Department of Medicine,
McMaster University,
Hamilton, Ontario

¹ Morgan, W P, and Pollock, M L, *Annals of the New York Academy of Sciences*, 1977, **301**, 382.

² *Medicine and Science in Sports*, 1975, **7**, vii.

³ Sutton, J R, and Harrison, H C, *Medical Journal of Australia*, 1977, **1**, 193.

⁴ Sutton, J R, et al, *Medical Journal of Australia*, 1972, **2**, 127.

⁵ *Medical Journal of Australia*, 1977, **1**, 165.

Natural history and prognosis of recurrent breast cancer

SIR,—Since the publication of our paper on the natural history and prognosis of recurrent breast cancer (9 September, p 730) we have been told by medical statisticians that there is a flaw in that part of the life-table data which we present in graphical form (figures 1, 2, and

3). This deals with the time of onset of local compared with distant recurrence and the effect of radiotherapy on the time of onset of local recurrence. The error, which we acknowledge and for which we should like to apologise, is to have excluded from the life-table analysis patients who did not have recurrence.

We should like to present amended data, dealing with these points, derived from a further analysis. Firstly, the time to onset of local recurrence is slightly longer than that to distant metastasis. The actuarial relapse rates for local recurrence at 2, 4, and 6 years are 15%, 23%, and 27% respectively, while for distant recurrence the rates are 23%, 36%, and 40%. For both groups there was a constant risk of relapse during the period of observation. Secondly, a slight, but not significant ($P=0.06$), delaying effect of radiotherapy on the time of onset of local recurrence was observed. The actuarial local recurrence rates, at 2, 4, and 6 years were 13%, 21%, and 25% for those receiving radiotherapy and 21%, 30%, and 33% for those who had not had radiotherapy.

Survival after recurrence is essentially the same as reported in our original figure 4. Five-year actuarial survival of distant and local recurrence was 5% and 32% respectively. The effects of menopausal status, age, and site of tumour on onset of local and distant recurrence and survival after recurrence are unaffected by further analysis, as is the lack of effect of prior radiotherapy on survival after recurrence.

The main conclusion of our paper is unchanged. Local recurrence is an event which implies a relatively poor prognosis. Since these patients, who have measurable disease, have presumably a smaller tumour mass than those with distant metastasis they are a group of patients in which to consider trials of cytotoxic chemotherapy for later use in adjuvant chemotherapy trials.

ROBERT SOUHAM
C G CLARK

University College Hospital,
London WC1

N J O'HIGGINS

University College,
Dublin

Surma and lead poisoning

SIR,—Dr Aulfat R. Ali and his colleagues (30 September, p 915) very elegantly demonstrate that the use of surma in Asian children can be correlated with high blood lead concentrations. After citing several other authorities in addition to their own work in Nottingham, the authors question the need for further evidence to establish beyond doubt that surma is responsible for the lead poisoning. I beg to differ.

Firstly, it is quite possible that those parents who use surma in Britain differ socially and economically from those who do not. The population using surma despite extensive propaganda may very well be less educated, more socially deprived, and reside in old houses decorated with peeling lead paint within the inner cities. Secondly, the amount of blood lead in control children is not insubstantial, though admittedly below the level considered to be harmful at present. Is it possible that surma merely acts as a precipitant to other contributory causes of excessive body lead? Thirdly, the numbers of boys among the subjects is more than twice that in the control group. Birmingham lead studies have shown

that boys have a higher blood lead level than girls.¹

I do not wish to minimise the part surma can play in precipitating lead poisoning, but the authors are wrong to assume that surma is solely responsible for lead poisoning in their study group. It is just one of the many complex causal factors—albeit uniquely important in this group—in the aetiological chain giving rise to excessive body lead levels which find their ultimate expression in clinical lead poisoning.

S S BAKHSI

Birmingham Area Health
Authority (Teaching),
Birmingham

¹ Department of the Environment, Central Unit on Environmental Pollution, *Lead Pollution in Birmingham*, Pollution Paper No 14. London, HMSO, 1978.

Beta-blockers in treatment of hypertension

SIR,—The experimental design used by Dr R G Wilcox (5 August, p 383) to evaluate seven antihypertensive drugs is such that a statistician must feel inclined to warn the physician-reader of the conclusions drawn from the resulting data in the paper. The main criticism concerns the use of a crossover design (particularly with as many as seven application periods) when the requirements for this design do not appear to be fulfilled at all.

A first requirement for any crossover design is that the state (expressed in values of the variables of interest) of all patients in the study would show a parallel course through all application periods—except for random deviations—if all patients in all periods were treated alike. With essential hypertension this requirement is probably not fulfilled.

A second requirement for a crossover is that all drugs to be compared have equal carry-over effects where equality includes the ideal case of no such effects at all, although some very special crossover designs have been proposed for the case of different carry-over or "residual" effects.¹ Since there was no wash-out between the application periods it is hard to believe that carry-over effects were not present in Dr Wilcox's study and presumably not all seven were alike.

Finally, for an unbiased comparison of the drug effects a balanced arrangement of the drug applications in so-called Latin squares is necessary, a condition obviously not met in the present study. A Latin square design requires $n=d \times k$ patients, where d is the number of drugs=number of application periods=number of patients in each square and k is the number of Latin squares. In each of the k squares each drug is applied, in a restricted randomised manner, exactly once to each of the d patients and once in each of the d application periods. In such a randomised Latin square design there is a possibility of testing for the condition of equal carry-over effects only in the case of $d=2$.²

In light of the fact that none of the three main requirements for a crossover design appear to be fulfilled in Dr Wilcox's study it seems not to be necessary to criticise the other flaws of the experiment, not to speak of those of the statistical analysis. (For example, adding another drug to the design during the course of the experiment, as done here with labetalol, represents one of the basic sins of experimental design.)

In conclusion, therefore, the resulting data from Dr Wilcox's study appear highly questionable, as do data from any crossover design when the requirements for its application are not fulfilled.

K ABT

Department of Biomathematics,
Division of Human Medicine,
University of Frankfurt am Main,
Frankfurt am Main, Germany

¹ Cochran, W G, and Cox, G M, *Experimental Designs*. New York, Wiley, 1957.

² Grizzle, J E, *Biometrics*, 1965, 21, 467.

SIR,—Dr R G Wilcox (5 August, p 383) reports on a randomised cross-over study of six beta-blockers, a diuretic, and placebo in essential hypertension. A study of this sort, designed to compare the therapeutic effect of several drugs, can be most useful. However, in such a study there is the possibility of several errors in respect of both the methodology and the statistics. Therefore the conclusions drawn by Dr Wilcox may be misleading. It is well known that the effect of a beta-blocker remains long after the drug is no longer demonstrable in the plasma. A carry-over effect may persist for several weeks, as demonstrated by Koldstrand.¹ In a comparative trial, therefore, either long enough wash-out periods with placebo have to be included or each treatment period should last correspondingly longer. In Dr Wilcox's study each period was too short and for each period the previously given treatment may have strongly influenced the result.

A possible way to avoid or at least diminish this disadvantage would be not only to randomise the order of administration but also to balance it according to a Latin square design to reduce the influence of the natural course of the disease (which is of paramount importance in a long study like this) and the carry-over effect. As eight different treatments were included in this randomised study a total of 15 patients cannot fulfil the second condition. In fact, 16 different treatment periods were included as the dose was changed each fortnight. Furthermore, there were three essentially different kinds of treatment, as beta-blockers, a combined alpha- and beta-blocker, and a diuretic were used; this calls for further caution in the interpretation of the results. All drugs in this study produced rather unsatisfactory effects on the blood pressure, which we think is mainly due to the frequent changes in therapy and the short period of only two weeks for each dose.

The author states that he used the manufacturers' recommended doses; however, he gave 5 and 15 mg of pindolol/day whereas the data sheets recommend 15–45 mg/day. As we have demonstrated,² 5 mg of pindolol is not sufficient in a once-daily dose regimen. On the other hand 100 and 200 mg of atenolol were used whereas the data sheet indicates that most patients respond to only 100 mg.

The author writes that both supine and standing blood pressures were measured, but no results of the supine measurements (which would be of great interest) are reported. The author also states that labetalol and pindolol produced twice as many side effects as the other drugs without giving the frequency and severity of any side effects or saying whether the differences between the drugs were significant.

In a study of over 700 patients treated with beta-blockers Henningsen³ found a lower frequency of side effects with pindolol