disguise of such terminology is a reasonable step—and indeed when one considers that the death certificate is required to be produced by the relatives for a variety of reasons such as legal or for burial of the deceased this would seem a humane practice.

Perhaps some euphemism or code could be introduced to cover the more emotive terms used on death certificates. This could spare the feelings of the relatives and indeed the doctor and could also contribute indirectly to more accurate death certification.

K N A MILLAR

Defence NBC Centre,
Winterbourne Gunner,
nr Salisbury, Wilts

multitude of sources of cultural knowledge and as they influence patients' everyday relation with their bodies and with the professionals who minister to them. The consequent action seems likely to involve social reform as much as health education. The first requirement, though, is a greater willingness on the part of the medical profession to keep an open mind on the apparent foolishness of its patients.

SHEILA ADAM
ROBERT DINGWALL

Department of Social and
Community Medicine,
University of Oxford

Deaths under 50

SIR,—Sir Cyril Clarke and Dr George Whitfield (14 October, p 1061) draw attention once more to the large proportion of deaths under 50 which are potentially preventable. In their sample of 250 deaths in medical wards they state that "no fewer than 98 patients contributed to their own deaths." These statistics provide a valuable illustration of one of the major problems of health care and are worthy of further discussion. However, we consider that by labelling these cases as "self-destruction" the authors fail to take account of the many stages between an individual ceasing to be healthy (as in the World Health Organisation definition) and their death. We would like to highlight two factors which are of known significance.

(1) The brief summary of their data indicates that the mechanism of "self-destruction" was cigarette smoking in 39%, gross overweight in 12%, and alcohol misuse in 7%. While it is common knowledge that these behaviours are detrimental to health, major institutions of our society not merely condone but actually encourage smoking, drinking, and the consumption of an unsuitable or excessive diet. In attributing specific "self-destructive" motives to those individuals who are "caught out" the authors fail to take account of the complexity of social and cultural influences in a modern society. The promotion of sickness for individual gain is an honoured activity. Its inevitable consequences are swept under the carpet by a philosophy of "blaming the victim."

(2) This same philosophy emerges when the authors interpret their findings by placing the responsibility for change on individual patients who are required to modify their response to an established pattern of medical care. The patients are charged with "fecklessness or a psychopathic attitude to life and to doctors in particular." This medical definition of rational behaviour dismisses any non-conforming actions as irrational, rendering any scientific attempt to investigate them pointless. The authors' assumptions obscure the degree to which patients' action or inaction may have an intelligible foundation in their experience of encounters with health services and the competing priorities in their everyday lives. Indeed, they may serve only to exacerbate the problem they identify by hardening resentment against patients who do not share medical conceptions of the good life.

This paper contributes little to our understanding of the social context of preventable death. Real advances will only come from the detailed study of lay conceptions of health and health care as they are acquired from the

Cimetidine and oral anticoagulants

SIR,—We have received preliminary details of an interaction study of cimetidine and warfarin in healthy subjects, a full report of which is to be published later. The results indicate that if cimetidine in the usual dosage of 1 g a day is given to patients receiving a stable dose of warfarin an increase in blood clotting ratio and prothrombin time of the order of 20% may be observed. The mechanism is not yet known.

Reports to this company of relevant clinical interactions have been very few (17 cases worldwide), but most of them give support to the findings of the study in that moderate rises of prothrombin time have been found and reduction of dosage of warfarin or other oral anticoagulants has been necessary. Patients receiving anticoagulants who are given concomitant cimetidine should, therefore, be carefully monitored and a reduction in anticoagulant dosage may be necessary. The text of the data sheet for cimetidine will be revised accordingly.

ANTHONY C FLIND
Medical Director,
Smith Kline and French Laboratories Ltd
Welwyn Garden City, Herts

Comparison of the tine and Mantoux tuberculin tests

SIR,—We have been particularly interested by Drs J A Lunn and A J Johnson's report (3 June, p 1451) and the subsequent correspondence on the diagnostic value of the tine test. We are in an unusual position in this district, as it was not the policy of the former local authority to undertake BCG vaccination. As a result we have a number of girls applying to be nurses at the Friarage Hospital who are tuberculin negative and who subsequently are given BCG vaccination on appointment. BCG vaccination where the tine test has been absolutely negative is frequently followed by conversion to a small but palpable response on repetition. As a result of this experience we were satisfied that the false-negative rate for tine testing is low and that any palpable response should be regarded as evidence of tuberculin sensitivity. However, we strongly suspected that the false-negative rate was substantial, though we were surprised that Dr Lunn reported it at such a high level.

We therefore studied 81 nurses, all of whom were reported either as previously tuberculin positive or as having had BCG vaccination. They were each given a test on each arm. Two batches bought commercially were used, but the same batch was used for each individual. Our results are summarised in the table, positive being taken as any palpable response.

We did attempt to grade the responses in a more detailed breakdown and this is available on request. If the left arm is taken as the test arm and the right as the check arm the false-negative rate lies between 13% and 28%, allowance being made for the occurrence of double false-negatives due to change and variation in individual response.

Percentage of 81 nurses (presumed tuberculin positive) giving positive and negative responses to tine test

	Left arm:	
	Negative	Positive
Right arm:		
Negative	18	8
Positive	10	63

This high false-negative rate makes it essential that a repeat tuberculin test is done on all patients who show a negative tine reaction. This makes the test quite unsuitable for use epidemiologically, particularly in populations where a high negative rate is anticipated. It does not necessarily preclude its use, if it is particularly convenient, in populations where a high positive rate is anticipated, provided it is appreciated by the operator that the false-negative rate is of the order of 20%. No action should be taken on the basis of a negative tine test until this has been confirmed by formal Mantoux testing.

C K CONNOLLY
T NAIK
E SHUTTLEWORTH

Friarage Hospital,
Northallerton,
North Yorks

SIR,—Drs J A Lunn and A J Johnson (3 June, p 1451) report an astonishing difference between the results of tine and Mantoux tests simultaneously performed on 307 mainly young subjects. They found only 3.9% positive tine tests among the 59.0% Mantoux-positive persons. We have the impression of a much better correlation between these two tests.

We studied 100 consecutive patients (59 men, 41 women, mean age 66.9 years) with tine tests on the left and Mantoux tests on the right forearm, using the Lederle tine test (expiry date 7 October 1979) and 5 units of tuberculin PPD (Berna). All tests were performed by two specially trained nurses. The results, read independently by two physicians, are shown in the table, which corresponds to table I in the study of Drs Lunn and Johnson. According to the manufacturer's recommendation tine tests with two of the four papules greater than 2 mm in diameter should be read as positive, whereas the American Lung Association¹ interprets a multiple-puncture test as positive only if vesiculation is present. For the sake of com-

Comparison of tine and Mantoux tuberculin tests on 100 consecutive subjects

Reading of Mantoux test	Reading of tine test			Total
	Negative (0-1 mm)	Doubtful (2-4 mm)	Positive (≥5 mm)	
Negative: (0-4 mm)	35	2	9	46
Positive: (5-9 mm)	—	5	6	11
(≥10 mm)	—	5	38	49
Total	35	12	53	100