Instead has been widely distributed to householders with the rate demands and well received.

Our health education free booklet *Clean Food and Clean Air*, sponsored by private advertisers, was distributed from 1971 onwards. An attack was made on stubble burning and some progress made by limiting this self-indulgence by those involved in the rural revolution and industrial pollution of the countryside. Environmentalists, the fire services, and the National Farmers' Union have helped to curb the more pyromanic farmers by introducing a code of practice, which is regrettably seldom followed.

Letters in the local press have exhorted gardeners at this time of year not to squander leaf fall by burning it and sickening their neighbours. Noise and smoke are the commonest causes of neighbourly disaffection and source of public complaint to the health authorities. Recently reports have been received from a rurally situated hospital where staff and patients were discomfited by stubble burning and smuts contaminating patients' food. Unfortunately our current district bylaw, which dates from 1974, is effective only against fire spreading, not smoke and ash spreading.

The civic amenity collections service has helped to reduce the problem of individuals burning furniture and other potentially lethal plastic materials which, if allowed to smoulder, can produce noxious gases even in the open. We hope that the civic conscience of the population will be alerted to the health and safety hazards of incomplete combustion epitomised by bonfires. Perhaps they should be called *malfires* instead. We have nothing against a good blaze consuming its own smoke.

In our published material we have drawn attention to the risk of small valleys filling up with smoke, especially at night, when atmospheric layers of air become "inverted." Could the weather centre and the BBC weather forecasters not be requested to give warnings when these conditions are likely to occur? They might suggest when stubble burning and bonfires are likely to lead to the outdoor air being fouled in this way, and when they ought to be discouraged. We should be interested in their response and so no doubt would bronchitic patients and non-smokers. A similar service is offered to pollen sufferers during the summer months.

BRIAN H BURNE
Medical Officer for Environmental Health

NOEL J BILLINGTON
Chief Environmental Health Officer

Chiltern District Council,
Chesham, Bucks

Postgraduate education in therapeutics

SIR,—The description of the Liverpool Therapeutics Group by Professor A Breckenridge and his colleagues (2 September, p 671) is an interesting and important contribution to the current debate on continuing education in clinical pharmacology. However, we would wish to raise two issues.

Firstly, the comment that community clinics in clinical pharmacology "... undoubtedly benefit the participant practitioners greatly, but have limited application," is misleading. Community or general practice clinics have been shown to have applicability in a variety of differing specialties. By combining clinical care of patients with continuing

education of doctors they fulfil a dual role. Conducted in group practice premises, they also have the advantage of involving all the partners in a practice, not only those who would have been attracted to a more conventional postgraduate course. They undoubtedly do benefit the participating practitioners, and we have shown that the number of practitioners need not be restricted¹; a single consultant could relate to 100 doctors serving a population of 240 000 with relative ease. The major limitation on this type of development is the reluctance of consultants to move outside the perimeter of the hospital walls.

Our second point is to express surprise that such a distinguished group of clinical pharmacologists doubted their ability to assess whether their teaching sessions affected patient management. Although outcome measures are notoriously difficult, a convenient index of the effect of education in this subject is the practitioner's prescription. To experiment without any form of measurement is rather like mounting a clinical trial which defines the drug, the dose, and the duration of treatment but does not study what happens to the patient.

ANDREW SMITH
JOHN H WALKER
MICHAEL D RAWLINS

Department of Family and Community
Medicine,
The Medical School,
Newcastle upon Tyne

¹ Smith, A, Walker, J H, and Rawlins, M D, *British Medical Journal*, 1977, 2, 169.

Carcinoma of the breast in women under the age of 30

SIR,—We read with considerable interest the letter from Mr A S Purandare and Dr J A H Finbow (9 September, p 771) on the incidence of breast cancer under the age of 30. We have recently assessed the age distribution of patients with breast cancer referred to this centre since 1955. Our findings are shown in the table. Women with non-carcinomatous malignancies (for example, lymphoma or sarcoma) together with men have been excluded. The total number of women with carcinoma and the age distribution expressed as a percentage for each of the five years since 1955 are shown in the table below.

These figures indicate that we have not seen a marked change in the age incidence during this 20-year period. The number of women under the age of 30 presenting with breast cancer has remained at 1.1% or less. We feel that the difference between Purandare and Finbow's figures of 5.1% over a two-year period and our own of 0.3-1.1% at period intervals of five years covering a total of 20 years may be due to the fact that we are a specialised unit, but this factor alone would not account for the difference in our figures. We would like to point out that the area we serve covers all types of communities.

It is interesting that our figures are similar

to those of Haagensen, as quoted by Purandare and Finbow. We therefore conclude that there has been no increase in breast cancer among women under the age of 30 years since 1955.

VERA JONES
ALAN SLATER

Radiotherapy Centre,
Velindre Hospital,
Cardiff

How to organise an international medical meeting

SIR,—I too have read with admiration the series of articles by Mr Ian Capper and Mr A I S Macpherson on organising international meetings. While sharing some of Professor John Dobbing's (16 September, p 827) doubts about the usefulness of enormous gatherings I think he goes to the other extreme in suggesting that no more than 20 people should ever be brought together for the presentation of papers and an exchange of views. One point he makes, however, and to this Mr Capper and Mr Macpherson give scant attention, is that the published proceedings of a well-planned medical meeting are liable to reach a very much wider audience than those present on the day. Thus in the long term they are potentially more important than the (at least in Professor Dobbing's view) agonising ephemera of the actual congress.

However, the resulting book or journal supplement is unlikely to be of the highest standard if the organisers merely tape-record the papers; having been briefed well in advance by the publishers, contributors should be asked to hand in manuscripts, if possible, on the day of the meeting. Publication of the proceedings, including the relevant discussion, will, contrary to Professor Dobbing's belief, need skilled editing; those who have occasion to search the medical literature, and know the value of having a body of knowledge on a particular topic conveniently collected in this form, will understand that that is well worth while.

LYN WILSON
Editor,
Proceedings Publications, ADIS Press

Auckland,
New Zealand

Mycosis fungoides treated by PUVA and mustine hydrochloride

SIR,—We were interested in the observation by Dr G Volden and others (23 September, p 865) that contact dermatitis to mustine hydrochloride diminished after oral psoralen plus long-wave ultraviolet light (PUVA) therapy in five cases of mycosis fungoides. Thinking along similar lines, we have started a pilot study of treating mycosis fungoides with topically applied mustine hydrochloride and PUVA therapy simultaneously. We argued that since PUVA therapy has been shown to inhibit

Percentage of women with carcinoma of the breast in different age groups

Year	No of patients	Age						
		<30	30-40	40-49	50-59	60-69	70-79	80+
1955	274	1.1	6.6	18.6	32.8	27.0	12.4	1.5
1960	312	1.0	7.4	19.9	23.1	25.3	18.6	4.8
1965	400	0.5	6.5	22.5	26.0	26.8	13.8	4.0
1970	304	0.7	4.9	19.1	30.0	23.0	15.1	7.2
1975	364	0.3	6.3	18.4	25.3	26.6	19.2	3.8