recorded 62 of their 100 patients and 35 of their 100 controls would have been classified as regular cornflake eaters ($\chi^2=14.6$, $P<0.001$).

They have thus shown that 53% of regular eaters of cornflakes abandoned the habit when they developed Crohn's disease, bearing out my comments (9 September, p 767) on the findings of Drs L N J Archer and R E Harvey (19 August, p 540). I accept that recollection may be faulty, but the authors have recorded these data and the inference is there to be drawn.

A H JAMES

Hillingdon Hospital,
Uxbridge, Middx

¹ James, A H, *British Medical Journal*, 1977, 1, 943.

Inadequacy of information on side effects

SIR,—At this hospital three patients have developed haemolytic anaemia due to phenazopyridine (Pyridium) in the past five years. They all showed exactly the same haematological features as the patient described by Drs H C Drysdale and M D Hellier (7 October, p 1021) and in all three the haemolysis ceased when the drug was stopped.

With this experience in one small health district, I find it difficult to believe that this particular complication of Pyridium therapy is as rare as the literature and Dr Witherspoon (p 1021) suggest. It should certainly not be classed as a "rather rare suspect event."

C J T BATEMAN

Royal West Sussex Hospital,
St Richard's,
Chichester, W Sussex

Computer confidentiality

SIR,—The second of the three ethical principles which the Child Health Computing Committee has put forward to govern computing systems (11 November, p 1382) does not protect the right of the patient's parents to control the distribution of medical information about their child. I consider the following addition (in italics) to be essential:

(2) "Access to identifiable information held in medical records is to be confined to the author and to the person clinically responsible for the patient during the episode for which the data has been collected (or their successors) unless specifically authorised by the clinician in the clinical interest of the patient and *by the patient (or his or her parent or guardian).*"

I would have thought that this point was elementary and am very surprised by the apparently widespread British practice of sending copies of children's medical reports to school medical authorities without the knowledge and consent of the parents. Such behaviour, irrespective of its motives, should be made clearly illegal. It cannot be excused by saying that it is "well meant" or "for the ultimate good of the patient"; it is for the patient (or his parents) to decide what is for his ultimate good and it is for him to decide whether a copy of a medical report should be sent to the school medical authorities, Uncle Tom Cobley, or anyone else. As soon as patients learn that a "mailing list" (of which they have no knowledge) may be appended to any information which they divulge in the consulting room they will learn to be selective when talking to their doctors. And that, as you

probably realise, would be the beginning of the end of the doctor/patient relationship which people often enthuse about.

PATRICK A CASEY

Wohlen, Switzerland

Scombrototoxic fish poisoning?

SIR,—I recently observed a reaction which occurred 30 minutes after eating a Bombay duck, which is a type of dried fish. The patient developed headache and nausea concurrently with a flushing of the face, chest, and shoulders, and the lips became swollen. The redness of the face and upper trunk gave way to a blotchy urticarial pattern with weals after a further five minutes. An intramuscular injection of chlorpheniramine maleate 10 mg was followed by a dramatic improvement and one hour later the patient was completely recovered and had no further relapses.

I am not sure whether this reaction was due to scombrototoxic fish poisoning (9 September, p 739) or to simple allergy, but the patient has no allergic history and has eaten Bombay duck since with no ill effects.

A J WARING

Limavady, Co Londonderry

Intrauterine hiccup

SIR,—Having just returned from five years of North American investigative medicine I find it delightful to read that the diagnosis of intrauterine hiccup is yet another use for real-time ultrasound scanning (Dr I Swann, 25 November, p 1497).

Most senior house officers who deal with postnatal patients would confirm our own clinical impression that intrauterine hiccups occur frequently (approximately one per 50 live births). They are noted by the end of the second or beginning of the third trimester and recur in subsequent pregnancies. They frequently occur at the same time of day for each patient and persist after birth. All the babies that we have seen have been normal. Most of our mothers correctly diagnosed themselves antenatally. This was usually confirmed by their midwife and so far there has been no need for a scan.

J DEARLOVE

Department of Paediatrics,
Charing Cross Hospital,
London W6

Conservation surgery for laryngeal cancer

SIR,—Your leading article (11 November, p 1318) on this subject was timely, factual, and well informed. In the last paragraph you ask whether British laryngologists are perhaps unduly hesitant to use these techniques—presumably supraglottic partial laryngectomy—for irradiation failures. I therefore write to assure you that our unit at this hospital is devoting time and study to this problem.

Patients with suitable lesions are uncommon and assessment of the extent of disease after irradiation so difficult that a firm decision for a partial operation can be made only on the operating table. In my experience no more than a small minority prove to be treatable and curable in this way, the majority requiring total laryngectomy. Nevertheless, such efforts

are rewarding and further salvage may be possible by total resection if recurrence takes place. It must also be said, with emphasis, that all patients with laryngeal cancer at any site must be observed before and regularly throughout irradiation treatment by the radiotherapist and the surgeon, who will operate if treatment fails. Only in this way can the surgeon hope to gain an approximate idea of the extent of disease. It is also my belief that by ignoring the curative value of treatment by initial supraglottic laryngectomy as now demonstrated in many centres outside Britain we are condemning more patients to eventual total laryngectomy if irradiation fails.

H J SHAW

Head and Neck Unit,
Royal Marsden Hospital,
London SW3

Nutrition and the cancer patient

SIR,—Mr A J Strain and his colleagues (4 November, p 1295) discuss the unavailability of appropriate animal models for the study of cancer-associated cachexia. I would certainly agree that the readily available rapidly growing transplantable tumours of laboratory rodents do not provide realistic models of cachexia in that significant loss of body weight does not usually ensue until tumours approach 40% of the body weight. Large tumours, especially when ulcerated, are commonly associated with extreme anaemia; the coincident weakness may itself discourage animals from eating and so give rise to a constitutional state indistinguishable from starvation.

In my experience of over 30 mouse tumours which have been maintained by serial transplantation for prolonged periods wasting is rarely seen in mice bearing tumours of up to 3 g ($\approx 10\%$ of body weight). The exceptions have been three squamous carcinomas (one arising in skin and two in the fore-stomach); all have been associated with extreme wasting at relatively small tumour sizes. In a comparative study of one of these tumours with a mouse mammary carcinoma (MMC) growing at about the same rate we observed that the squamous tumours when grown to a mean weight of only 1.1 ± 0.8 g were associated with a mean loss of body weight of 16% (range 0-30%). In contrast no loss of body weight was associated with growth of the MMCs to a mean weight of 4.6 ± 2.3 g. The wasted mice bearing squamous carcinomas rapidly became moribund and showed a depression of blood glucose concentration from the normal 8 mmol/l (145 mg/100 ml) to between 5.6 and 4.4 mmol/l (100 and 80 mg/100 ml) (as seen in mice starved for 24 h). Wasted mice did not appear to reduce their food intake (as measured by reduction in the weight of the food pellets to which they had access), but we did have some evidence that the food taken was not being masticated or swallowed. Since one of our squamous carcinomas has been shown to produce epidermal chalone¹ we entertained a hypothesis that such chalone inhibited mitoses in the normal squamous epithelia of tumour bearers and that this could reduce cell replacement in squamous mucosa and so give rise to painful mastication and swallowing. In support of this hypothesis was our finding that the mitotic rate in squamous epithelia (tongue, oesophagus, and fore-stomach) of wasting mice was reduced to between 50% and 20% of normal. However, we could not establish whether the mitotic inhibition preceded or followed the wasting.

It is of interest that the diet given to mice which exhibited cachexia was a standard formulation and fully supported weight output of weanlings. Wasting was not seen when the diet was changed to one with a higher fat content.

The point to be made here is that distinctive