The pattern of lower referral rates for singlehanded practices was present in both urban and rural practices. This contrasts directly with the findings of a study on the use of laboratory facilities by general practitioners in the north-east of Scotland¹⁰ which showed that single-handed rural practitioners were the highest users of laboratory services.

We talked to 10 general practitioners chosen from those with high and those with low referral rates. Those with low referral rates claimed to refer correspondingly more patients to the outpatient departments, and the rural practitioners adopted the policy that if the patient is going to have to travel to Aberdeen anyway then it is just as well to refer him to an outpatient clinic. It was suggested that in Aberdeen city the lower referral rate of singlehanded general practitioners might be due to a tendency for them to be older than general practitioners in group practices. This was not shown, but the suggestion is supported by the findings of Morrison and Riley¹¹ and Davis and Williams,⁵ who showed that the less frequent users of direct access diagnostic facilities were older doctors.

We thank Professor I. M. Richardson of the department of general practice for his advice and the following general practitioners who gave up their time and co-operated in this survey: Dr. J. Chalmers, Inch; Dr. C. I. T. Cruickshank, Inverurie; Dr. C. L. Forbes, Aberdeen; Dr. J. Gauld, Aberdeen; Dr. F. F. Mair, Banchory; Dr. I. Ross, Aberdeen; Dr. G. G. Shirriffs, Aberdeen; Dr. G. Sorrie, Aberdeen; Dr. R. F. F. Steven, Aberdeen; Dr. A. V. R. Watson, Torphins.

References

- ¹ Scottish Home and Health Department, *Doctors in an Integrated Health Service*. Edinburgh, 1972. H.M.S.O.
- ² Whitfield, M. J., *Practitioner*, 1973, 210, 780.
- ³ Steiner, R. E., *Proceedings of the Royal Society of Medicine*, 1965, 58, 448.
- ⁴ Cook, P. L., *British Medical Journal*, 1966, 2, 351.
- ⁵ Davis, R. H., and Williams, J. E., *British Medical Journal*, 1968, 1, 502.
- ⁶ Petrie, J. C., Needham, C. D., and Gillanders, L. A., *British Medical Journal*, 1972, 1, 78.
- ⁷ Lennon, E. A., *Health Trends*, 1969, 1, 17.
- ⁸ Wallace, B. B., et al., *Journal of the Royal College of General Practitioners*, 23, 337.
- ⁹ Fry, J., et al., *Medical World (London)*, 101, 23.
- ¹⁰ Brodie, J., and Porter, I. A., *Health Bulletin*, 1972, 31, 244.
- ¹¹ Morrison, S. L., and Riley, M. M., *Medical Care*, 1963, 1, 137.

Hospital Topics

Treatment of Urinary Retention

J. J. SHIPMAN, B. M. NEWMAN, G. MORRIS, K. A. MISCH

British Medical Journal, 1974, 3, 734-735

Summary

In 50 patients retention of urine due to adenomatous enlargement of the prostate was relieved, in most cases by intraprostatic injections given for varying periods of time. No complications were encountered. Subsequently, some of the patients who did not respond adequately, and whose medical conditions improved, underwent prostatectomy. Histologically the removed prostates showed coagulation necrosis, with or without haemorrhage, but no inflammatory reaction.

Introduction

Interest has continued in the treatment of acute retention of urine by intraprostatic injection treatment since the successful relief of retention in 17 patients illustrated by Shipman and Akilie.¹ This paper describes a four-year follow-up of 50 patients treated for acute retention of urine due to an adenomatous enlargement of the prostate by the intraprostatic injection of a solution containing phenol, glacial acetic acid, and glycerin.²

Technique and Selection of Cases

The prostate was injected during the first three days in hospital, as follows. The patient was placed in the left lateral position and the prostate injected with 2 to 3 ml of a solution of the following ingredients: phenol 0.6 ml; glacial acetic acid 0.6 ml; glycerin 1.2 ml; and distilled water to 28 g. A 20-gauge lumbar puncture needle was used.

A preliminary injection of local anaesthetic was made in only one of our patients. The injections were not painful. A finger placed in the rectum acted as a guide. The needle point required repositioning if pain was felt down the penis or blood could be aspirated into the syringe.

FIG. 1—Seven days after prostatic injection. Low power microphotograph of prostate showing an area of fresh necrosis and adjacent haemorrhages. (H. & E. $\times 25$.)

Department of Surgery and Pathology, Lister Hospital, Stevenage
J. J. SHIPMAN, M.S., F.R.C.S., Consultant Surgeon
B. M. NEWMAN, F.R.C.S., Surgical Registrar
G. MORRIS, M.B., F.R.C.S., Surgical Registrar
K. A. MISCH, D.PATH., F.R.C.PATH., Consultant in Pathology

FIG. 2—Multiple unsuccessful injections between ten to two weeks before prostatectomy. Edge of necrotic area with squamous metaplasia. (H. & E. $\times 100$.)

FIG. 3—Successful prostatic injections one year before prostatectomy. Low power microphotograph of fibrosed infarcted area with squamous metaplasia of glands along edge. (H. & E. $\times 25$.)

FIG. 4—As FIG. 3, higher power to show metaplastic glands at edge of infarct. (H. & E. $\times 400$.)

We have used this form of treatment in patients who were poor operative risks owing to intermittent chest disease or cardiovascular disease and who presented with acute urinary retention or acute-on-chronic retention, and failed to pass water after catheter trials. No side effects were encountered in any of our patients. Patients have complained of no more discomfort than would be expected with any injection in this region. The contraindications to injections were causes other than benign enlargement, associated vesical calculus, and cases in which a catheter could not be passed.

A total of 189 patients were admitted with retention. In 45 of these there was a cause other than benign enlargement, and 94 passed urine spontaneously after 48 hours trial, or had a prostatectomy. The remaining 50 received the injection

treatment. The average age of this group was 73.4 years.

Each patient had an average of 3.2 injections for the relief of symptoms. No complications were associated with injections of the prostate, there was remarkably little discomfort, and no analgesics were necessary afterwards.

Follow-up Study

In view of the old age of the patients in this series, and their poor general medical condition, only 29 were still alive in the follow-up period. The causes of death were mainly cardiovascular and respiratory complications; none was attributed to the treatment or renal impairment. Of the 29 survivors, 13 had a normal stream for three years or more, 12 had a normal stream for one to three years, and four from two months to a year.

Pathology

We examined 22 prostates, which had been injected at varying periods (one week to five years) before transvesical prostatectomy, by serial slicing and histological examination. We excluded three cases as they were found to be carcinomatous. Almost all of the prostates showed macroscopic evidence of trauma varying from haemorrhages to gross necrosis with infarction. In cases which had been injected over six months or more previously, there were also fibrous scars. The lesions were often multiple and were found in any part of the prostate. The largest was 3 cm in diameter occupying almost a whole lateral lobe. Microscopically, there was coagulation necrosis, with or without haemorrhages (fig. 1). There was a striking absence of any inflammatory reaction, but the necrotic areas were often surrounded by metaplastic squamous epithelium, though this depended on the age of the lesion (fig. 2). Old infarctions showed as fibrous areas with complete loss of muscle and were invariably surrounded by squamous metaplasia of the adjacent glands (figs. 3 and 4). Thromboses were rare and were seen only in one case. The lesions found differ in no way from the infarcts occasionally found during routine examination of adenomatous prostates, first described by Abeshouse³ and later by Moore.⁴

Discussion

The patients were chosen initially for treatment because they could not be operated on owing to their poor general condition. Thus the intraprostatic injection treatment has opened up another technique in the management of the adenomatous prostate—saving the patient from a catheter life. The mechanism of the relief is unknown. Hubly and Thompson⁵ suggested that contracture of a healing prostatic infarct could relieve urinary obstruction by mechanically reducing the prostatic volume. This is unlikely to be the explanation in our cases, as the infarcts produced by the injections were almost always very small and relief occurred within a few hours of the injection.

The rational fear of the prostatic abscesses or urethral stricture appears to be unfounded with the technique used. It may be necessary to give three or more injections before a satisfactory result is obtained, the average number of injections given—3.2—corresponding well with the findings of Talwar and Pande.² We think that this form of treatment is useful in selected cases of benign glandular enlargement presenting with acute retention.

References

- 1 Shipman, J. J., and Akilie, A. N., *British Medical Journal*, 1967, 2, 418.
- 2 Talwar, G. L., and Pande, S. K., *British Journal of Surgery*, 1966, 53, 421.
- 3 Abeshouse, B. S., *Journal of Urology*, 1933, 30, 97.
- 4 Moore, R. A., *Journal of Urology*, 1943, 50, 680.
- 5 Hubly, J. W., and Thompson, G. J., *Journal of Urology*, 1940, 43, 459.