bmj.com news roundup

Full versions of these stories are available at: bmj.com/content/vol333/issue7574/#NEWS_ROUNDUP

Doctors are warned against having sex with former patients

Pursuing a sexual relationship with any former patient will "usually be inappropriate, regardless of the length of time since the therapeutic relationship ended," warns supplementary guidance on maintaining boundaries, issued by the General Medical Council alongside its latest updated guidance on acceptable professional practice.

Doctors must not have sexual relationships with vulnerable former patients, such as those with mental health problems or insufficient maturity, it states.

In these circumstances it may be difficult to be certain that the professional relationship is not being abused, it says. Doctors who are unsure of their position should seek advice from an impartial colleague, professional body, or the GMC.

The main change to the new guidance—last updated in 2001—is the need to take account of patients' views in assessing their condition and to support them in self care.

Other new recommendations include guidance on equality and diversity and on doctors' duty to report to the GMC any convictions or findings against them by any professional body.
Lynn Eaton *London*

Good Medical Practice is available at www.gmc-uk.org/guidance.

Incidence of breast cancer is rising in Australia, while death rate falls

The annual number of new cases of breast cancer has more than doubled in Australia over the past 20 years, says a report from the National Breast Cancer Centre and the Australian Institute of Health and Welfare. A total of 12 027 women were given a diagnosis of the disease in 2002, up from 5318 in 1983.

Women now have a one in eight chance of receiving a diag-

Screening with computed tomography detects early lung cancer

Screening with spiral computed tomography (CT) in people aged 40 or more who are at risk of lung cancer detects early cancers, giving a 10 year survival rate of 80% to 92%, shows new research.

The international early lung cancer action project (I-ELCAP) screened 31 567 asymptomatic people who were current or previous smokers, or who had been exposed to secondhand smoke, or potential carcinogens at work.

The baseline and annual screenings detected lung cancer in 484 participants. Eighty five per cent (412) had stage I lung cancer, and their estimated 10 year survival rate with surgery or other treatments was 88% (95% confidence interval 84% to 91%). The survival rate at 10 years of the 302 people with stage I cancer who had surgical resection within a month of diagnosis was 92% (88% to 95%).

Screening is "highly cost-effective... similar to that for mammography," say Claudia Henschke of New York Presbyterian Hospital-Weill Cornell Medical College and colleagues (*New England Journal of Medicine* 2006;355:1763-71).

Janice Hopkins Tanne *New York*

nosis of breast cancer by the age of 85, up from one in 12 in 1983. The rise was mostly seen among those aged 50-69.

But a woman's risk of dying from breast cancer before the age of 85 years fell from one in 29 in 1983 to one in 36 in 2004. The death rate in Australia was lower than that in New Zealand, the United Kingdom, and North America.

Women in high socio-economic groups had the greatest risk of developing breast cancer. But the disease was the most common cancer among Aboriginal and Torres Strait Islander women, despite their having a lower incidence than non-indigenous people.

Stephen Pincock *Sydney*

Breast Cancer in Australia: An Overview, 2006 is available at www.nbcc.org.au.

NICE issues new guidance on incontinence in women

New guidance for doctors in England and Wales aims to improve the care and treatment of millions of women who have urinary incontinence.

But surgeons should have specific competences before they perform surgery in cases of urinary incontinence, says the guidance, which was issued this week by the National Institute for Health and Clinical Excellence (NICE).

Women should also be categorised as having either stress urinary, mixed, or urge incontinence (overactive bladder syndrome).

The guidelines contain around 100 recommendations

and advise on diagnosis, treatment, and training, in an area still regarded by many as a "hidden" subject, says the institute.

Treatment recommendations include:

- Lifestyle changes, such as changing how much a person drinks and losing weight
- At least three months of pelvic floor muscle training
- At least six weeks of bladder training.

Experts believe that the condition affects five million women in England and Wales aged over 20 and one in three women aged over 40.

Adrian O'Dowd *London*

The guidance is available at www.nice.org.uk/CG40.

NHS launches library for research into ethnicity and health

The NHS in England and Wales has launched a free electronic library to bring together quality assured medical research about different ethnic groups and cultures, in a bid to develop good practice and support the investigation of health inequalities in the United Kingdom.

Created by researchers at Warwick Medical School and De Montfort University in Leicester, the web based library will collect and index relevant evidence, and allow NHS staff and community and voluntary organisations to see what multilingual information is available for patients.

Muir Gray, director of clinical knowledge, process, and safety for Connecting for Health—which oversees the implementation of the NHS's national programme for information technology—said that the web based library would highlight substantial gaps in the knowledge of how treatments affect specific ethnic minorities.

Resources on the site are listed under four main headings: diseases and conditions, service delivery, cultural competence, and management and policy.

Robert Short *London*

The resource can be accessed at www.library.nhs.uk/ethnicity.