group as a whole ($P < 0.025$). The antipyrine half life of the 20 non-deficient people was 11.3 ± 2.1 h, which was significantly shorter than in either the group as a whole ($P < 0.05$) or the 16 people with low vitamin levels ($P < 0.001$).

We conclude that poor nutrition in the elderly is at least in part responsible for the decline in microsomal enzyme function with age.

D J SMITHARD
M J S LANGMAN

University Department of
Therapeutics,
City Hospital,
Nottingham

national interests at heart and by those responsible for the public health.

If the Royal College of Physicians and other medical authorities take up the challenge as they have done effectively in regard to smoking much can be accomplished to control the incidence of alcoholism and to instruct the public in the facts of this increasing major national health problem.

J E CAUGHEY

National Iranian Oil Company,
Non-basic Operations Abadan
and Fields—Khuzistan,
Iran

A J ISAACS
D LESLIE
J GOMEZ
P R RAGGATT

Westminster Hospital,
London SW1

¹ O'Malley, K, *et al*, *British Medical Journal*, 1971, **3**, 607.

² Vestal, R E, *et al*, *Clinical Pharmacology and Therapeutics*, 1975, **18**, 425.

³ Briant, R H, *et al*, *New Zealand Medical Journal*, 1975, **82**, 136.

⁴ Griffiths, L L, *et al*, *Gerontologia Clinica*, 1967, **9**, 1.

⁵ Smithard, D J, and Langman, M J S, *British Journal of Clinical Pharmacology*. In press.

⁶ McLennan, W J, *et al*, *Quarterly Journal of Medicine*, 1973, **42**, 1.

⁷ Baylis, E M, *et al*, *Lancet*, 1971, **1**, 62.

⁸ Colby, H D, *et al*, *Biochemical Pharmacology*, 1975, **24**, 1644.

The cowardice continues

SIR,—Under the above title (4 June, p 1428) you berate the Government for its lack of effective action against the lethal effects of smoking. I am in complete agreement with the tenor of your article. However, a further reaction was to ask myself when we as a profession are going to overcome our cowardice and take up the cudgels against alcoholism as we have done quite effectively against smoking.

Having worked among alcoholics in New Zealand in the past and been president of the National Society on Alcoholism for seven years I have had the opportunity to study the many facets of the problem as it affects the individual and his family, the national economy, the cost to medical services, and the effect on crime and accident rates. Competent surveys initiated by the society and the public health authorities give some impression of the extent of the problem in New Zealand—a small country with approximately 3 million inhabitants. In a Sir Charles Burns lecture Dr Fraser McDonald, a leading psychiatrist, gave a small sample of the statistics as follows. (1) Thirty per cent of all male admissions to both general hospital, psychiatric units, and mental hospitals are due to alcoholism. (2) It is estimated that 30% of acute surgical beds in general hospital are occupied by patients in whom a major contributory factor is alcohol. A study in Christchurch Hospital showed that 70% of road accident victims would not have been able to pass the breath test. (3) One out of 15 adult social drinkers will become an alcoholic. (4) It is estimated that there are 35 000 alcoholics in New Zealand and some 200 000 people will have their lives severely disrupted by alcoholism.

These statistics indicate some distressing personal aspects of the problem but do not touch on the many medical aspects of the problem, the part alcohol plays in the increasing crime rate, and the impact on the economy of the country in lost man hours and loss of efficiency. Clearly alcoholism is a great national health problem and the adverse effects of social drinking and of alcoholism on such a large section of the community cannot be continually ignored by those of us who have the

Prolactin in anorexia nervosa

SIR,—Dr A D B Harrower and his colleagues (16 July, p 156) report finding raised serum prolactin concentrations in anorexia nervosa. Our own results, to be published in detail shortly, are in marked contrast. We examined basal serum prolactin concentrations and the response to thyrotrophin-releasing hormone (TRH) in 11 female patients with anorexia nervosa, both in the acute stage and after supervised weight gain in hospital. No psychotropic medication was given.

The basal prolactin level (mean $\pm$ S.D.) before weight gain was 4.8 ± 2.5 μ g/l, and after weight gain 6.6 ± 3.1 μ g/l. These results do not differ significantly from each other nor from the mean level in a group of 23 healthy female controls (6.3 ± 2.1 μ g/l). The ranges of prolactin concentrations were: before weight gain 2–11 μ g/l; after weight gain 3–13 μ g/l; healthy controls 2–12 μ g/l. There was thus no evidence of hyperprolactinaemia in the anorexic patients—if anything the results were marginally lower before weight gain.

When an intravenous injection of 200 μ g of TRH was given before weight gain the mean prolactin level rose to 20.5 ± 7.7 μ g/l after 20 min and was 16.2 ± 5.4 μ g/l after 60 min. After weight gain the corresponding values were 35.5 ± 11.0 and 21.5 ± 6.7 μ g/l. The difference between the 20-min levels is highly significant ($P = 0.001$), while there is no significant difference between the 60-min levels. These results are consistent with the delay in peak prolactin secretion after TRH in anorexia nervosa shown by Vigersky *et al*¹ and the recovery in pituitary prolactin reserve after weight gain is similar to that demonstrated for gonadotrophins.²

On examination of Dr Harrower and his colleagues' data it is clear that the apparent hyperprolactinaemia of their group with anorexia nervosa is really due to the effect of the one subject with a considerably elevated serum prolactin concentration of 90 μ g/l, which requires investigation in its own right and rigorous exclusion of the possibility of drug administration. If this subject is excluded the mean serum prolactin concentration falls to 16.6 μ g/l, not significantly different from that of their obese group. Similarly, the report of Travaglini *et al*³ of hyperprolactinaemia in five out of six anorexic patients is unsatisfactory in that therapy with dopamine antagonists was not excluded.

We would therefore conclude that there is no unequivocal evidence of hyperprolactinaemia in anorexia nervosa. This is important to resolve as it may help to elucidate the central mechanisms involved. Both depletion of brain dopamine⁴ and increased activity of dopamine at central receptors⁵ have been postulated as mechanisms for the development

of anorexia nervosa. The evidence from prolactin studies is against the dopamine depletion hypothesis. Most reports suggest normal basal prolactin levels in anorexia nervosa, but our finding of impaired pituitary prolactin reserve and the study of Kalucy *et al*⁶ showing decreased nocturnal prolactin secretion are more consistent with the second hypothesis.

¹ Vigersky, R A, *et al*, *Clinics in Endocrinology and Metabolism*, 1976, **5**, 517.

² Warren, M P, *et al*, *Journal of Clinical Endocrinology and Metabolism*, 1974, **40**, 601.

³ Travaglini, P, *et al*, *Acta Endocrinologica*, 1976, **81**, 252.

⁴ Mawson, A R, *Psychological Medicine*, 1974, **4**, 289.

⁵ Barry, V C, and Klawans, H L, *Journal of Neural Transmission*, 1976, **38**, 107.

⁶ Kalucy, R S, *et al*, *Journal of Psychosomatic Research*, 1976, **20**, 595.

Starting on the pill

SIR,—Women starting on oral contraceptives are advised to take the first pill on the fifth day of their menstrual period and told that additional contraceptive precautions are essential for the first 14 days.

I suggest that the time has come to question whether this is the most sensible regimen to adopt. Three years ago I took part in a clinical trial of a pill containing 150 μ g of D-norgestrel combined with 30 μ g of ethinyl oestradiol (Ovranette). Patients were instructed to start taking the pill on the first day of the next menstrual period and told that the contraceptive effect was immediate, so that no additional contraception was necessary. No pregnancies occurred in the first cycle among 178 patients for whom the pill was prescribed in this way. Since then it has been my practice to advise patients starting oral contraception to follow this regimen and not one has become pregnant. The first cycle on the pill is shorter (22–23 days), but provided patients are warned about this they do not mind. The incidence of spotting and breakthrough bleeding is not increased. Subsequent packets of pills are started as usual after the standard number of pill-free days and normal cycle length is re-established.

It is also routine practice to advise patients to use additional contraceptive precautions for the first 14 days when changing to a new pill, particularly to one of a lower dose. This advice was reiterated in a recent circular from one of the drug firms, Schering. However, we find that many of our patients object so strongly to this that they refuse to change over to a pill containing 30 μ g of oestrogen, even when we explain to them that such pills are now to be preferred. I believe that the need for additional contraceptive precautions under these circumstances could safely be avoided if patients were advised to start their new pill on the first day of withdrawal bleeding or even proceed to take it on the day after finishing their present preparation—that is, to take two packets of pills consecutively without a break. I have used both these regimens successfully and find that they prove effective and acceptable to the majority of women.

At present patients attending family planning clinics who require additional contra-

ceptives, usually condoms and spermicides, are provided with these free of charge. If we could avoid the need to use two forms of contraception at the same time this would result in a considerable saving of money to the NHS, but above all it would relieve couples from the nuisance and inconvenience of having to use mechanical contraceptives, which so many of them dislike.

NANCY LOUDON

Lothian Health Board,
Family Planning Services,
Edinburgh

Geriatrics in isolation

SIR,—It is difficult to share Dr W B Wright's view (30 July, p 315) that the confinement of doctors exclusively to an older group of patients would not provide the incentive for recruitment. Clearly there is no lack of incentive for recruitment of doctors to provide care exclusively to children (paediatrics). Some regions have already tried to make combined appointments, with some sessions in general medicine and most sessions devoted to geriatric medicine; this does not seem to have improved the situation significantly. It does not appear that Dr W E Bagnall and his colleagues (9 July, p 102) were advocating that the department of geriatric medicine should work "in isolation." It seems that they are able to provide a comprehensive service which is not in isolation. This can be done with the facilities, including the siting of most geriatric beds in three general hospitals where all the work in "acute" specialties are carried out, and adequate staffing, including 13 full-time doctors for 433 beds, which are available to them. In addition they seem to have good understanding and close co-operation from other clinical colleagues and administrators working in the same hospitals. There are also other units up and down the country which are providing efficient geriatric services and which have very similar facilities and staffing. With the redistribution of existing resources in the district, as recommended in the consultative document in 1976,¹ the difficulties in recruitment could be overcome and consequently it should be possible to provide an efficient geriatric service in all districts.

Many clinicians view geriatric medicine as an "unattractive" specialty mainly because of lack of knowledge of the "content" of geriatric medicine and very often junior doctors are dissuaded from taking up geriatrics by their seniors. On the other hand most geriatricians find the job to be busy, interesting, challenging, and rewarding. The other main factors causing difficulties in recruitment are lack of beds in general hospitals and the fact that in some districts there are still 200 or more beds allocated to one consultant geriatrician, while the level of junior medical staff available to him may be unacceptably low. In 1948 there were four consultant geriatricians and by 1973 more than 300.² In 1975 about 10% of geriatric consultant posts in England and Wales were unfilled, but this partly reflects expansion of the specialty. Although recruitment has improved within the past few years, speedy expansion itself has contributed to a significant number of unfilled posts. In comparison with 25-30 years ago much has already been achieved and the system of geriatric care which has been developed in Great Britain is acknowledged as among the best in the world. The number of people over 75 is expected to rise by half a million over 10 years. It is obvious that the care of the elderly and recruitment to the specialty have to be improved. It will be unfortunate if we have to throw away what has been achieved within the last quarter of a century.

Elderly people have a special need. This can be met satisfactorily by the medical and paramedical staff who have developed expertise by working in departments of geriatric medicine over many years. This cannot be abandoned simply because some would like to think of this as being an "unattractive" specialty despite the fact that the number of doctors, juniors and seniors, working in this specialty has increased considerably over the past few years.

N K CHAKRAVORTY

St Luke's Hospital,
Huddersfield

¹ Department of Health and Social Security, *Priorities for Health and Personal Social Services in England*. London, HMSO, 1976.

² Working Party of the Royal College of Physicians, *Lancet*, 1977, **1**, 1092.

Erythromycin in renal failure

SIR,—I was very interested in Dr Paul Sharpstone's paper "Prescribing for patients with renal failure" (2 July, p 36), but was surprised at the omission of any mention of erythromycin. May I be permitted to supply the missing information?

There is no need to alter dosage regimens of erythromycin in any degree of impairment of renal function.¹ Erythromycin is mainly excreted in the liver,³ with some enterohepatic circulation⁴ and elimination in the faeces.⁵ Only 2-5% of orally administered erythromycin appears in the urine, or up to 15% after intravenous administration.^{1,6} Neither peritoneal dialysis nor haemodialysis makes any significant alteration in serum levels of the drug.¹

M ST G WHEELEY

Medical Adviser,
Abbot Laboratories Ltd

Queensborough, Kent

¹ *The Pharmacological Basis of Therapeutics*, ed L S Goodman and A Gilman, 5th edn, p 1225. New York, Macmillan, 1970.

² Bennett, W M, et al, *Journal of the American Medical Association*, 1974, **230**, 1554.

³ *The Pharmacological Basis of Therapeutics*, ed L S Goodman and A Gilman, 5th edn, p 1108. New York, Macmillan, 1970.

⁴ Hammond, J B, and Griffith, R S, *Clinical Pharmacology and Therapeutics*, 1961, **2**, 308.

⁵ Heilman, F R, et al, *Proceedings of the Mayo Clinic*, 1952, **27**, 285.

⁶ Martindale: *The Extra Pharmacopoeia*, ed A Wade, 27th edn, p 1131. London, Pharmaceutical Press, 1977.

Prevention and health

SIR,—Professor I M Richardson (9 July, p 128) calls for "a more clearly identifiable preventive responsibility." In many valuable contributions that he has made as a member of the Council for the Education and Training of Health Visitors he has been fully supported when he has raised the issue of renewed emphasis in the field of prevention—in this instance in relation to the development of health visiting. Recently this statutory body, which of course has responsibility for health visitor education and training, has been encouraged by the many reports in which the role of the health visitor has been acknowledged and the request made for more health visitors, a service running at the moment possibly at about half strength. At the same time the council is gravely concerned that recruitment to health visitor courses fell in the United Kingdom as a whole by 5.8% in 1976,

and preliminary returns suggest that it may fall again by upwards of 10% in 1977.

The Government has suggested a move towards resources for the preventive and community services.¹ Professor Thomas McKeown has pointed out to doctors that their role must be to educate, inform, advise, stimulate, and direct patients towards the basic rules of good health.² The CETHV for its part is ready and willing to continue to develop health visitor courses. When, how, and by whom will the shift of emphasis to the preventive service be begun?

NORMA BATLEY

Director,
Council for the Education and
Training of Health Visitors

London NW1

¹ Department of Health and Social Security, *Prevention and Health: Everybody's Business*. London, HMSO, 1976.

² McKeown, T, *The Role of Medicine: Dream, Mirage or Nemesis?* Rock Carling lecture 1976. London, Nuffield Provincial Hospitals Trust, 1976.

Maintenance digoxin after an episode of heart failure

SIR,—Concerning the comments of Drs S J Warrington and N A J Hamer (23 July, p 265) on our paper (19 March, p 749), our subsequent letter (21 May, p 1350) re-emphasises that all 13 patients with atrial fibrillation had impaired atrioventricular conduction ("slow fibrillators") and gives the numbers of these among those who deteriorated clinically on placebo (4/16) and those who did not (9/30). There was no significant difference in composition between the two groups in this respect ($P=0.26$, Fisher's exact test).

"Rapid fibrillators" were deliberately excluded from the study to avoid the confounding influence of controlling rate.¹ Historically, none of the 13 fibrillators had a ventricular rate exceeding 120/min and on placebo in only one of these did it exceed 100/min (110/min). It is dangerous to assume that only "slow fibrillators" are at risk of bradycardia on digoxin. Slowing of ventricular rate in atrial fibrillation is dependent on change in serum digoxin concentration, whether the untreated rate was "rapid" or "slow."² Since by definition this rate is lower in the "slow fibrillators" we recommend a lower dose for such patients, in whom the only indication for digoxin is heart failure, than that required in "rapid fibrillators" for control of ventricular rate.

Contrary to the implications of Drs Warrington and Hamer, had fewer patients with cor pulmonale presented during the study period the chest x-ray taken during the placebo phase might have shown frank pulmonary oedema in more cases. Similarly, the inclusion of patients with severe lung disease, whose respiratory function tests are more likely to exhibit irreversible changes, would militate against finding a significant difference in the within-patient comparison of spirometric values on digoxin and placebo. They state that the spirometric values were especially low in those who deteriorated clinically on placebo, but in fact there was no significant difference between the measurements on digoxin in this group and those who did not deteriorate clinically (FEV_1 , $P>0.1$; FVC , $P>0.2$; FEV_1/FVC , $P>0.1$; unpaired t -test).

Measurements of systolic time intervals were made under standard conditions. They were recorded at 10 am, after the patient's