CORRESPONDENCE

Medical hazards of air travel I D Rennie, MD..... 515	Pinning down the diagnosis in breast cancer W W Park, MD..... 519	Erythromycin in renal failure M St G Wheeley, MB..... 522
Epilepsy and drowning in childhood S Livingston, MD, and others..... 515	Which arm squeezes the bag? H G Magill, MB..... 519	Prevention and health Norma B Batley, SRN..... 522
A case of intrinsic asthma I W B Grant, FRCPED; D Honeybourne, MRCP..... 516	Photo-onycholysis caused by tetracyclines H Baker, FRCP..... 519	Maintenance digoxin after an episode of heart failure R J Dobbs, MRCP, and others..... 522
Accident or suicide? H H Pilling, MB..... 516	Barium meal in uncomplicated dyspepsia J R W Lyall, MRCP, and T E T West, MD... 520	Sterilisation reversal requests Marion Gillett, MB..... 523
Induction of labour and perinatal mortality R W Beard, FRCOG, and P J Steer, MB..... 516	Chloroquine-induced involuntary movements S Singhi, MD, and others..... 520	Genital yeast infections A B Maclean, MB..... 523
Rubella vaccination during oral contraceptive care A J Rose, MRCGP, and K F Mole, MRCGP.... 517	Strawberry pickers' foot drop P A M Weston, FRCS, and M L E Espir, FRCP 520	Poisoning with antidepressants C L Brewer, MRCPsych..... 523
Late infection after total hip replacement J D Langdon, MB, FDSRCS; M Webley, MRCP, and M L Snaith, MD..... 517	Paternity testing A Usher, FRCPATH..... 520	How effective is measles immunisation? W M Jordan, BM..... 523
Comparison of treatment with fast neutrons and photons Mary Catterall, FRCR, and others..... 518	Drug metabolism in the elderly D J Smithard, MRCP, and M J S Langman, FRCP..... 520	Postgraduate training in environmental health A D C S Cameron, MFCM..... 524
Evolution of poliovirus since introduction of attenuated vaccine H V Wyatt, PhD..... 518	The cowardice continues J E Caughey, FRCP..... 521	Need part time be second-rate? J M Kerry Bluglass, MRCPsych, and R S Bluglass, FRCPsych..... 524
Running away from accidents P A Lawrence, MB..... 518	Prolactin in anorexia nervosa A J Isaacs, MRCP, and others..... 521	Health centre running costs A Elliott, FRCP..... 524
Immersion and drowning in children C D Auld, MB, and J N Norman, FRCSed... 519	Starting on the pill Nancy B Loudon, MB..... 521	How many medical students? T D Brogan, FRCPATH..... 524
Entry to medicine Anne-Marie Feeley; B A Bell, FRCSed..... 519	Geriatrics in isolation N K Chakravorty, MRCPED..... 522	A gross exaggeration T J Millin, FRCS..... 524

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Medical hazards of air travel

SIR,—Like Mr J G Callanan (4 June, p 1473) I seem to have an Ancient Mariner effect in jumbo jets. In December 1975, travelling from Chicago to San Francisco, I spent an hour or so looking after a middle-aged man with chest pains while fuel was jettisoned for an unscheduled stop in Denver to ship him off to hospital. In March 1976, flying directly from Honolulu to Chicago, I spent a yet more energetic two hours looking after a 60-year-old woman with severe angina and tachycardia. She told me (but had not previously told the airline) that she had just taken her discharge from hospital against medical advice and was flying to the mid-West for cardiac investigation. Once again an unscheduled stop, this time in San Francisco. Getting a history was difficult—she had a recurrent laryngeal palsy after thyroidectomy. Six weeks after this I went to O'Hare to meet a British physician arriving from London. Meet him I did: he was returning from a nearby hospital whither he had accompanied the body of the co-pilot, he having spent two hours performing cardiac massage en route.

Several points arise. Firstly, I would endorse Dr Callanan's suggestion that we start to put together a screening system for high-risk passengers, such as the second patient above, but the co-pilot had presumably been thoroughly screened by his employers already, so the chances of success here are small. Secondly, it seems to me that no one can call himself a physician who is not prepared to help in such an emergency (which is terrifying to patient, fellow passengers, and crew), though legally he does not have to and is indeed advised not to. The doctor's mere presence and

authority have a dramatic calming effect on all. As important as this is the attitude of the airlines. My inquiries elicited the fact that such incidents were vanishingly rare (one airline physician told me, "one death for every three million passengers": I wonder whether this statistic is correct). On none of these occasions was any medical equipment, diagnostic or therapeutic, available apart from oxygen and a trivial first-aid box. I was told by one airline that no general agreement had been reached about adequate medical equipment largely because the airlines were afraid that a physician-volunteer would not be able to use the equipment if it was presented to him and there would be important legal implications if the volunteer could not cope. They gave as an example an ophthalmologist suddenly faced with a tracheostomy. This is a silly argument and of small comfort to the reasonably competent physician who would like to pass a laryngeal tube, relieve pain, or even use a stethoscope.

On the second of the two incidents mentioned above passengers were asked if they had any drugs which might be suitable. Trinitrin was produced by a tourist and it relieved the patient's pain. A very large amount of diazepam was also collected from numerous passengers; presumably it would have effectively sedated me. It certainly gave me food for sociological thought.

It was pointed out to me that it was hard for the airlines to identify any volunteer as indeed being a physician (I was able to give numerous evidences of this in the shape of medical licences and so on and would not have thought it hard for others to do so). They also pointed

out the problem of carrying any form of drugs around on planes, especially those making international flights, partly because of the threat of theft. This can scarcely apply to such drugs as trinitrin. It would not be difficult to devise a system for securing an adequately stocked drug and instrument chest. Certainly the doctor flying as a paying passenger cannot afford the weight or space to carry his own emergency bag: such threatening instruments would be confiscated at the airport and the doctor probably imprisoned while the authorities decided whether to press drug charges in addition.

Clearly the airlines and regulatory agencies should get together soon to sort out these problems.

I DRUMMOND RENNIE

Section of Nephrology,
Rush-Presbyterian-St Lukes
Medical Center,
Chicago, Illinois

Epilepsy and drowning in childhood

SIR,—Unfortunately, many physicians still restrict the participation of epileptic children in physical activities, particularly body contact sports and swimming. We were therefore very pleased to note that analysis of the data relative to water immersion accidents in 149 cases led Dr J H Pearn (11 June, p 1510) to conclude that "with effective supervision . . . epileptic children may swim with confidence." Our experience over 41 years in managing approximately 20 000 epileptic children, thousands of whom have engaged in swimming under supervised circumstances, supports his findings. Although some of our patients experienced a convulsion while swimming, none, as in Dr Pearn's series, drowned.

Our experience is also consistent with Dr Pearn's finding that "seizures occurring in the bathtub . . . may be life-threatening," and in