with schistosomal ova is an essential part of the syndrome and that the immunological response a consequence of the infestation and not the cause of the neurological lesion.

In 41 cases of schistosomal spinal cord disease I have traced in the literature¹⁻⁵ the neurological deficit was caused by cord compression due to a discrete granuloma in 13. It would be superfluous to implicate an immunological cause in these cases. Of the remaining 28 cases, in which cord compression was not reported, the spinal cord was examined histologically in 18 and ova found in all of these. I can find no recorded case of this condition in which ova were looked for in the spinal cord and not found. This is in remarkable contrast to schistosomiasis without clinical spinal cord involvement, in which ova are rarely found in the cord.⁶

As ova appear to be invariably present in the affected part of the spinal cord in these cases it is difficult to avoid the conclusion that the lesion is a direct consequence of their presence and there is no need to postulate an immunological cause.

JOHN SIDDORN

School of Medicine,
University of Zambia,
Lusaka, Zambia

¹ Marcial-Rojas, R A, and Fiol, R E, *Annals of Internal Medicine*, 1963, **59**, 215.

² Bird, A V, *Neurology*, 1964, **10**, 647.

³ Levy, L F, and Taube, E, *Central African Medical Journal*, 1969, **15**, 52.

⁴ Rosenbaum, R M, et al, *American Journal of Tropical Medicine and Hygiene*, 1972, **21**, 182.

⁵ Herskowitz, A, *Journal of Neurosurgery*, 1972, **36**, 495.

⁶ Gelfand, M, *Schistosomiasis in Southern Africa*. Juta, Cape Town, 1950.

Poisoning with mianserin

SIR,—I would like to make the following point regarding the letter from Dr P Crome and Miss Belinda Newman (23 July, p 260) concerning overdosage with maprotiline and mianserin.

They report the death of a 72-year-old woman who took an overdose of 60 × 10 mg mianserin and an unknown quantity of lorazepam. On admission her plasma level of mianserin was 110 µg/l and that of lorazepam 500 µg/l. While the level of mianserin was in the therapeutic range, that of lorazepam was in overdose.

W L SHAW
Medical Adviser,
Organon Laboratories Ltd

Morden, Surrey

Immunisation rates in Kent

SIR,—I was concerned to hear an extract from the BMA conference being reported on the national news in which Dr W J Appleyard quoted figures for the uptake of childhood immunisation in Kent. He stated that only 67% of children were being immunised against polio, diphtheria, and tetanus and only a third were being immunised against whooping cough.

The immunisation programme in Kent is computerised and I believe the computer's results to be suspect. I recently received a print-out showing that 46 children on my list had not been immunised. When these children were followed up it was found that six were either not on my list, had come on to it after their immunisations were complete, or mainly lived abroad. Twenty-three had been

fully immunised and in the remaining 17 cases the parents had elected not to have the whooping-cough component of the programme. Thus all the relevant children were accounted for. Discussion with several other GPs in south-east Kent revealed a similar picture. I reported these findings to the area health authority, but I believe that the quoted figures are still a gross underestimate of the true immunisation state of the county, which can be checked only by using the clinic and surgery records rather than relying on computer returns.

A R CRAWFORD

Tenterden, Kent

Danger of saline emetics

SIR,—The *Daily Telegraph* of 2 July drew attention to the dangers inherent in the use of salt and water as an emetic. This specially arose in relation to the unfortunate death of a patient given such an emetic following an overdose of drugs. The news item stated that death was attributed to a dangerous surfeit of sodium in the patient's body which caused a brain haemorrhage. It appeared from the evidence given at the inquest that there is a general unawareness of the danger of salt emetics.

Some 14 years ago I described and discussed this hazard in a case study of fatal hypernatraemia after a saline emetic.¹ The practice of attempting to induce vomiting by this method has become especially dangerous since the advent of the phenothiazines as this group of psychotropic drugs is so frequently prescribed and so frequently figures in overdoses, and many of them are potent antiemetics. Perphenazine, for instance, is 16.6 times more powerful than chlorpromazine as an antiemetic. Nor is it always known with patients who have taken drug overdoses exactly what drugs or quantities have been taken.

All things considered it is probably safest to discard this practice from our therapeutic armamentarium.

DERMOT J WARD

St Loman's Hospital,
Dublin

¹ Ward, D J, *British Medical Journal*, 1963, **2**, 432.

Dextrostix-Eyetone in the insulin hypoglycaemia test

SIR,—Dr M A Preece and Mr R G Newall (16 July, p 152) draw attention to the need for rapidly available blood glucose values during the insulin tolerance test (ITT) to avoid severe hypoglycaemia and to ensure a sufficiently low blood glucose concentration to elicit a growth hormone response. The Dextrostix-Eyetone procedure they describe goes a long way, in principle, to meet this need, but our own experience in routine rather than "research" conditions makes us urge caution.

We would make the following points: (1) Results improved with experience, even in the hands of one enthusiastic clinician. In practice, however, several of our housemen have had too little experience and time to develop the needed expertise. The very simplicity of the system encourages carelessness. (2) No quality control of the procedure is described. This is surely unacceptable for so important a test; the manufacturers themselves recommend the

regular use of Lab-Chek control materials. (3) Even under the favourable conditions of this evaluation the quoted precision is unimpressive. The error, so far as we can calculate from the figures quoted, is twice that of our routine laboratory method. (4) The major potential advantage, avoiding severe hypoglycaemia, is undermined by the one large, inexplicable error which the authors are honest enough to admit.

The Dextrostix-Eyetone system is tantalisingly close to being a major advance in monitoring of ITTs. We would stress the importance of adequate training and experience of the operator and regular quality control procedures.

A M BOLD
S E EVANS
K W GARVEY

Clinical Chemistry Department,
Queen Elizabeth Hospital,
Birmingham

Getting close to the patient

SIR,—With reference to the letter from Dr J L Ogle (16 July, p 191), surely we require more information before we consider altering the system? It is my experience that the sort of advice which he quotes and ridicules, even when issued by an area health authority, has usually been prepared by or after consultation with practitioners in the relevant specialty. This practice is certainly neither new nor the creation of the reorganised Health Service.

If such advice is given to hospital staff which is not the result of consultation it should, of course, be repudiated. If staff are asked to participate in the preparation of such advice they should formally decline to do so if they consider the task inappropriate and make it known to the members of the health authority that the issuing of such advice is unnecessary.

NORMAN MILLS

Ebbw Vale, Gwent

Debendox overdosage in children

SIR,—Following a previous report on Debendox overdosage in children¹ we would like to report a case treated here recently. It illustrates the dangers inherent in accidental ingestion of slow-release preparations when there is ignorance of their nature on the part of those for whom the drug is prescribed, particularly when the drug is commonly prescribed to an age group which is likely to have young children in the home, such as mothers suffering from vomiting of pregnancy.

A 3½-year-old boy was admitted from the accident and emergency department some time after midnight, having apparently been found by his mother 7 h earlier with an empty bottle which had contained an unknown quantity of Debendox tablets, each containing dicyclomine hydrochloride 10 mg, doxylamine succinate 10 mg, and pyridoxine hydrochloride 10 mg in slow-release form (57 were thought later to be unaccounted for). Because he seemed well at the time he had been put to bed and it was not until midnight that he became agitated and restless, precipitating his admission.

On arrival he was hallucinating and restless, with dilated pupils, a dry mouth, hot dry skin, and a tachycardia of 150/min. He was initially treated with pilocarpine 3 mg subcutaneously plus chlorpromazine 15 mg intramuscularly, with the oral admission of magnesium hydroxide as a purgative. He seemed to improve 90 min later after a further