the family they report since all heterozygotes will carry the same mutant gene.

J V LEONARD
O H WOLFF

Department of Child Health,
Institute of Child Health,
London WC1

JUNE LLOYD

Department of Paediatrics,
St George's Hospital,
London SW17

ANDREW WHITELAW

Northwick Park Hospital,
Harrow, Middx

JOAN SLACK

MRC Clinical Genetics Unit,
Institute of Child Health,
London WC1

¹ Betteridge, D J, Higgins, M J P, and Galton, D J,
British Medical Journal, 1975, 4, 500.

More compassion needed

SIR,—Regarding Dr R M Whittington's article on dextropropoxyphene overdosage (16 July, p 172) we had a tragic incident in our family when our daughter was found dead within a period of 2½ hours after ingestion of some 50 Distalgic tablets, source unknown. The post-mortem findings showed collapse at the bases of both lungs. She was found with her hand on the receiver of the telephone, suggesting some sudden distress. This ingestion was possibly a *cri de coeur* following refusal of the staff of a psychiatric teaching hospital to see her, an application she often made. Indeed, after a termination of pregnancy in January all they would offer her was a fortnightly hour with the community nurse, although this was increased when her behaviour improved.

I am writing this because behind so many poisonings lies mental instability of one kind or another, cause usually unknown. Treatment even in this day and age is often sophisticatedly cruel—indeed, if the case is deemed incurable the patient is often pushed back on to the general practitioner, who may or may not be sympathetic but is always busy. Community care is often only of a voluntary type. In this case a local church is trying to help these people but finding little help and support from the psychiatric hospital.

When choosing medical students it should be born in mind that all doctors, be they consultants or general practitioners, need the quality of compassion as much as three As at A level and that we shall all be incurable one day and shall be at their mercy.

MARY LEVINSON

Richmond, Surrey

Tryptophan and rheumatic diseases

SIR,—In "Any Questions" recently (18 June, p 1590) your expert pointed out that there is no published evidence to suggest that L-tryptophan has an antirheumatic effect. However, it should be noted that almost all antirheumatic drugs possess the common property of displacing L-tryptophan from serum protein binding sites, thus increasing the plasma levels of the free amino-acid.¹⁻³ It is still not clear how, if at all, this displacement and the increased availability of free L-tryptophan are related to the antirheumatic

effect of these drugs. Some workers¹ suggest that the freeing of other substances, especially short-chain peptides, may be of greater significance.

Interestingly, those antirheumatic drugs which do free L-tryptophan from serum proteins appear to possess some antidepressive effect. At least one group of workers⁴ has demonstrated an inverse relationship between the level of free tryptophan in patients treated with three different antirheumatic drugs and their depressive scores on the Hamilton Rating Scale.

We have used L-tryptophan extensively as an antidepressant, and in the course of treating a large number of patients have inevitably used the amino-acid in some persons with rheumatoid arthritis. A few of these have shown a dramatic lessening of rheumatoid symptoms associated with a fall in the erythrocyte sedimentation rate. Subsequent discontinuation of the L-tryptophan in several of these cases has been followed by an increase in joint pain. It would be quite wrong to draw specific conclusions from clinical impressions in such small numbers of patients, but it may be considered desirable that formal clinical testing of the effect of L-tryptophan in rheumatoid arthritis should be undertaken.

ALAN D BROADHURST

West Suffolk Hospital,
Bury St Edmunds, Suffolk

¹ McArthur, J N, *et al*, *British Medical Journal*, 1971, 2, 677.

² Smith, M J H, Dawkins, P D, and McArthur, J N, *Journal of Pharmacy and Pharmacology*, 1971, 23, 451.

³ Smith, H G, and Lakatos, C, *Journal of Pharmacy and Pharmacology*, 1971, 23, 180.

⁴ Aylward, M, and Maddock, J, *Lancet*, 1973, 1, 936.

Shortage of anaesthetists

SIR,—Over the past five years I have been the surgeon attached to a large regional health centre in Stockholm with its base hospital (Karolinska Sjukhus) some five miles (8 km) away. During this time I have performed many operations as outpatient procedures using only the various methods of local anaesthesia. The accompanying table is not a comprehensive list, but rather to give some idea of what can be done.

It seems that the time has come for British surgeons to attempt to break some of the bands that bind them to their anaesthetists. A return to old-fashioned and outdated practices is not suggested but rather that more interest be taken in the modern forms of local anaesthesia which, although widely practised in Scandinavia, seem to have been largely ignored (or suppressed) by their British colleagues.

No special equipment is required; in recommended doses the drugs are perfectly safe, the technique is easily learnt, and the costs are low. Simple tourniquets, a sphygmomanometer, and a variety of syringes and needles are all that is required. The anaesthetic agent I use is prilocaine hydrochloride in concentrations of ½% and 1% both with and without adrenaline. The methods most applicable and readily learnt are:

(1) Local infiltration with ½% solution—this is already widely used and needs no further discussion.

(2) Local nerve or plexus blocks—these can be accomplished with 1% prilocaine, though I invariably use ½% with equally good results and avoid local motor blockade. The method allows

Outpatient surgery under local anaesthesia

Operation	No performed
Varicose vein surgery	422
Removal of joint or tendon ganglion	132
Extirpation of Dupuytren's contracture	47
Tendon sheath splitting	64
Peripheral neurolysis	20
Circumcision	49
Cystourethrosopy (males)	34
Removal of hydrocoele/spermatocele	8
Extirpation of sacral pilonidal sinus	22
Herniorrhaphy	34
Extirpation of axillary sweat glands	28
Reposition of wrist fracture	26
Removal of exostoses	44
Extirpation of larger subcutaneous tumours	83
Vasoresection (allowed only since 1976)	28

2 h of surgery to an isolated limb with or without a bloodless field and is also suitable for groin (hernia) and genital operations. Each type of block is excellently described in Eriksson's *Illustrated Handbook of Local Anaesthesia*.¹

(3) Regional intravenous anaesthesia gives the operator about 20 min and is, for example, suitable for reduction of fracture of the wrist. The technique is also described in Eriksson's handbook.

(4) Finally, topical anaesthesia, which, although widely used for oral surgery, is perhaps underestimated for use in other mucosal areas. The urethra is easily anaesthetised after about 10 min contact with a local anaesthetic gel and will allow painless endoscopic examination of the urethra and urinary bladder even in males.

I have seen thyroidectomies and laminectomies performed under local anaesthesia in Sweden, but do not advocate a total redundancy of anaesthetists. Many procedures can, however, be performed with advantage using the above methods. I prefer my patients to have fasted, but this is not essential in acute situations. Nervous patients can be given 10 mg diazepam intravenously before starting the operation. Before embarking on outpatient surgery agreement with a nearby hospital is obligatory so that rapid admission can be obtained if necessary. Only one of my patients has required acute admission to the Karolinska after surgery and that was a 75-year-old man who developed a clot retention after cystoscopy. Three others have needed overnight observation, one after excision of a pilonidal sinus and one after a vasectomy because of bleeding, while a third patient developed abdominal cramps after an inguinal herniorrhaphy.

British surgeons have too long been pampered and protected by a readily available corps of the best-trained anaesthetists in the world. My advice is to stop belly-aching and be practical. Learn the basic principles of local anaesthesia and, by doing it yourself, allow the highly skilled anaesthetists to employ their talents where most required.

DAVID FREEDMAN

Stockholm

¹ Eriksson, E, (ed), *Illustrated Handbook of Local Anaesthesia*. Copenhagen, Munksgaard, 1969.

Diagnosis in cases of haematuria

SIR,—Mr A G Turner and his colleagues (2 July, p 29), describing a diagnostic service for haematuria, have understandably concentrated on the yield of unsuspected carcinoma. They have, however, failed to mention the possibility of a glomerular site for haematuria despite the fact that over 10% of their cases remained undiagnosed.

We have previously reported¹ cases illustrating the importance of renal biopsy as a diagnostic tool in those cases of haematuria

undiagnosed by cystoscopy and routine radiology. When symptoms are recurrent or there is proteinuria between attacks of haematuria a positive histological diagnosis is more likely and has the advantage of making repetitive radiology or even surgery less likely.

J MICHAEL
N F JONES
D R DAVIES
J R TIGHE

St Thomas's Hospital and
Medical School,
London SE1

¹ Michael, J, *et al*, *British Medical Journal*, 1976, **1**, 686.

Distalgesic

SIR,—An examination of the latest available mortality statistics¹ shows that since 1964 the decreasing availability of the traditional methods of self-poisoning—for example, coal gas—has coincided with an increase in the abuse of other agents such as psychotropics and analgesics (aspirin, paracetamol, etc), the majority of the latter being obtained from retail sources.

The recent article by Dr R M Whittington (16 July, p 172) confirms previous reports² that Distalgesic (paracetamol + dextropropoxyphene) is the most widely prescribed analgesic in the United Kingdom and it is perhaps not surprising that, unfortunately, a small minority should seek to abuse it. However, statistics³ indicate that annually many more deaths result from ingestion of aspirin and paracetamol.

The problem associated with the taking of quantities of alcohol with centrally acting drugs is well known to physicians and we stress this in our literature.⁴ However, the effectiveness and safety of Distalgesic are confirmed by its 13 years of widespread clinical use, which has shown that it does not present a problem when taken as directed.

C J WICKS

Medical Director,
Lilly Industries Ltd

Basingstoke, Hants

¹ Office of Population Censuses and Surveys, *Mortality Statistics 1974*, series DH1, No 1. London, HMSO, 1976.

² Skegg, D C G, Doll, R, and Perry, J, *British Medical Journal*, 1977, **1**, 1561.

³ Office of Population Censuses and Surveys, *Mortality Statistics 1974*, series DH4, No 1. London, HMSO, 1976.

⁴ ABPI Data Sheet Compendium, 1977.

Perforating diverticulitis

SIR,—I am prompted to write by a statement in your recent leading article headed "Perforating diverticulitis" (2 July, p 5). While agreeing that a conservative policy of management is often rewarding in inflammatory diverticular disease and welcoming the re-emergence of the old-fashioned expression "a phlegmon," I cannot accept that the use of intravenous fluids and broad-spectrum antibiotics should be advocated for an appendix abscess.

The management of two patients recently may illustrate the point. The first, a 42-year-old man, had a 3½-year history of a sinus in the right loin. A suspected perinephric abscess had originally been opened, but the drainage site had intermittently discharged. He presented several times with right iliac fossa pain and the discharging sinus, each time managed conservatively with antibiotics. Eventual laparotomy revealed an abscess

cavity around the appendix which had a chronic perforation at its centre, through which threadworms moved freely.

The second case involved a suspected carcinoma of the caecum in a 74-year-old woman. Her story of right iliac fossa pain and a febrile illness of three months before was not forthcoming until after a laparotomy had revealed an abscess cavity in the right iliac fossa. This had formed around a small bone from the flying fox (a species of bat which is a culinary delicacy here), which had perforated the caecum.

I feel that neither of these patients would have been greatly served by a regimen of intravenous fluids and antibiotics and, although not absolutely typical of the case which you presumably had in mind, serve to emphasise that adequate surgical drainage remains the basis of successful treatment of an abscess.

JOHN C NICHOLLS

Victoria Hospital,
Mahe, Seychelles

SIR,—I was interested to read Mr D D Rao's report (16 July, p 192) of a solitary perforated diverticulum of the ascending colon. This is a rare condition which the average surgeon meets on only one or two occasions during his "emergency surgery period." I had the doubtful privilege of extending this period beyond all reasonable lengths and was able, in 1962, to report¹ a personal series of six such cases encountered during a 10-year senior registrar stint. The condition can occur at any age—my patients' ages ranging from 12–62—and is always misdiagnosed as acute appendicitis. On opening the abdomen a further misdiagnosis is nearly always made and a right hemicolectomy performed on the assumption that one is dealing with carcinoma. This is because the ruptured diverticulum is surrounded with a firm inflammatory exudate. I made this mistake in my first case, but in the remaining five was forewarned and was able to resect out the offending diverticulum and remove it.

ROBIN BURKITT

Ashford Hospital,
Ashford, Middx

¹ Burkitt, R T, *Postgraduate Medical Journal*, 1962, **38**, 442.

Percutaneous peritoneal biopsy

SIR,—We were interested to read your leading article (18 June, p 1557) concerning abdominal tuberculosis in Britain as we recently diagnosed tuberculous peritonitis in a woman presenting with ascites by the technique of percutaneous peritoneal biopsy. The ascitic fluid contained 43 g protein/l and moderate numbers of leucocytes, but examinations for mycobacteria on direct smear were repeatedly negative. Percutaneous peritoneal biopsy was performed with an Abram's needle through a small skin incision in the right iliac fossa. Histological examination revealed caseating granulomata and large numbers of acid alcohol-fast bacilli.

It is instructive to compare the findings in this patient with those in patients with tuberculous pleural effusion, in whom pleural biopsy is positive in approximately 80% of cases, although mycobacteria are rarely seen on direct smear on pleural fluid. As far as we are aware, peritoneal biopsy is not commonly practised in Britain, although it is used more extensively in the United States,¹ where the

diagnostic yield in tuberculous peritonitis (with the blunt-ended Cope needle) is as high as 60–90%.²

In the presence of ascites percutaneous peritoneal biopsy is a safe and relatively simple bedside technique; its diagnostic yield is high and we suggest that it may be used to advantage in the investigation of ascites and particularly when tuberculous peritonitis is suspected.

P JENKINS
RICHARD BURDEN
C R ELSTON

City Hospital,
Nottingham

¹ Viranuvatti, V, *et al*, *American Journal of Proctology*, 1966, **17**, 488.

² Levine, H, *Archives of Internal Medicine*, 1967, **120**, 542.

Disulfiram blood levels

SIR,—Drs L Ranek and P B Andreassen (9 July, p 94) claim that "the pharmacokinetics of disulfiram have not been elucidated because there are no sensitive and specific methods of determining disulfiram and its metabolites."

Disulfiram is a very difficult drug to understand, but we have described a method of measuring blood levels clinically.¹ After oral disulfiram therapy levels of 2–8 mg/l were obtained, and immediately after implantation levels of 1–3 mg/l were recorded.³ A final metabolite of disulfiram is carbon disulphide. This gas was measured in the breath of disulfiram-treated patients and levels of 0.1–0.4 mg/m³ were obtained.³

M T MALCOLM

West Cheshire Hospital,
Chester

¹ Porter, G S, and Williams, A E, *Journal of Pharmacy and Pharmacology*, 1972, **24**, suppl, p 144.

² Brown, M, Porter, G S, and Williams, A E, *Journal of Pharmacy and Pharmacology*, 1974, **26**, suppl, p 95.

³ Malcolm, M T, Madden, J S, and Williams, A E, *British Journal of Psychiatry*, 1974, **125**, 485.

Trace elements in intravenous feeding of infants

SIR,—Your leading article "Intravenous feeding in infancy" (11 June, p 1490) quotes Fox and Krasna,¹ supporting the recommendation to supply the need for trace elements by giving "10 ml/kg fresh frozen plasma every week." Fox and Krasna, like many other authors, simply state that they rely on this procedure but do not provide any data to support its adequacy. A simple calculation based on normal plasma levels and on recommended allowances for zinc, for example, easily shows that such a procedure will scarcely change anything and will certainly not supply any substantial amounts of zinc. "Normal" adult human plasma (the source of "fresh frozen plasma") contains 960 ± 120 µg Zn/l.² Infants on parenteral nutrition require 40 µg Zn/kg/day³ or 280 µg Zn/kg/week. Thus fresh frozen plasma in a dose of 10 ml/kg/week will supply at the best 10.8 µg/kg/week—that is, only about 1/30th of the required amount.

Infants on parenteral nutrition are sick babies, not normally thriving ones. Catabolism, wound healing, and catch-up growth enormously increase the demand for zinc. It is therefore obvious that 10 ml plasma/kg/week will in no case, not even in healthy ones, supply