Diabetic feet

SIR,—Your leading article on this subject (5 February, p 338) is misleading. You quote a paper by Goodman *et al*¹ which advises suturing of amputation wounds for infected toes and metatarsals; advises the use of a tourniquet during surgery; advises drainage of wounds with iodoform wicks; and dismisses any correlation between palpable leg pulses and results of treatment as “not relevant” in one short paragraph.

My own experience² and that of others³⁻⁶ strongly suggests that suturing of wounds after local amputation for diabetic infection is completely contraindicated. Although well-localised infections can be treated with antibiotics and bed rest prior to limited local surgery,⁷ acute infections with swelling, redness, and systemic manifestations invariably require an urgent, wide, radical, local debridement as this type of infection tends to spread rapidly and extensively in tissue planes despite local measures and intravenous antibiotics.

Although no longer life-threatening, each of these lesions may lead to loss of a limb unless active local medical and surgical treatment are initiated without delay. Finally, unless all peripheral pulses are present and strong, angiography and arterial reconstruction are an important additional method for improving results in this type of patient.

ADOLF SINGER

Department of Surgery,
Mount Sinai School of Medicine,
New York

¹ Goodman, J, *et al*, *Surgery, Gynecology and Obstetrics*, 1976, **143**, 587.

² Singer, A, *et al*, *Journal of Mount Sinai Hospital*, 1968, **35**, 390.

³ Catterall, R C F, in *Clinical Diabetes*, ed W G Oakley, D A Pyke, and K W Taylor. Oxford, Blackwell, 1968.

⁴ Baddeley, R M, *Surgery Digest*, April 1966, p 29.

⁵ Rosenberg, N, *et al*, *AMA Archives of Surgery*, 1956, **72**, 160.

⁶ Pratt, T C, *Medical Clinics of North America*, 1965, **49**, 987.

⁷ Singer, A, *Archives of Surgery*, 1976, **111**, 964.

Hygiene in NHS hospitals

SIR,—We feel it is necessary to reply to Dr H Gordon and Mr G Aston (11 June, p 1536) since they seem to have misunderstood some points in our previous letter (7 May, p 1215). The standards of hygienic practices we recommend are certainly as high as, if not higher than, those required in other establishments.¹ Generalisations such as “the trouble [health authorities] are in today” are often misleading and unnecessarily harmful to patient morale. In this region hygienic practices in hospital catering establishments have considerably improved in recent years owing to the excellent collaboration between environmental health, catering, and infection control staff at both regional and local levels.

We are surprised that it is considered patronising to suggest that environmental health officers should act in an advisory capacity as this is the role of many specialists in the Health Service, including ourselves.

Approximately 5% of patients admitted to hospital acquire an infection, and food poisoning constitutes a rather small though not unimportant part of these infections. We agree that the views of the environmental health officer must be carefully considered, but statutory powers affecting one area only are likely to exert an undue influence on priorities. Many hospitals have deficient ventilation systems and

inadequate facilities in operating theatres and difficulty in obtaining essential equipment as well as problems of overcrowding and understaffing. Decisions on priorities are often difficult and should be made by staff with an overall knowledge of the service.

Infectious disease units play an important role in preventing the spread of infection in hospital and the community but so far as we are aware are not usually very closely involved in catering hygiene and cannot be blamed for past deficiencies.

G AYLIFFE
B J COLLINS

Regional Health Service Infection
Research Laboratory,
Birmingham

¹ *Hygiene in Catering Departments and Food Service Areas*. Health Service Catering Manual, vol II. London, Catering and Dietetic Branch, DHSS, 1974.

A confidence trick

SIR,—The statistics recently published by the Department of Health and Social Security¹ include the following figures concerning hospital inpatient activity in Great Britain. The numbers of beds are given in thousands.

	1968	1975	Change
<i>Beds (total)</i>			
Average occupied daily	439	382	−13.2%
Deaths and discharges	5829	5994	+3%
<i>Beds (non-psychiatric excluding geriatrics and units for younger disabled)</i>			
Average occupied daily	172	151	−12%
Deaths and discharges	5414	5525	+2%

This is a fine record of making better use of available resources. The NHS year by year needs fewer, not more, beds to do more work. Since 1949 in England alone there are now over 2 000 000 more admissions each year using over 60 000 fewer beds. The closure of pay-beds is no help to the NHS, as Professor Harold Ellis has pointed out (28 May, p 1408). It was extraordinarily misleading for Mrs Barbara Castle to say that the closure of private beds in NHS hospitals would release 4000 beds, “the equivalent of four 1000-bedded district hospitals” as she put it, for “public use.” However, Mrs Castle may be given credit for giving £5m a year to facilitate reduction of waiting lists. These fell sharply by 1972, only to rise again as a result of political interventions causing great loss of morale. Their reduction depends not only on overcoming the persisting bottlenecks but on restoring the splendid hospital staff morale which was so evident in the ‘sixties.

British medicine has made fantastic progress in recent years and there has been steady improvement in our hospital services, but there is still a long way to go. There is probably more administrative and medical professional expertise proportionately in this country than anywhere else in the world. Politicians should concentrate on restoring morale and leave the professionals to finish the job of giving us the Health Service we all wish to achieve.

F AVERY JONES

London W1

¹ Department of Health and Social Security, *Health and Personal Social Services Statistics for England (with summary tables for Great Britain) 1975*. London, HMSO, 1976.

SIR,—Professor Harold Ellis writes (28 May, p 1408) about the sad state of private beds at the Gordon Hospital “gathering dust.” It may be of interest to consider the position of the private wing at University College Hospital. This is—or was—the largest private wing in any of the London teaching hospitals, or indeed in the country. It was built in 1937 by public appeal which enabled the site to be bought and a building containing 75 beds and a suite of three operating theatres to be erected. It is completely separate from the rest of the hospital and has its own kitchens, services, and staff; most of the last will presumably become redundant if closure takes place. Part of the building was also originally constructed as a nurses’ home because of a shortage of this accommodation at that time, but as it became possible to house the nursing staff elsewhere these rooms were later altered and absorbed into the wing, which could then provide 120 rooms for private patients.

In 1948, with the passing of the Health Act, the wing, in common with the rest of UCH, was taken over by the Ministry of Health. But, as Aneurin Bevan had agreed, it remained for the use of those who wished to avail themselves of private paying facilities. In the 1950s the advantage to the hospital and medical school as well as to the NHS patients of “geographical whole-time” practice became apparent and the somewhat limited consulting rooms expanded and new appointees to the consultant staff were encouraged to set up practice in the wing. A number of other consultants left their rooms in the Harley Street area to join this service, which by now provided its own x-ray diagnostic facilities. The use of the wing is not confined to UCH staff but has been open to, and used by, consultants from many other hospitals in the London area.

In the 1960s and early ‘70s 25 rooms were closed because they were used for NHS purposes while reconstruction was going on in the main hospital. Subsequently these rooms were turned over to the district management team offices and the major computer experimental project, freeing space for teaching and other facilities elsewhere in the hospital. In 1974 a further nine rooms were given up to help accommodate the heavy demands made by radiotherapy. In 1977, with the phasing out of the first 1000 private beds, 16 more rooms were shut, leaving a total of 72 rooms out of the original 120. Nearly half the private beds have therefore disappeared without any noticeable benefit to NHS patients.

It is worth emphasising that the principle of “geographical whole-time” was recognised and encouraged by the Royal Commission on Medical Education in the Todd Report¹ and has been policy since at UCH. Leaving aside the fact that it will be more than difficult for a group of some 80 consultants to find comparable accommodation elsewhere, both the hospital and medical school are deeply concerned that closure of the wing (a decision, be it noted, taken without any consultation with the university) will gravely damage both teaching and research activities, since university academic staff make use of the wing and thus are able to augment their departmental funds. In addition to teaching and research the availability of part-time NHS staff for administrative responsibility within both the school and hospital (dean, vice-dean, postgraduate sub-dean, chairmen of Medical and Medical Executive Committees, etc) must be seriously affected. It may also not be sufficiently realised that the part-time consultant staff involved could not, even if they wished to, become full-time NHS employees since neither sessions nor money is available for them.

The aim behind the new Act is allegedly to provide more inpatient accommodation for NHS patients by freeing private beds for them. Purpose-built private rooms in separate blocks are quite unsuitable for this for a number of reasons. They require a much higher ratio of nursing staff to patients than an open ward of