CORRESPONDENCE

Design of motor-cycle crash helmets G Hadfield, FRCS.....	43	Suicide and life insurance B M Barraclough, FRACP, and Daphne M Shepherd, MA.....	46	A confidence trick Sir Francis Avery Jones, FRCP; B J Harries, FRCS, and G L Bunton, FRCS; L S Carstairs, FRCR.....	48
Congenital dislocation of the hip P M Dunn, MD.....	43	Serological test for tuberculosis A C Nicholls, PHD.....	46	Court Report and community medicine Agnes M Gordon, MFCM, and others.....	49
Medical hazards of air travel H C McLaren, FRCOG; F S Preston, MB.....	44	Urinary tract infections B M Buckley, DPHIL; A W Asscher, FRCP.....	47	No room for complacency W H Bond, FRCR, and others.....	49
How effective is measles immunisation? J A Dudgeon, FRCP; L E Hilburg, MD.....	44	When is significant not significant? H A F Dudley, FRCS.....	47	Review Body Report J P Lewin, MB.....	49
Cardiac arrhythmias and epilepsy M Winter, MD.....	45	Role of the occupational health adviser F J Robertson, MRCP; A R Broadbent, MFCM.....	47	One-sided contract N H Barley, MRCP.....	50
Anoxic seizures or epilepsy? J B P Stephenson, MRCP.....	45	SOMA I J T Davies, MRCP.....	47	"Inducement practices" S A Nairn, MB.....	50
Apples and the teeth G B Winter, MB, FDSRCS.....	45	Diabetic feet A Singer, FRCS.....	48	Points from letters Overheated wards (B Gaunt); Relative time (D J Vicary); Purpura in malaria (Colonel E E Vella); Wax in the ear causing cough (J Saperia); Speed limits and the public health (A O Griffiths); Protection for motor- cyclists (V H Needham); "Wait for that bike" (R F Irvine).....	50
Insurance companies' attitude to psychiatric illness D E Yarham.....	45	Hygiene in NHS hospitals G A J Ayliffe, FRCPATH, and B J Collins, AIMLS.....	48		

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Design of motor-cycle crash helmets

SIR,—Reading your leading article on this subject (11 June, p 1491), it would appear that you are advocating the use of the full-face helmet for motor-cyclists. The support for this type of helmet seems to be based on the decreased incidence of facial injury due to the presence of a bar across the lower part of the face. This bar was originally built in to the helmet not to protect the face but to increase the stability of the cranial shell.

As chief honorary medical officer to the British Motor Cycle Racing Club for 30 years, and now chairman of the medical committee of the Auto-Cycle Union, I was involved six years ago in trying to get these helmets banned for racing use, as they already were in two European countries and a State of Australia. My main contention was that the weight of the full-face helmet was almost double that of the jet type helmet and the diameter greater, thereby increasing the forces

transmitted to the cervical spine, particularly rotation. The ban was not imposed for lack of evidence. Since then I have personally treated five motor-cycle racers with neck injuries received while wearing full-face helmets, four of these being fatal, and have knowledge of another fatal neck injury only two weeks ago. I think it is likely that many deaths attributed to head injuries are, in reality, high cervical injuries. A further hazard of these helmets is inhalation of vomit which cannot readily escape.

I feel that overprotection of one part of the body to the detriment of another should be carefully considered before full approval is given to articles of protective clothing for motor-cycles. I will continue to wear a jet type helmet.

GORDON HADFIELD

St Peter's Hospital,
Chertsey

Congenital dislocation of the hip

SIR,—I fear that your leading article (21 May, p 1303) is likely to increase the confusion that exists regarding "clicks" and "clunks" and their significance. You suggest that hips which subluxate at birth are "clicks" and do not require treatment. You support your view by reference to a short abstract, which indeed you reproduce almost verbatim, omitting only the critical finding—namely, that the author reported a 15.9% incidence of hip subluxation among the 866 infants he examined at birth.¹ You also omit the last three lines of the abstract, which comment: "In the discussion it was suggested that this high figure might be caused by misrepresentation of clicks which

are easily elicited in the newborn on manipulating the hips, but which do not necessarily indicate dislocation or subluxation."

My own experience, like that of Barlow² and others, suggests that the incidence of hips which "clunk" or subluxate at birth ("Ortolani positive") in this country is approximately 1.5%. Clicking hips are commonplace and should be ignored. Hips which "clunk" are abnormal³ and in my view should be treated at once. It is true that the majority of such hips become stable without treatment. However, some 10% will progress to full dislocation, while a similar proportion of the hips that become stable will be shallow and/or

dysplastic, and may either dislocate later on or eventually become osteoarthritic. Nor do I know of any way of predicting at birth the outcome in any individual case. Furthermore, once the legs extend after birth simple subluxability may progress to full dislocation with intumed limbus within a matter of weeks.

Limitation of abduction, which is an uncommon sign at birth, becomes the rule rather than the exception by the age of 3 months. This greatly increases the danger of ischaemic necrosis of the femoral head in association with abduction splinting.⁴ The leading article implies that this serious complication may arise, albeit infrequently, from simple treatment in the neonatal period. It is worth noting that all the cases reported by Gore⁵ were treated by plaster cast after the neonatal period, the average age of commencing treatment being 8 months. In contrast, Fredensborg⁶ states that in Malmö between 1956 and 1973 no such complication occurred among 510 infants treated from birth with a von Rosen splint. My own experience over the last 18 years has been the same.

My practice is as follows. All infants are examined by a paediatrician on the first day of life. Suspected hip abnormalities are checked at once by a senior member of the staff. When CDH is present, treatment is commenced at once using the Aberdeen⁷ or von Rosen⁸ splint for 6 to 12 weeks depending on whether the hip is subluxatable or dislocated at birth. All infants are followed up with regular radiological checks until at least 5 years old. With the help of Dr Rosemary Evans and the co-operation of Mr Peter Witheroe I am currently reviewing the last 300 cases. There have been no significant complications, and the end results have been excellent. Among over 20 000 infants screened since 1969 there have been four diagnoses of CDH after the neonatal period, an incidence of 0.2/1000. All these particular infants were screened at birth by junior doctors who had just joined the staff, two of them being temporary locum tenens. It seems very likely

that the hip abnormality was overlooked in these cases.

PETER M DUNN

Southmead Hospital,
Bristol

- ¹ Nelson, M. A. *Journal of Bone and Joint Surgery*, 1966, **48B**, 388.
² Barlow, T. G. *Journal of Bone and Joint Surgery*, 1962, **44B**, 292.
³ Dunn, P. M. *Clinical Orthopaedics*, 1976, **119**, 23.
⁴ Ogden, J. A. *Journal of Bone and Joint Surgery*, 1974, **56A**, 941.
⁵ Gore, D. R. *Journal of Bone and Joint Surgery*, 1974, **56A**, 493.
⁶ Fredensborg, N. *Lancet*, 1974, **2**, 780.
⁷ MacKenzie, I. G. *Journal of Bone and Joint Surgery*, 1972, **54B**, 18.
⁸ Rosen, S. von. *Journal of Bone and Joint Surgery*, 1962, **44B**, 284.

Medical hazards of air travel

SIR,—I write to support Mr J G Callanan (4 June, p 1473) in his comments on the use of travelling doctors. In my regular visits to take part in examinations in our sister university in Salisbury, Rhodesia, and on other occasions I have had to spend a good deal of the night attending to sick travellers. For example, a fishbone in the throat, two unexplained syncope possibly related to the 8000 feet pressure, a coronary thrombosis, and gross emphysema with dyspnoea. On these occasions the equipment produced has been inadequate and twice there was no oxygen in the labelled cylinders.

On uninterrupted flights of up to 10 hours the average age of the modern traveller may be over 60. Although no one would wish to curtail visits of senior citizens to their children or grandchildren overseas, especially in the modern package-trip era, nevertheless it would be of value for the chief steward of an aircraft to know how many passengers might be in need of attention during these long flights. With 365 passengers and certain potential patients it might be possible for airlines to have a free return ticket available for an accredited young doctor, who might be able to arrange his off-duty in North America or Southern Africa. Alternatively, doctors who arrive exhausted after attending to sick travellers might be able to sign a form which would ensure that the airline using his services paid a subscription to a named charity.

Finally, litigation against a doctor: surely the wealthy airlines should be prepared to underwrite any possible damages awarded for "malpractice."

HUGH CAMERON McLAREN

University Department of
Obstetrics and Gynaecology,
Birmingham Maternity Hospital,
Birmingham

SIR,—With reference to the letter from Mr J G Callanan (4 June, p 1473) I feel some comment is called for on a number of the points he makes.

Firstly, responsible airlines do carry out very careful medical clearance of elderly, sick, and infirm passengers before accepting them for travel and in fact require the completion of a medical questionnaire from the passenger's own doctor. These, in British Airways, are personally cleared by our medical officer (passenger services) and his staff at Heathrow. Last year, for instance, he dealt with over 26 000 such cases on our route network. Regarding Dr Callanan's second point on the lack of uniformity on the contents of first-aid kits in IATA airlines, this is, of course, true.

The main problem surrounding this is the great variance in regulations concerning the carriage of drugs and medicines in aircraft between countries, and what is legal in one country can be illegal in another.

Regarding myocardial infarction or other acute emergency, for example, renal colic, we in British Airways carry morphine under strictly controlled conditions set by the Home Office, but not all airlines carry drugs of this kind. We also carry cardiac stimulants for use by any doctor on board. The IATA Medical Advisory Committee has spent a great deal of time in trying to get uniformity in the contents of first-aid kits, and this has not yet been achieved. On international journeys there are differences in nomenclature as well as language problems.

As regards help from a physician on board an aircraft, we in British Airways greatly appreciate the help given by many physicians of many nations to our passengers when a call is put out by the cabin service officer on the public address system, and we are always very grateful for the help they give on these occasions. I know of no occasions where such Good Samaritan acts have led to subsequent litigation as far as treatment on UK aircraft is concerned, but we agree it is a sad reflection of our times that some doctors do not volunteer their services because of this possibility. Regarding ground-to-air medical advice, this, of course, has been practised for many years, and with the introduction of very long-range single side-band radio fitted to most long-range transport aircraft this has enabled the captain of the aircraft to seek medical advice from airline medical departments often thousands of miles away from base.

Finally, I think that doctors should be aware that airline medical departments are only too pleased to help with their elderly or infirm patients and to give advice where necessary. Prior warning is essential so that cabin crew and staff on the ground en route can be adequately briefed on the individual requirements.

F S PRESTON

Principal Medical Officer (Air),
British Airways Medical Service

London (Heathrow) Airport

How effective is measles immunisation?

SIR,—I am glad to see that Dr Christine L Miller (11 June, p 1532) has been able to reassure Drs M D Coulter and B M Jones (21 May, p 1347) that the live measles vaccine used in the MRC vaccine trials has given a high degree of protection of the order of 90% of vaccinated children. This is also in accord with recent evidence from the United States reported by Krugman.¹ However, I do not think that all the queries raised by Drs Coulter and Jones have been answered. They reported that they had observed an excessive number of cases of a measles-like illness in immunised children. It is difficult to believe that doctors with such long experience in general practice could make a misdiagnosis of measles in so many cases, but it is conceivably possible that measles in the immunised or partially immunised child may present unusual features. We have recently had a patient admitted to this hospital with atypical measles in that the admission diagnosis was rubella, but serological tests showed that it was a clear-cut primary infection with measles. An unusual aspect of the case was

that the child had no cough or other respiratory symptoms whatsoever, merely a prolonged and rather unusual rash. However, in the circumstances reported by Drs Coulter and Jones some other explanation should be sought for possible vaccination failures.

Basically there are three possible reasons why this could happen. Firstly, even in the best circumstances when measles vaccine is administered strictly in accordance with the recommended procedures, approximately 2—5% of children given vaccine will not develop a detectable immune response. Therefore these children are at risk and may subsequently develop measles. It is important to realise that "inoculation" of measles vaccine is not synonymous with "immunisation." Several studies, including the MRC 1964 trials with live measles vaccine,² have shown that approximately 8% failed to develop an antibody response. Secondly, measles vaccine is a very labile product, being extremely sensitive to light and to changes in temperature. Unless vaccine is held at the correct temperature as suggested by the manufacturers in their leaflet, both before and especially after reconstitution, then there is a risk of a drop in potency. It is also important to use the manufacturer's diluent for reconstitution of the freeze-dried vaccine. A third possibility, but not very likely in Britain, is that if measles vaccine is administered before the first year of life the take-rate will be considerably less than 100%.

In view of the findings reported by Drs Coulter and Jones it would seem desirable to carry out some virological tests to try and determine whether such cases were in fact measles or resulted from some other viral infection.

J A DUDGEON

Hospital for Sick Children,
Great Ormond Street,
London WC1

¹ Krugman, S. *Pediatrics*, 1977, **90**, 1.

² MRC Measles Vaccination Committee, *British Medical Journal*, 1968, **2**, 449.

SIR,—In response to the letter entitled "How effective is measles immunisation?" (21 May, p 1347) let me present data collected on this side of the Atlantic. In a recent survey¹ antibody was detected in 92% of those vaccinated over 13 months of age, 80% at 12 months, and 67% of infants vaccinated under 1 year of age. In another series² it was reported that 80 million doses of live attenuated measles vaccine have been distributed in the United States since 1963. Fourteen-year follow-up has shown persistence of immunity. However, 15 months³ is now the recommended age for measles vaccination when 95% of children should be protected by such immunisation. If an epidemic occurs, vaccination at 6 months of age is also recommended.²

I hope these findings will help Drs M D Coulter and B M Jones in deciding how effective measles vaccination is.

LELAND E HILBURG

Family Doctor Medical Group,
Vallejo,
California,
USA

¹ Shasby, D M, et al, *New England Journal of Medicine*, 1977, **296**, 585.

² Krugman, S, and Katz, S L, *Journal of the American Medical Association*, 1977, **237**, 2228.

³ Krugman, S, *Medical World News*, 29 November 1976, p 30.