paltry gain. Young doctors working for the deputising services would do well to ponder on the advantages of spending those valuable hours learning to be their own plumber, electrician, motor mechanic, or gardener, since attaining proficiency in these trades will certainly save more money than they are ever likely to earn as doctors' deputies.

Ultimately it is degrading for doctors to accept such conditions, a thought that the indulged and somewhat subsidised doctor, at his safe distance from the firing line of emergency medicine, might care to consider.

P O'DONNELL
D TANT

Luton, Beds

*The Secretary writes: "The Central Advisory Committee, BMA Deputising Services, is responsible for reaching agreement with the operating companies of those services on the charges to subscribing general practitioners. It is not a part of the remit of the CAC to negotiate the pay of deputies employed by the BMA services or to negotiate either charges or rates of pay for any other services. The pay of deputies employed by the BMA services is currently under consideration by the Association's Private Practice Committee."—ED, *BMJ*.

Allocation of additional consultant posts

SIR,—The review of allocation of additional consultant posts by Dr D H Vaughan (24 September, p 842) does not relate to the type of problem existing in the hospital service of the North Tees Health District. In 1974 the balance between the anaesthetic establishment and the surgical work load was radically altered by the opening of a large district general hospital and the retirement within a few years of each other of three general surgeons, the work load of two of whom had been reduced by ill health, and their replacement by three able and active young men. As a result the surgical throughput more than doubled, and, according to figures released by the Cleveland Area Health Authority, exceeded by 50%, the national average per full-time equivalent consultant surgeon. At this time the consultant surgeon:consultant anaesthetist ratio was 3.3:1 and no addition to the senior anaesthetic establishment was contemplated despite repeated strident warnings until one of us was dead from coronary thrombosis and two more had extended periods of sickness due to stress illnesses. After this a medical assistant post was replaced by consultant, giving us a ratio of 2.75:1.

As a result of all this I work nine theatre sessions, three of which are in two separate hospitals in a neighbouring district, take my part on a 1 in 4 emergency rota, administer the anaesthetic service, and run the intensive care unit. The district, area, and regional authorities are now aware of this situation and for the past two years have done their best to help us, but their efforts have always foundered on the Central Manpower Committee's insistence that regionally our establishment exceeds the national norm. Surgeons have tried to be helpful to us, but their problems remain, and their problems are ill people who do not get better and sometimes, if not helped, die; so we continue to overwork. If the service is not

to break down completely in districts like this, and the anaesthetic service does not suffer alone here, some way that short-circuits the present ponderous and unfeeling machinery must be found to relieve local anomalies.

E N S FRY

Department of Anaesthesia,
North Tees General Hospital,
Stockton-on-Tees, Cleveland

Royal Medical Benevolent Fund Christmas Appeal

SIR,—Every year I ask members of the medical profession to help us provide the beneficiaries of the Royal Medical Benevolent Fund with a little extra for Christmas. Each year the profession responds most generously and in 1976 over £16 000 was raised and distributed. I am sure it is unnecessary to remind readers that the Royal Medical Benevolent Fund is a special charity supported by doctors for the benefit of doctors and their dependants.

The fund sends a Christmas gift to every one of its beneficiaries and we know from the letters of thanks we receive how much these gifts mean to them. For many, particularly the children, our help means the difference between another ordinary day and a happy family occasion. Please help us to make this difference.

Contributions may be passed direct to the treasurers or medical representatives of the local Guilds of the Royal Medical Benevolent Fund or sent, marked "Christmas Appeal," to the Director, Royal Medical Benevolent Fund, 24 King's Road, Wimbledon, London SW19 8QN.

T HOLMES SELLORS

London SW19

Points from Letters

Which arm squeezes the bag?

Dr M D EDMONDS (Rugby, Warwicks) writes: The profession is deeply indebted to Dr H G Magill (20 August, p 519) and Mr G E Rose (17 September, p 775) . . . for the essential truth about left-handed pipers. . . . No more will second- and third-generation Caledonian exiles, popping out from BMA conferences in Scotland to arrange for a quick complete kitting out with the authentic clan tartan etc before flying back to the New World, be taken aback by the traditional inquiry of the Scottish tailor: "... and on which side d'ye carry your drones?"

Mr M BECKETT (King's College Hospital Medical School) writes: Mr G E Rose (17 September, p 775), arguing the historical precedent of left-handed pipers, refers to a self-portrait of Joseph MacDonald in 1761 in which he is carrying his pipes in this fashion. However, self-portraits almost invariably show us a mirror image of the artist. Perhaps there is other evidence for his left-handedness?

Death from ischaemic heart disease

Dr G SYMONDS (London NW3) writes: One of the reasons why there appears to be an

increase in the numbers of deaths from heart disease (leading article, 27 August, p 537) may be that no one is allowed to die from natural causes. . . . Since technically everyone dies from heart failure it may be that in many elderly people, in order to expedite the formalities, the cause of death is given as "heart failure as a consequence of arteriosclerosis" when the patient has simply come to the end of his natural life.

Finger clubbing

Dr G H HALL (Exeter) writes: Your leading article (24 September, p 785) mentions that the "mysterious material" invoked as a cause of clubbing by "arteriovenous shunting across the lung" has never been isolated. True, but there is experimental evidence¹⁻³ that it is reduced ferritin and as far as I know this suggestion has neither been superseded nor disproved.

¹ Hall, G H, *Lancet*, 1959, **1**, 750.

² Hall, G H, and Laidlaw, D, *Clinical Science*, 1963, **24**, 121.

³ Hall, G H, and Sachner, M A, *Journal of Applied Physiology*, 1966, **21**, 923.

Dr M HAZELL (Nether Edge Hospital, Sheffield) writes: Your leading article (24 September, p 785), while providing an impressive list of causes of finger clubbing, unfortunately omits congenital clubbing, the diagnosis of which might save the unfortunate sufferer from investigation.

Urinary incontinence

Mr E S GLEN (Urology Unit, Southern General Hospital, Glasgow) writes: . . . It should be emphasised that in cases of urinary incontinence the urologist can be the link between the specialties. In this context it is beneficial for trainees in specialties such as geriatrics, gynaecology, and paediatrics to have exposure to urology units with special experience in incontinence. . . . Mutual benefit accrues, the width of experience being increased by the multidisciplinary approach, and the non-urologist becomes better able to determine when to refer patients to the urologist.

Related ancillary staff

Dr F S BLACK (Audlem, Cheshire) writes: . . . Should not all doctors' wives who work whole or part time in the practice join NUPE or COHSE? They would then be given a label and an analogue with other NHS clerical/nursing workers and perhaps the union would fight for their rights.

The placenta and twins

Dr A J BRADBURY (Birmingham Maternity Hospital) writes: In reply to Dr R B Williams (24 September, p 835) concerning my case report of malaria in one twin (3 September, p 613) I should like to explain that at the time of the acute illness it was clear that the features of the two twins were quite different and the fact that they were non-identical was subsequently confirmed by determination of their blood groups, which were also different.